

# THE KONTSEVICH CONSTANTS FOR THE VOLUME OF THE MODULI OF CURVES AND TOPOLOGICAL RECURSION

KEVIN M. CHAPMAN, MOTOHICO MULASE, AND BRAD SAFNUK

ABSTRACT. We give an Eynard-Orantin type *topological recursion formula* for the canonical Euclidean volume of the combinatorial moduli space of pointed smooth algebraic curves. The recursion comes from the edge removal operation on the space of ribbon graphs. As an application we obtain a new proof of the Kontsevich constants for the ratio of the Euclidean and the symplectic volumes of the moduli space of curves.

## CONTENTS

1. Introduction	1
2. The combinatorial model of the moduli space	5
3. Topological recursion for the number of integral ribbon graphs	7
4. The Laplace transform of the number of integral ribbon graphs	13
5. The Euclidean volume of the moduli space	14
6. The symplectic volume of the moduli space and the Kontsevich constants	16
7. The Eynard-Orantin theory on $\mathbb{P}^1$	20
Appendix A. Calculation of the Laplace transforms	27
Appendix B. Examples	32
References	36

## 1. INTRODUCTION

The purpose of this paper is to identify a combinatorial origin of the *topological recursion formula* of Eynard and Orantin [15] as the operation of edge removal from a ribbon graph. As an application of our formalism, we establish a new proof of the formula for the Kontsevich constants  $\rho = 2^{5g-5+2n}$  of [28, Appendix C].

In moduli theory it often happens that we have two different notions of the *volume* of the moduli space. The volume may be defined by the push-forward measure of the canonical construction of the moduli space. Or it may be defined as the symplectic volume with respect to the intrinsic symplectic structure of the moduli space. An example of such situations is the moduli space of flat  $G$ -bundles on a fixed Riemann surface for a compact Lie group  $G$  [24, 25, 29, 49]. In this case, the two definitions of the volume agree.

The space we study in this paper is the combinatorial model of moduli space  $\mathcal{M}_{g,n}$  of smooth algebraic curves of genus  $g$  with  $n$  distinct marked points. It also has two different families of volumes parametrized by  $n$  positive real parameters. One comes from the push-forward measure, and the other comes from the intrinsic symplectic structure depending on these parameters. And again these two notions of volume agree.

The moduli space  $\mathcal{M}_{g,n}$  admits orbifold cell-decompositions parametrized by the collection of positive real numbers assigned to the marked points. This orbifold is identified as

---

2000 *Mathematics Subject Classification.* Primary: 14N35, 05C30, 53D30, 11P21; Secondary: 81T30.

the space of ribbon graphs of a prescribed perimeter length, using the theory of Strebel differentials. In his seminal paper of 1992, Kontsevich [28] calculated the symplectic volume of orbi-cells, and compared it with the standard Euclidean volume. He found that the ratio was a constant depending only on the genus of the curve and the number of marked points. This constant plays a crucial role in his *main identity*, and hence in his proof of the Witten conjecture. He wrote in Appendix C of [28] that his proof of the evaluation of this constant “presented here is *not nice*, but we don’t know any other proof.” In this article we give another proof of the formula for the Kontsevich constant, based on the topological recursion for ribbon graphs.

The idea of topological recursion has been used as an effective tool for calculating many quantities related to the moduli space  $\mathcal{M}_{g,n}$  and its Deligne-Mumford compactification  $\overline{\mathcal{M}}_{g,n}$ . The quantities we can deal with include tautological intersection numbers and certain Gromov-Witten invariants. Suppose we have a collection of quantities  $v_{g,n}$  for  $g \geq 0$  and  $n > 0$  subject to the *stability* condition  $2g - 2 + n > 0$ , which guarantees the finiteness of the automorphism group of an element of  $\mathcal{M}_{g,n}$ . By an Eynard-Orantin type topological recursion formula [15] we mean a particular inductive formula for  $v_{g,n}$  with respect to the *complexity*  $2g - 2 + n$  of the form

$$(1.1) \quad v_{g,n} = f_1(v_{g,n-1}) + f_2(v_{g-1,n+1}) + \sum_{\substack{\text{stable} \\ g_1+g_2=g \\ n_1+n_2=n-1}} f_3(v_{g_1,n_1+1}, v_{g_2,n_2+1})$$

with linear operators  $f_1, f_2$  and a bilinear operator  $f_3$ , where the sum is taken for all possible partitions of  $g$  and  $n - 1$  subject to the stability conditions  $2g_1 - 1 + n_1 > 0$  and  $2g_2 - 1 + n_2 > 0$ . We refer to Section 7 for more detail.




FIGURE 1.1. The topological recursion. The reduction of  $2g - 2 + n$  by 1 corresponds to cutting off of a pair of pants from an  $n$ -punctured surface.

There are many examples of such formulas.

- (1) The Witten-Kontsevich theory for the tautological cotangent class (i.e. the  $\psi$ -class) intersection numbers

$$(1.2) \quad \langle \tau_{d_1} \cdots \tau_{d_n} \rangle_{g,n} = \int_{\overline{\mathcal{M}}_{g,n}} c_1(\mathbb{L}_1)^{d_1} \cdots c_1(\mathbb{L}_n)^{d_n}$$

on the moduli stack  $\overline{\mathcal{M}}_{g,n}$  of stable algebraic curves of genus  $g$  with  $n$  distinct smooth marked points. The Dijkgraaf-Verlinde-Verlinde formula [9], which is equivalent to the Virasoro constraint condition, is a topological recursion of the form (1.1).

- (2) The Mirzakhani recursion formula for the Weil-Petersson volume of the moduli space of bordered hyperbolic surfaces with prescribed geodesic boundaries [32, 33] is a topological recursion.

- (3) Mixed intersection numbers

$$\langle \tau_{d_1} \cdots \tau_{d_n} \kappa_1^{m_1} \kappa_2^{m_2} \kappa_3^{m_3} \cdots \rangle_{g,n}$$

of  $\psi$ -classes and the Mumford-Morita-Miller  $\kappa$ -classes satisfy a topological recursion, first found in [35] for the case with  $\kappa_1$  and later generalized in [30].

- (4) The expectation values of the product of resolvents of various matrix models satisfy a topological recursion (see for example, [12]). This is the origin of the work [15].
- (5) Indeed, the first three geometric theories turned out to be examples of the general theory [15] of topological recursions [13, 16], though geometric theories had been discovered earlier than the publication of [15].
- (6) Both open and closed Gromov-Witten invariants of an arbitrary toric Calabi-Yau threefold are expected to satisfy a topological recursion. This is the *remodeling* conjecture of [31, 4].
- (7) Simple Hurwitz numbers satisfy a topological recursion. It was first conjectured in [5] based on a limit case of the remodeling conjecture, and was recently proved in [3, 14, 36].
- (8) The simplest case of the remodeling conjecture for  $\mathbb{C}^3$  was proved in [7, 50, 51] based on the Laplace transform technique of [14].
- (9) As shown below, the number of metric ribbon graphs with integer edge lengths for a prescribed boundary condition satisfies a topological recursion.

Our current paper provides an elementary approach to the idea of topological recursion that uniformly explains the combinatorial nature of the geometric examples (1), (2), (3), (7), (8) and (9).

The work of Harer [21], Mumford [37], Strebel [45], Thurston and others [44] show that there is a topological orbifold isomorphism

$$\mathcal{M}_{g,n} \times \mathbb{R}_+^n \cong RG_{g,n},$$

where

$$RG_{g,n} = \coprod_{\substack{\Gamma \text{ ribbon graph} \\ \text{of type } (g,n)}} \frac{\mathbb{R}_+^{e(\Gamma)}}{\text{Aut}(\Gamma)}$$

is the space of metric ribbon graphs of genus  $g$  and  $n$  boundary components, and  $e(\Gamma)$  is the number of edges of a ribbon graph  $\Gamma$ . We denote by  $\pi : RG_{g,n} \rightarrow \mathbb{R}_+^n$  the natural projection, and its fiber at  $\mathbf{p} \in \mathbb{R}_+^n$  by  $RG_{g,n}(\mathbf{p}) = \pi^{-1}(\mathbf{p})$ . To give a combinatorial description of tautological intersection numbers (1.2) on  $\overline{\mathcal{M}}_{g,n}$ , Kontsevich [28, Page 8] introduced a combinatorial symplectic form  $\omega_K(\mathbf{p})$  on  $RG_{g,n}(\mathbf{p}) \cong \mathcal{M}_{g,n}$  and its *symplectic*

volume

$$(1.3) \quad v_{g,n}^S(\mathbf{p}) = \int_{RG_{g,n}(\mathbf{p})} \exp(\omega_K(\mathbf{p})).$$

The definition of this symplectic form is given in Section 6. At each orbi-cell level, the derivative  $d\pi$  of the projection map  $\pi$  is determined by the edge-face incidence matrix

$$A_\Gamma : \mathbb{R}_+^{e(\Gamma)} \longrightarrow \mathbb{R}_+^n$$

of a ribbon graph  $\Gamma$ . Note that we have the standard volume forms  $dl_1 \wedge \cdots \wedge dl_{e(\Gamma)}$  on  $\mathbb{R}_+^{e(\Gamma)}$  and  $dp_1 \wedge \cdots \wedge dp_n$  on  $\mathbb{R}_+^n$ . We can define the Euclidean volume of the inverse image  $P_\Gamma(\mathbf{p}) = A_\Gamma^{-1}(\mathbf{p})$  of  $\mathbf{p} \in \mathbb{R}_+^n$  using the push-forward measure by

$$\text{vol}(P_\Gamma(\mathbf{p})) = \left. \frac{(A_\Gamma)_*(dl_1 \wedge \cdots \wedge dl_{e(\Gamma)})}{dp_1 \wedge \cdots \wedge dp_n} \right|_{\mathbf{p}},$$

where  $(A_\Gamma)_*(dl_1 \wedge \cdots \wedge dl_{e(\Gamma)})$  is the  $n$ -form on  $\mathbb{R}_+^n$  obtained by integrating the volume form on  $\mathbb{R}_+^{e(\Gamma)}$  along the fiber  $A_\Gamma^{-1}(\mathbf{p})$ . The *Euclidean* volume of the moduli space is defined by

$$v_{g,n}^E(\mathbf{p}) = \sum_{\substack{\Gamma \text{ ribbon graph} \\ \text{of type } (g,n)}} \frac{\text{vol}(P_\Gamma(\mathbf{p}))}{|\text{Aut}(\Gamma)|}.$$

In Appendix C of [28], Kontsevich proved the following.

**Theorem 1.1** ([28]). *The ratio of the symplectic volume and the Euclidean volume of  $RG_{g,n}(\mathbf{p})$  is a constant depending only on  $g$  and  $n$ , and its value is*

$$(1.4) \quad \rho = \frac{v_{g,n}^S(\mathbf{p})}{v_{g,n}^E(\mathbf{p})} = 2^{5g-5+2n}.$$

**Remark 1.2.** The Euclidean volume of the polytope

$$P_\Gamma(\mathbf{p}) = \{\mathbf{x} \in \mathbb{R}_+^{e(\Gamma)} \mid A_\Gamma \mathbf{x} = \mathbf{p}\}$$

is a *quasi-polynomial* and is difficult to calculate in general. It is quite surprising that the ratio  $\rho$  of the two functions is indeed a constant. Although he says “not nice,” Kontsevich’s original proof is a beautiful application of homological algebra to the complexes defined by the incidence matrix  $A_\Gamma$ .

The new proof we present here uses an elementary argument on the topological recursion of ribbon graphs corresponding to the edge removal operation. We show that both  $v_{g,n}^S(\mathbf{p})$  and  $2^{5g-5+2n} \cdot v_{g,n}^E(\mathbf{p})$  satisfy exactly the same induction formula based on  $2g - 2 + n$ , *after taking the Laplace transform*. We then calculate the initial condition for the recursion formula, i.e., the cases for  $(g, n) = (0, 3)$  and  $(1, 1)$ , and see that the equality holds. Since the topological recursion uniquely determines the value for every  $(g, n)$  subject to the stability condition  $2g - 2 + n > 0$ , we conclude that

$$v_{g,n}^S(\mathbf{p}) = 2^{5g-5+2n} \cdot v_{g,n}^E(\mathbf{p}).$$

Here the appearance of the Laplace transform is significant. The Laplace transform plays a mysterious as well as a crucial role in each of the works [13, 14, 16, 28, 36, 42]. In the light of the Eynard-Orantin recursion formalism [15] and the remodeling conjecture due to Mariño [31] and Bouchard-Klamm-Mariño-Pasquetti [4], we find that the Laplace transform appearing in these contexts is the *mirror map*. Usually mirror symmetry is considered as

a duality, and hence a family of Fourier-Mukai type transforms naturally appears [23, 46]. In our context, however, the nature of duality is not apparent. On one side of the story (the A-model side) we have a combinatorial structure. The mirror symmetry transforms this combinatorial structure into the world of complex analysis (the B-model side). In the complex analysis side we have such objects as the residue calculus of [15] and integrable nonlinear PDEs such as the KdV equations [28, 30, 35, 48], the KP hierarchy [26, 27, 41], Frobenius manifold structures [10, 11], the Ablowitz-Ladik hierarchy [6], and more general integrable systems considered in [17, 18, 19]. The mathematical apparatus of the mirror map hidden in these structures is indeed the Laplace transform.

This paper is organized as follows. In Section 2 we review ribbon graphs and combinatorial description of the moduli space  $\mathcal{M}_{g,n}$  that are necessary for our investigation. Although the definition of the Euclidean volume of  $RG_{g,n}(\mathbf{p})$  is straightforward, it seems to be difficult to calculate it and there is no concrete formula. The approach we take in this paper is to appeal to the counting of *lattice points* of  $RG_{g,n}(\mathbf{p})$ . Thus Section 3 is devoted to proving an effective topological recursion formula for the number of lattice points in the space of metric ribbon graphs with prescribed perimeters. Our proof is based on counting ciliated ribbon graphs. Once we find the number of lattice points in  $RG_{g,n}(\mathbf{p})$ , we can obtain its volume by taking the limit as the mesh of the lattice tends to 0. To compare the number of lattice points and the volume, the simplest path is to take the Laplace transform. Thus we are led to calculating the Laplace transform of the topological recursion for the number of lattice points in Section 4. After establishing the Laplace transform formula, one can read off the information of the Euclidean volume of  $RG_{g,n}(\mathbf{p})$  as the leading terms of the Laplace transform, by introducing the right coordinate system. This is carried out in Section 5. The Kontsevich symplectic form is defined in Section 6, and the topological recursion for the symplectic volume due to [2] is reviewed. With these preparations, we give a new and simple proof of (1.4). In Section 7 we explain the Eynard-Orantin formalism. This formalism is independent on the context and provides the same formula. We then convert our recursion formulas into this formalism, and observe how they all fit together in a single formula. This is the beauty and strength of the Eynard-Orantin formalism.

We present a full detail of the calculations of the Laplace transform in this paper, hoping it may lead to a deeper understanding of the Eynard-Orantin theory and the mirror map. Appendix A is thus devoted to giving a proof of (4.6) and (6.11). These recursion formulas start with the initial values  $(g, n) = (0, 3)$  and  $(g, n) = (1, 1)$ . The Eynard-Orantin theory also uses the unstable case  $(g, n) = (0, 2)$ . All these values are calculated in Appendix B, together with a few more examples.

**Acknowledgement.** M.M. thanks Soheil Arabshahi, Zainal bin Abdul Aziz, Minji Kim, and Jian Zhou for useful discussions. He is particularly grateful to Michael Pankava and Andy Port for discussions on the Laplace transform formulas. During the preparation of this work, the research of K.C. was supported by NSF grant DMS-0636297, M.M. received support from the American Institute of Mathematics, NSF, Universiti Teknologi Malaysia, and Tsinghua University in Beijing, and the research of B.S. was supported by Central Michigan University.

## 2. THE COMBINATORIAL MODEL OF THE MODULI SPACE

Let us begin with reviewing basic facts about ribbon graphs and the combinatorial model of the moduli space  $\mathcal{M}_{g,n}$  due to Harer [21], Mumford [37], and Strebel [45]. We refer to [34] for precise definitions and more detailed exposition.

A *ribbon graph* of topological type  $(g, n)$  is the 1-skeleton of a cell-decomposition of a closed oriented topological surface  $\Sigma$  of genus  $g$  that decomposes the surface into a disjoint union of  $v$  0-cells,  $e$  1-cells, and  $n$  2-cells. The Euler characteristic of the surface is given by  $2 - 2g = v - e + n$ . The 1-skeleton of a cell-decomposition is a graph  $\Gamma$  drawn on  $\Sigma$ , which consists of  $v$  vertices and  $e$  edges. An edge can form a loop. We denote by  $\Sigma_\Gamma$  the cell-decomposed surface with  $\Gamma$  its 1-skeleton. Alternatively, a ribbon graph can be defined as a graph with a cyclic order given to the incident half-edges at each vertex. By abuse of terminology, we call the boundary of a 2-cell of  $\Sigma_\Gamma$  a *boundary* of  $\Gamma$ , and the 2-cell itself as a *face* of  $\Gamma$ .

A *metric ribbon graph* is a ribbon graph with a positive real number (the length) assigned to each edge. For a given ribbon graph  $\Gamma$  with  $e = e(\Gamma)$  edges, the space of metric ribbon graphs is  $\mathbb{R}_+^{e(\Gamma)}/\text{Aut}(\Gamma)$ , where the automorphism group acts through permutations of edges (see [34, Section 1]). We restrict ourselves to the case that  $\text{Aut}(\Gamma)$  fixes each 2-cell of the cell-decomposition. If we also restrict that every vertex of a ribbon graph has degree (i.e., valence) 3 or more, then using the canonical holomorphic coordinate system of a topological surface [34, Section 4] and the Strebel differentials [45], we obtain an isomorphism of topological orbifolds [21, 37, 44]

$$(2.1) \quad \mathcal{M}_{g,n} \times \mathbb{R}_+^n \cong RG_{g,n}.$$

Here

$$RG_{g,n} = \coprod_{\substack{\Gamma \text{ ribbon graph} \\ \text{of type } (g,n)}} \frac{\mathbb{R}_+^{e(\Gamma)}}{\text{Aut}(\Gamma)}$$

is the orbifold consisting of metric ribbon graphs of a given topological type  $(g, n)$  with degree 3 or more. The degree condition is necessary to bound the number of edges  $e(\Gamma)$  for a given topological type  $(g, n)$ . If we allow degree 2 vertices, then there are infinitely many different ribbon graphs for every  $(g, n)$ . By restricting to ribbon graphs of degree 3 or more, we have the bound  $e(\Gamma) \leq 3(2g - 2 + n)$ , which gives the dimension of each orbi-cell  $\mathbb{R}_+^{e(\Gamma)}/\text{Aut}(\Gamma)$ . The gluing of orbi-cells is done by making the length of a non-loop edge tend to 0. The space  $RG_{g,n}$  is a smooth orbifold (see [34, Section 3], [44]). We denote by  $\pi : RG_{g,n} \rightarrow \mathbb{R}_+^n$  the natural projection via (2.1), which is the assignment of the collection of perimeter length of each boundary to a given metric ribbon graph.

Take a ribbon graph  $\Gamma$ . Since  $\text{Aut}(\Gamma)$  fixes every boundary component of  $\Gamma$ , they can be labeled by  $N = \{1, 2, \dots, n\}$ . For a moment let us give a label to each edge of  $\Gamma$  from an index set  $E = \{1, 2, \dots, e\}$ . The edge-face incidence matrix is defined by

$$(2.2) \quad \begin{aligned} A_\Gamma &= [a_{i\eta}]_{i \in N, \eta \in E}; \\ a_{i\eta} &= \text{the number of times edge } \eta \text{ appears in face } i. \end{aligned}$$

Thus  $a_{i\eta} = 0, 1$ , or  $2$ , and the sum of entries in each column is always 2. The  $\Gamma$  contribution of the space  $\pi^{-1}(p_1, \dots, p_n) = RG_{g,n}(\mathbf{p})$  of metric ribbon graphs with a prescribed perimeter  $\mathbf{p} = (p_1, \dots, p_n)$  is the orbifold polytope

$$P_\Gamma(\mathbf{p})/\text{Aut}(\Gamma), \quad P_\Gamma(\mathbf{p}) = \{\mathbf{x} \in \mathbb{R}_+^e \mid A_\Gamma \mathbf{x} = \mathbf{p}\},$$

where  $\mathbf{x} = (\ell_1, \dots, \ell_e)$  is the collection of edge lengths of a metric ribbon graph  $\Gamma$ . We have

$$(2.3) \quad \sum_{i \in N} p_i = \sum_{i \in N} \sum_{\eta \in E} a_{i\eta} \ell_\eta = 2 \sum_{\eta \in E} \ell_\eta.$$

The canonical Euclidean volume  $\text{vol}(P_\Gamma(\mathbf{p}))$  of the polytope  $P_\Gamma(\mathbf{p})$  is the ratio of the push-forward measure of the Lebesgue measure on  $\mathbb{R}_+^e$  by  $A_\Gamma$  and the Lebesgue measure on  $\mathbb{R}_+^n$  at the point  $\mathbf{p} \in \mathbb{R}_+^n$ :

$$(2.4) \quad \text{vol}(P_\Gamma(\mathbf{p})) = \left. \frac{(A_\Gamma)_*(dl_1 \wedge \cdots \wedge dl_e)}{dp_1 \wedge \cdots \wedge dp_n} \right|_{\mathbf{p}},$$

where  $(A_\Gamma)_*(dl_1 \wedge \cdots \wedge dl_e)$  is the  $n$ -form on  $\mathbb{R}_+^e$  obtained by integrating the volume form on  $\mathbb{R}_+^e$  along the fiber  $\pi^{-1}(\mathbf{p})$ . This definition is equivalent to imposing

$$(2.5) \quad \int_D \text{vol}(P_\Gamma(\mathbf{p})) dp_1 \wedge \cdots \wedge dp_n = \int_{A_\Gamma^{-1}(D)} dl_1 \wedge \cdots \wedge dl_e$$

for every open subset  $D \subset \mathbb{R}_+^n$  with compact closure. We define the *Euclidean volume function* by

$$(2.6) \quad v_{g,n}^E(\mathbf{p}) = v_{g,n}^E(p_1, \dots, p_n) = \sum_{\substack{\Gamma \text{ trivalent ribbon} \\ \text{graph of type } (g,n)}} \frac{\text{vol}(P_\Gamma(\mathbf{p}))}{|\text{Aut}(\Gamma)|}.$$

This is the Euclidean volume of the moduli space  $\mathcal{M}_{g,n}$  considered as the orbi-cell complex

$$(2.7) \quad RG_{g,n}(\mathbf{p}) \stackrel{\text{def}}{=} \pi^{-1}(\mathbf{p}) = \coprod_{\substack{\Gamma \text{ ribbon graph} \\ \text{of type } (g,n)}} \frac{P_\Gamma(\mathbf{p})}{\text{Aut}(\Gamma)} \cong \mathcal{M}_{g,n}$$

with the prescribed perimeter length  $\mathbf{p} \in \mathbb{R}_+^n$ . Only degree 3 (or trivalent) graphs contribute to the volume function because they parametrize the top dimensional cells. Since  $\dim_{\mathbb{R}} RG_{g,n}(\mathbf{p}) = 2(3g - 3 + n)$ , we expect that the definition of the push-forward measure and the relation (2.5) imply that the volume function  $v_{g,n}^E(\mathbf{p})$  has the polynomial growth of order  $2(3g - 3 + n)$  as  $\mathbf{p} \rightarrow \infty$ . We will verify this growth order in Section 5, (5.3).

### 3. TOPOLOGICAL RECURSION FOR THE NUMBER OF INTEGRAL RIBBON GRAPHS

It is a difficult task to find a topological recursion formula for the Euclidean volume functions  $v_{g,n}^E(\mathbf{p})$  directly from its definition. Although it is somewhat a detour, we first derive a recursion formula for the number of metric ribbon graphs with integer edge lengths, take its Laplace transform, and then extract the topological recursion for the Euclidean volume functions.

Thus our main subject of this section is the set of all metric ribbon graphs  $RG_{g,n}^{\mathbb{Z}_+}$  whose edges have integer lengths. We call such a ribbon graph an *integral ribbon graph*. Let us begin with defining the weighted number  $|RG_{g,n}^{\mathbb{Z}_+}(\mathbf{p})|$  of integral ribbon graphs with prescribed perimeter lengths  $\mathbf{p} \in \mathbb{Z}_+^n$ :

$$(3.1) \quad N_{g,n}(\mathbf{p}) = |RG_{g,n}^{\mathbb{Z}_+}(\mathbf{p})| = \sum_{\substack{\Gamma \text{ ribbon graph} \\ \text{of type } (g,n)}} \frac{|\{\mathbf{x} \in \mathbb{Z}_+^{e(\Gamma)} \mid A_\Gamma \mathbf{x} = \mathbf{p}\}|}{|\text{Aut}(\Gamma)|}.$$

Since the finite set  $\{\mathbf{x} \in \mathbb{Z}_+^{e(\Gamma)} \mid A_\Gamma \mathbf{x} = \mathbf{p}\}$  is a collection of lattice points in the polytope  $P_\Gamma(\mathbf{p})$  with respect to the canonical integral structure  $\mathbb{Z} \subset \mathbb{R}$  of the real numbers,  $N_{g,n}(\mathbf{p})$  can be thought of as counting the number of *lattice points* in  $RG_{g,n}(\mathbf{p})$  with a weight factor  $1/|\text{Aut}(\Gamma)|$  for each ribbon graph. The function  $N_{g,n}(\mathbf{p})$  is a symmetric function in  $\mathbf{p} = (p_1, \dots, p_n)$  because the summation runs over all ribbon graphs of topological type  $(g, n)$  whose boundaries are labeled by the index set  $N$ .

**Remark 3.1.** The function (3.1) was first considered in [38]. Note that we do not allow the integer vector  $\mathbf{p} \in \mathbb{Z}_+^n$  to have any 0 entry, since each face of a ribbon graph must have a positive perimeter length. The natural extension of the definition (3.1) gives  $N_{g,n}(0) = 0$  since  $\mathbf{x}$  is a positive vector representing the edge lengths of  $\Gamma$ . An edge can be contracted if it is not a loop, but then we consider the result as a different graph. Therefore,  $A_{\Gamma}\mathbf{x} = 0$  has no positive solutions.

Using the lattice point interpretation, it is easy to see that the relation between this function and the Euclidean volume function is the same as that of the Riemann sum and the Riemann integral. Let  $k$  be a positive integer and  $D \subset \mathbb{R}_+^n$  an open domain with compact closure. Then for every continuous function  $f(\mathbf{p})$  on  $D$ , the definition of the Riemann integration in terms of Riemann sums gives

$$(3.2) \quad \lim_{k \rightarrow \infty} \sum_{\mathbf{p} \in D \cap \frac{1}{k}\mathbb{Z}_+^n} N_{g,n}(k\mathbf{p})f(\mathbf{p}) \frac{1}{k^{3(2g-2+n)}} = \int_D v_{g,n}^E(\mathbf{p})f(\mathbf{p})dp_1 \cdots dp_n.$$

This equality holds because our definition of the volume uses the push-forward measure. We note that as a function in  $\mathbf{p}$  there is no simple direct relation between the values  $N_{g,n}(\mathbf{p})$  and  $v_{g,n}^E(\mathbf{p})$ . For example,  $N_{g,n}(\mathbf{p}) = 0$  if  $\sum_{i=1}^n p_i$  is odd because of (2.3), but the volume function is not subject to such a relation.

To derive a topological recursion for  $N_{g,n}(\mathbf{p})$ , we introduce the notion of *ciliation*.

**Definition 3.2.** A *ciliation* is an assignment of a cilium in a face attached to a bordering edge. Let  $\ell \in \mathbb{Z}_+$  be the length of the edge on which the ciliation is attached. We place the root of the cilium at a half-integer length away from the vertices bounding the edge. Thus no cilium is attached to a vertex of a ribbon graph.




FIGURE 3.1. A ciliation in a face. The cilium is placed on a bordering edge, 0.5 unit length away from the nearest vertex.

The number of ciliations of a metric ribbon graph  $\Gamma$  with integer edge lengths is given by (2.3). Indeed, if we count with respect to the edges, then there are  $2\ell$  ways for a ciliation to each edge because the cilium can be placed on each side of the edge. And each face  $i$  has  $p_i$  ways of ciliation. Thus the total number of ciliations is  $p_1 + \cdots + p_n$ .

For brevity of notation, we denote by  $p_I = (p_i)_{i \in I}$  for a subset  $I \in N = \{1, 2, \dots, n\}$ . The cardinality of  $I$  is denoted by  $|I|$ .

**Theorem 3.3.** *The number of integral metric ribbon graphs with prescribed boundary lengths satisfies the following topological recursion formula:*

$$(3.3) \quad p_1 N_{g,n}(p_N) = \frac{1}{2} \sum_{j=2}^n \left[ \sum_{q=0}^{p_1+p_j} q(p_1+p_j-q) N_{g,n-1}(q, p_{N \setminus \{1,j\}}) \right]$$



$$\begin{aligned}
& + H(p_1 - p_j) \sum_{q=0}^{p_1-p_j} q(p_1 - p_j - q) N_{g,n-1}(q, p_{N \setminus \{1,j\}}) \\
& - H(p_j - p_1) \sum_{q=0}^{p_j-p_1} q(p_j - p_1 - q) N_{g,n-1}(q, p_{N \setminus \{1,j\}}) \Big] \\
& + \frac{1}{2} \sum_{0 \leq q_1 + q_2 \leq p_1} q_1 q_2 (p_1 - q_1 - q_2) \left[ N_{g-1,n+1}(q_1, q_2, p_{N \setminus \{1\}}) \right. \\
& \qquad \qquad \qquad \left. + \sum_{\substack{\text{stable} \\ g_1 + g_2 = g \\ I \sqcup J = N \setminus \{1\}}} N_{g_1,|I|+1}(q_1, p_I) N_{g_2,|J|+1}(q_2, p_J) \right].
\end{aligned}$$

Here

$$H(x) = \begin{cases} 1 & x > 0 \\ 0 & x \leq 0 \end{cases}$$

is the Heaviside function, and the last sum is taken for all partitions  $g = g_1 + g_2$  and  $I \sqcup J = N \setminus \{1\}$  subject to the stability condition  $2g_1 - 1 + |I| > 0$  and  $2g_2 - 1 + |J| > 0$ .

*Proof.* The key idea is to count all integral ribbon graphs with a cilium placed on the face named 1. The number is clearly equal to  $p_1 N_{g,n}(p_N)$ . We then analyze what happens when we remove the ciliated edge from the ribbon graph. There are several situations after the removal of this edge. The right-hand side of the recursion formula is obtained by the case-by-case analysis of the edge removal operation. For any ciliated ribbon graph of type  $(g, n)$  subject to the condition  $2g - 2 + n > 1$ , removing the ciliated edge creates a new graph of type  $(g, n - 1)$  or  $(g - 1, n + 1)$ , or two disjoint graphs of types  $(g_1, n_1 + 1)$  and  $(g_2, n_2 + 1)$  subject to the stability condition and the partition condition

$$\begin{cases} g_1 + g_2 = g \\ n_1 + n_2 = n - 1. \end{cases}$$

Note that in each case the quantity  $2g - 2 + n$  is reduced exactly by 1.

Let  $\eta$  be the edge bordering face 1 on which the cilium is placed, and  $a_{1\eta}$  the incidence number of (2.2). Let  $\ell \in \mathbb{Z}_+$  be the length of edge  $\eta$ . There are two main situations:  $a_{1\eta} = 1$  and  $a_{1\eta} = 2$ . Each main situation breaks down further into three cases.

**Case 1.**  $a_{1\eta} = a_{j\eta} = 1$  for  $j \geq 2$ ,  $p_1 > \ell$  and  $p_j > \ell$ . Define  $q = (p_1 - \ell) + (p_j - \ell) > 0$ . Then we have

$$\begin{cases} q - p_1 + p_j = 2(p_j - \ell) > 0 \\ q + p_1 - p_j = 2(p_1 - \ell) > 0. \end{cases}$$

Therefore,  $q > |p_1 - p_j|$ . Geometrically,  $q$  is the perimeter length of the face created by removing edge  $\eta$  that separates faces 1 and  $j$  (see Figure 3.2).

To recover the original ribbon graph with a cilium on edge  $\eta$  of length  $\ell$  from the one without edge  $\eta$ , we need to place the edge on the face of perimeter  $q$ , and place a cilium on this edge. Here we note that the data  $p_i, p_j, q$  and  $\ell$  are all prescribed. The number of ways to place an endpoint of the edge on the face of perimeter length  $q$  is  $q$ . This point uniquely determines the edge we need, since the other endpoint is  $p_1 - \ell$  away from the first endpoint along the perimeter measured by the clockwise distance. The enclosed face of perimeter length  $p_1$  becomes face 1, and the other side of the newly placed edge is face

$j$ . Since the ciliation is done on face 1, there are  $\ell$  choices for the assignment of the root of the cilium. Altogether, the contribution of this case is

$$(3.4) \quad \sum_{q=|p_1-p_j|+1}^{p_1+p_j} q \frac{p_1+p_j-q}{2} N_{g,n-1}(q, p_{N \setminus \{1,j\}}).$$




FIGURE 3.2. Case 1:  $a_{1\eta} = a_{j\eta} = 1$ ,  $p_1 > \ell$  and  $p_j > \ell$ .

**Case 2.**  $a_{1\eta} = a_{j\eta} = 1$  for  $j \geq 2$ , and  $p_1 \geq p_j = \ell$ . Since  $p_j = \ell$ , face  $j$  and edge  $\eta$  are the same and forms a loop. This loop is connected to face 1 by an edge  $\eta'$  of incidence number 2. Let  $\ell'$  be the length of this connecting edge, which is bounded by  $(p_1 - p_j)/2 \geq \ell' \geq 0$  (see Figure 3.3, left). This time define  $q = p_1 - p_j - 2\ell'$ . This is the perimeter length of the face created by removing face  $j$  and edge  $\eta'$ . In this situation, removing edge  $\eta$  ( $=$  face  $j$ ) alone does not create an admissible ribbon graph, since edge  $\eta'$  remains with a vertex of degree 1 at one end. Therefore, we need to remove the entire *tadpole* consisting of a head of face  $j$  and a tail of edge  $\eta'$ . The cilium is on face 1, which is attached to the outer boundary of face  $j$ .

To recover the original graph from the result of this tadpole removal, we have  $q$  choices for the tadpole placement and  $p_j = \ell$  choices for ciliation. Therefore, the contribution from this case is

$$(3.5) \quad \sum_{q=0}^{p_1-p_j} qp_j N_{g,n-1}(q, p_{N \setminus \{1,j\}}).$$




FIGURE 3.3. Case 2 (left):  $a_{1\eta} = a_{j\eta} = 1$  and  $p_1 \geq p_j = \ell$ ; Case 3 (center):  $a_{1\eta} = a_{j\eta} = 1$  and  $p_j \geq p_1 = \ell$ ; and Case 4 (right):  $a_{1\eta} = 2$  and the edge  $\eta$  connects a loop  $j$  to the rest of the graph.

**Case 3.**  $a_{1\eta} = a_{j\eta} = 1$  for  $j \geq 2$ , and  $p_j \geq p_1 = \ell$ . The situation is similar to Case 2 (see Figure 3.3, center). Let  $\eta'$  be the edge of length  $\ell'$  that connects face 1 and face  $j$ . Define  $q = p_j - p_1 - 2\ell'$ . This is the perimeter length of the face created by removing the entire

tadpole consisting of face 1 with a cilium as its head and edge  $\eta'$  as its tail. We have  $q$  choices for tadpole placement and  $p_1$  choices for ciliation. Thus the contribution is

$$(3.6) \quad \sum_{q=0}^{p_j-p_1} qp_1 N_{g,n-1}(q, p_{N \setminus \{1,j\}}).$$

**Case 4.**  $a_{1\eta} = 2$  and removal of edge  $\eta$  separates a single loop  $j$  for some  $j \geq 2$  from the rest of the graph (see Figure 3.3, right). It is necessary that  $p_1 > p_j$  in this case. Since a single loop alone is not an admissible graph, we need to remove face  $j$  together when we remove edge  $\eta$ . Define  $q = p_1 - p_j - 2\ell$ , which is the perimeter length of the face created after the removal of the tadpole. This time the recovery process has  $q$  choices of tadpole placement and  $2\ell$  choices for ciliation, because the cilium can be placed on either side of the tail. Thus the contribution is

$$(3.7) \quad \sum_{q=0}^{p_1-p_j} q(p_1 - p_j - q) N_{g,n-1}(q, p_{N \setminus \{1,j\}}).$$

**Case 5.**  $a_{1\eta} = 2$  and removal of edge  $\eta$  creates a connected ribbon graph. The removal of edge  $\eta$  breaks face 1 into two separate faces of perimeter lengths  $q_1$  and  $q_2$  subject to the condition  $0 < q_1 + q_2 < p_1$ . The removal of the edge reduces the genus by 1, and increases the number of faces by 1. We have the equality  $p_1 = q_1 + q_2 + 2\ell$  (see Figure 3.4). To recover the original graph from the result of the edge removal, we have  $q_1$  choices for one endpoint of edge  $\eta$ ,  $q_2$  choices for the other endpoint, and  $2\ell$  choices for ciliation, again because the cilium can be placed on either side of edge  $\eta$ . Altogether the contribution is

$$(3.8) \quad \frac{1}{2} \sum_{0 \leq q_1 + q_2 \leq p_1} q_1 q_2 (p_1 - q_1 - q_2) N_{g-1,n+1}(q_1, q_2, p_{N \setminus \{1\}}).$$

Here we need the factor  $\frac{1}{2}$ , which is the symmetry factor of interchanging  $q_1$  and  $q_2$ .




FIGURE 3.4. Case 5:  $a_{1\eta} = 2$  and removal of edge  $\eta$  creates a connected ribbon graph.

**Case 6.**  $a_{1\eta} = 2$  and removal of edge  $\eta$  creates a disjoint union of two ribbon graphs. There are  $n$  faces in the original ribbon graph  $\Gamma$ . The removal of edge  $\eta$  breaks face 1 into two separate faces of perimeter lengths  $q_1$  and  $q_2$ . The other faces  $2, 3, \dots, n$  remain intact. Let  $I \subset N \setminus \{1\}$  be the label of faces that are connected to the new face of perimeter length  $q_1$ , and  $J \subset N \setminus \{1\}$  for  $q_2$ . Then the two disjoint ribbon graphs have types  $(g_1, |I| + 1)$  and  $(g_2, |J| + 1)$  satisfying the partition condition

$$\begin{cases} g_1 + g_2 = g \\ I \sqcup J = N \setminus \{1\}. \end{cases}$$

The contribution from this case is

$$(3.9) \quad \frac{1}{2} \sum_{0 \leq q_1 + q_2 \leq p_1} q_1 q_2 (p_1 - q_1 - q_2) \sum_{\substack{\text{stable} \\ g_1 + g_2 = g \\ I \sqcup J = N \setminus \{1\}}} N_{g_1, |I|+1}(q_1, p_I) N_{g_2, |J|+1}(q_2, p_J)$$

with the symmetry factor  $\frac{1}{2}$  corresponding to interchanging  $q_1$  and  $q_2$ .




FIGURE 3.5. Case 6:  $a_{1\eta} = 2$  and removal of edge  $\eta$  creates a disjoint union of two ribbon graphs.

Summing all contributions (3.4)-(3.9), we obtain

$$(3.10) \quad \begin{aligned} p_1 N_{g,n}(p_N) &= \sum_{j=2}^n \sum_{q=|p_1-p_j|+1}^{p_1+p_j} q \frac{p_1+p_j-q}{2} N_{g,n-1}(q, p_{N \setminus \{1,j\}}) \\ &+ \sum_{j=2}^n H(p_1-p_j) \sum_{q=0}^{p_1-p_j} q(p_1-q) N_{g,n-1}(q, p_{N \setminus \{1,j\}}) \\ &+ \sum_{j=2}^n H(p_j-p_1) \sum_{q=0}^{p_j-p_1} q p_1 N_{g,n-1}(q, p_{N \setminus \{1,j\}}) \\ &+ \frac{1}{2} \sum_{0 \leq q_1 + q_2 \leq p_1} q_1 q_2 (p_1 - q_1 - q_2) \left[ N_{g-1,n+1}(q_1, q_2, p_{N \setminus \{1\}}) \right. \\ &\quad \left. + \sum_{\substack{\text{stable} \\ g_1 + g_2 = g \\ I \sqcup J = N \setminus \{1\}}} N_{g_1, |I|+1}(q_1, p_I) N_{g_2, |J|+1}(q_2, p_J) \right]. \end{aligned}$$

If we allow the variable  $q$  to range from 0 to  $p_1 + p_j$  in the first summation of the right-hand side of (3.10), then we need to compensate the non-existing cases. Note that we have

$$(3.11) \quad \begin{aligned} & - \sum_{j=2}^n \sum_{q=0}^{|p_1-p_j|} q \frac{p_1+p_j-q}{2} N_{g,n-1}(q, p_{N \setminus \{1,j\}}) \\ & + \sum_{j=2}^n H(p_1-p_j) \sum_{q=0}^{p_1-p_j} q \frac{2p_1-2q}{2} N_{g,n-1}(q, p_{N \setminus \{1,j\}}) \\ & + \sum_{j=2}^n H(p_j-p_1) \sum_{q=0}^{p_j-p_1} q \frac{2p_1}{2} N_{g,n-1}(q, p_{N \setminus \{1,j\}}) \\ & = \sum_{j=2}^n H(p_1-p_j) \sum_{q=0}^{p_1-p_j} q \frac{p_1-p_j-q}{2} N_{g,n-1}(q, p_{N \setminus \{1,j\}}) \end{aligned}$$

$$- \sum_{j=2}^n H(p_j - p_1) \sum_{q=0}^{p_j - p_1} q \frac{p_j - p_1 - q}{2} N_{g,n-1}(q, p_{N \setminus \{1,j\}}).$$

Substituting (3.11) in (3.10), we obtain (3.3). This completes the proof.  $\square$

**Remark 3.4.** The topological recursion for  $N_{g,n}(\mathbf{p})$  was considered in [38], but the formulas there contain typos. The proof given here is based on a different idea using ciliation.

#### 4. THE LAPLACE TRANSFORM OF THE NUMBER OF INTEGRAL RIBBON GRAPHS

The limit formula (3.2) tells us that  $N_{g,n}(\mathbf{p})$  asymptotically behaves like a polynomial for large  $\mathbf{p} \in \mathbb{Z}_+^n$ , and the coefficients of the leading terms correspond to that of the Euclidean volume function  $v_{g,n}^E(\mathbf{p})$ . The lack of the direct relation between  $N_{g,n}(\mathbf{p})$  and  $v_{g,n}^E(\mathbf{p})$ , together with equation (3.2), suggest that we need to consider an integral transform, such as the Laplace transform of  $N_{g,n}(\mathbf{p})$ , to extract the information of the Euclidean volume of  $RG_{g,n}(\mathbf{p})$  from it. Since

$$\int_0^\infty x^m e^{-xw} dx = \frac{m!}{w^{m+1}}$$

for a complex variable  $w \in \mathbb{C}$  with  $\text{Re}(w) > 0$ , the coefficients of the highest order poles of the Laplace transform

$$(4.1) \quad L_{g,n}(w_1, \dots, w_n) \stackrel{\text{def}}{=} \sum_{\mathbf{p} \in \mathbb{Z}_+^n} N_{g,n}(\mathbf{p}) e^{-\langle \mathbf{p}, w \rangle}$$

should represent the Euclidean volume of  $RG_{g,n}(\mathbf{p})$ . Here  $\langle p, w \rangle = p_1 w_1 + \dots + p_n w_n$ . This section is devoted to the analysis of the Laplace transform of the topological recursion (3.3).

To relate our investigation with the Hurwitz theory and the Witten-Kontsevich theory, and in particular from the point of view of the *polynomial* expressions of [14, 36], we introduce new complex coordinates

$$(4.2) \quad e^{-w} = \frac{t+1}{t-1} \quad \text{and} \quad e^{-w_j} = \frac{t_j+1}{t_j-1},$$

and express the result of the Laplace transform in terms of these  $t$ -variables. This substitution makes sense because the Laplace transform is a rational function in  $e^{-w_j}$ 's.

**Theorem 4.1.** *Define  $\mathcal{L}_{g,n}(t_1, \dots, t_n)$  by*

$$(4.3) \quad \begin{aligned} \mathcal{L}_{g,n}(t_1, \dots, t_n) dt_1 \otimes \dots \otimes dt_n &= (d_1 \otimes \dots \otimes d_n) L_{g,n}(w_1(t), \dots, w_n(t)) \\ &= \frac{\partial^n}{\partial w_1 \dots \partial w_n} L_{g,n}(w_1, \dots, w_n) dw_1 \otimes \dots \otimes dw_n \end{aligned}$$

using the coordinate change (4.2). The differentials  $dt_j$  and  $dw_j$  are related by

$$dw_j = \frac{2}{t_j^2 - 1} dt_j.$$

Then every  $\mathcal{L}_{g,n}(t_1, \dots, t_n)$  for  $2g - 2 + n > 0$  is a Laurent polynomial of degree  $3g - 3 + n$  in  $t_1^2, t_2^2, \dots, t_n^2$ . The initial values are

$$(4.4) \quad \mathcal{L}_{0,3}(t_1, t_2, t_3) = -\frac{1}{16} \left( 1 - \frac{1}{t_1^2 t_2^2 t_3^2} \right)$$

and

$$(4.5) \quad \mathcal{L}_{1,1}(t) = -\frac{1}{128} \cdot \frac{(t^2 - 1)^3}{t^4}.$$

The functions  $\mathcal{L}_{g,n}(t_1, \dots, t_n)$  for all  $(g, n)$  subject to  $2g - 2 + n > 0$  are uniquely determined by the topological recursion formula

$$(4.6) \quad \mathcal{L}_{g,n}(t_N) = -\frac{1}{16} \sum_{j=2}^n \frac{\partial}{\partial t_j} \left[ \frac{t_j}{t_1^2 - t_j^2} \left( \frac{(t_1^2 - 1)^3}{t_1^2} \mathcal{L}_{g,n-1}(t_{N \setminus \{j\}}) - \frac{(t_j^2 - 1)^3}{t_j^2} \mathcal{L}_{g,n-1}(t_{N \setminus \{1\}}) \right) \right] - \frac{1}{32} \frac{(t_1^2 - 1)^3}{t_1^2} \left[ \mathcal{L}_{g-1,n+1}(t_1, t_1, t_{N \setminus \{1\}}) + \sum_{\substack{\text{stable} \\ g_1 + g_2 = g \\ I \sqcup J = N \setminus \{1\}}} \mathcal{L}_{g_1, |I|+1}(t_1, t_I) \mathcal{L}_{g_2, |J|+1}(t_1, t_J) \right].$$

Here we use the same convention of notations as in Theorem 3.3.

If we assume (4.4), (4.5), and (4.6), then it is obvious that  $\mathcal{L}_{g,n}(t_N)$  is a Laurent polynomial in  $t_1^2, \dots, t_n^2$  of degree  $3g - 3 + n$ . The proof of (4.6) is given in Appendix A. The initial values (4.4) and (4.5) are calculated in Appendix B.

## 5. THE EUCLIDEAN VOLUME OF THE MODULI SPACE

In this section we extract the information on the Euclidean volume function from the Laurent polynomial  $\mathcal{L}_{g,n}(t_N)$ . We then derive a topological recursion for the Laplace transform of the Euclidean volume. Let us recall the Euclidean volume function  $v_{g,n}^E(\mathbf{p})$  of (2.6).

**Proposition 5.1.** *Let  $V_{g,n}^E(t_N)$  be the homogeneous leading terms of  $\mathcal{L}_{g,n}(t_N)$  for  $(g, n)$  subject to  $2g - 2 + n > 0$ . Then we have*

$$(5.1) \quad V_{g,n}^E(t_N) dt_1 \otimes \cdots \otimes dt_n = d_1 \otimes \cdots \otimes d_n \int_{\mathbb{R}_+^n} v_{g,n}^E(\mathbf{p}) e^{-\langle w, \mathbf{p} \rangle} dp_1 \cdots dp_n,$$

where we change the  $w$ -variables to the  $t$ -variables according to the transformation (4.2).

*Proof.* From (3.2), we have

$$\begin{aligned} \int_{\mathbb{R}_+^n} v_{g,n}^E(\mathbf{p}) e^{-\langle w, \mathbf{p} \rangle} dp_1 \cdots dp_n &= \lim_{k \rightarrow \infty} \sum_{\mathbf{p} \in \frac{1}{k} \mathbb{Z}_+^n} N_{g,n}(k\mathbf{p}) e^{-\langle w, \mathbf{p} \rangle} \frac{1}{k^{3(2g-2+n)}} \\ &= \lim_{k \rightarrow \infty} \sum_{\mathbf{p} \in \mathbb{Z}_+^n} N_{g,n}(\mathbf{p}) e^{-\frac{1}{k} \langle w, \mathbf{p} \rangle} \frac{1}{k^{3(2g-2+n)}} \\ &= \lim_{k \rightarrow \infty} L_{g,n} \left( \frac{w_1}{k}, \dots, \frac{w_n}{k} \right) \frac{1}{k^{3(2g-2+n)}}. \end{aligned}$$

The coordinate transformation (4.2) has the expansion near  $w = 0$

$$(5.2) \quad \begin{aligned} t &= t(w) = -\frac{2}{w} - \frac{w}{6} + \frac{w^3}{360} - \frac{w^5}{15120} + \cdots, \\ w &= w(t) = -\frac{2}{t} - \frac{2}{3t^3} - \frac{2}{5t^5} - \cdots. \end{aligned}$$

Since

$$\mathcal{L}_{g,n}(t_N) = \frac{\partial^n}{\partial t_1 \cdots \partial t_n} L_{g,n}(w(t_1), \dots, w(t_n))$$

is a Laurent polynomial of degree  $2(3g - 3 + n)$ , and since the change  $w \mapsto w/k$  makes

$$t \mapsto k t + \mathcal{O}\left(\frac{1}{k}\right)$$

for a fixed value  $t$ , we have

$$\begin{aligned} & \frac{\partial^n}{\partial t_1 \cdots \partial t_n} \int_{\mathbb{R}_+^n} v_{g,n}^E(\mathbf{p}) e^{-\langle w(t), \mathbf{p} \rangle} dp_1 \cdots dp_n \\ &= \lim_{k \rightarrow \infty} \frac{\partial^n}{\partial t_1 \cdots \partial t_n} L_{g,n} \left( \frac{w(t_1)}{k}, \dots, \frac{w(t_n)}{k} \right) \frac{1}{k^{3(2g-2+n)}} \\ &= \lim_{k \rightarrow \infty} \mathcal{L}_{g,n} \left( kt_1 + \mathcal{O}\left(\frac{1}{k}\right), \dots, kt_n + \mathcal{O}\left(\frac{1}{k}\right) \right) \frac{k^n}{k^{3(2g-2+n)}} = V_{g,n}^E(t_N). \end{aligned}$$

This completes the proof.  $\square$

Since  $v_{g,n}^E(\mathbf{p})$  is defined by the push-forward measure of the incidence matrix  $A_\Gamma$  of (2.2) at each point  $\Gamma \in RG_{g,n}(\mathbf{p})$ , we have

$$\begin{aligned} (5.3) \quad & \int_{\mathbb{R}_+^n} v_{g,n}^E(\mathbf{p}) e^{-\langle w, \mathbf{p} \rangle} dp_1 \cdots dp_n \\ &= \sum_{\substack{\Gamma \text{ trivalent ribbon} \\ \text{graph of type } (g,n)}} \frac{1}{|\text{Aut}(\Gamma)|} \int_{\mathbb{R}_+^n} \text{vol}(P_\Gamma(\mathbf{p})) e^{-\langle w, \mathbf{p} \rangle} dp_1 \cdots dp_n \\ &= \sum_{\substack{\Gamma \text{ trivalent ribbon} \\ \text{graph of type } (g,n)}} \frac{1}{|\text{Aut}(\Gamma)|} \int_{\mathbb{R}_+^{e(\Gamma)}} e^{-\langle w, A_\Gamma \mathbf{x} \rangle} dx_1 \cdots dx_{e(\Gamma)} \\ &= \sum_{\substack{\Gamma \text{ trivalent ribbon} \\ \text{graph of type } (g,n)}} \frac{1}{|\text{Aut}(\Gamma)|} \prod_{\eta=1}^{e(\Gamma)} \frac{1}{\langle w, a_\eta \rangle}, \end{aligned}$$

where  $a_1, \dots, a_{e(\Gamma)}$  are columns of the edge-face incidence matrix

$$A_\Gamma = [a_1 | a_2 | \cdots | a_{e(\Gamma)}].$$

We note that  $e(\Gamma)$  takes its maximum value  $3(2g - 2 + n)$  for a trivalent graph. Thus the last line of (5.3) has a pole of order  $3(2g - 2 + n)$  at  $w = 0$ . This expression also shows that the leading terms of  $L_{g,n}(w(t_N))$  as a function in  $t_N$  using the expansion (5.2) around  $t_N \sim \infty$  are the Laplace transform of the Euclidean volume function. In particular, we deduce that  $N_{g,n}(\mathbf{p})$  behaves asymptotically like a polynomial of degree  $2(3g - 3 + n)$  for large  $\mathbf{p} \in \mathbb{R}_+^n$ .

Since  $V_{g,n}^E(t_N)$  is the leading terms of  $\mathcal{L}_{g,n}(t_N)$ , it is easy to obtain a topological recursion.

**Theorem 5.2.** *The Laplace transformed Euclidean volume function  $V_{g,n}^E(t_N)$  in the stable range  $2g - 2 + n > 0$  satisfies the following topological recursion:*

$$(5.4) \quad V_{g,n}^E(t_N) = -\frac{1}{16} \sum_{j=2}^n \frac{\partial}{\partial t_j} \left[ \frac{t_j}{t_1^2 - t_j^2} \left( t_1^4 V_{g,n-1}^E(t_{N \setminus \{j\}}) - t_j^4 V_{g,n-1}^E(t_{N \setminus \{1\}}) \right) \right]$$

$$-\frac{1}{32} t_1^4 \left[ V_{g-1, n+1}^E(t_1, t_1, t_{N \setminus \{1\}}) + \sum_{\substack{\text{stable} \\ g_1+g_2=g \\ I \sqcup J = N \setminus \{1\}}} V_{g_1, |I|+1}^E(t_1, t_I) V_{g_2, |J|+1}^E(t_1, t_J) \right].$$

*Proof.* The leading contribution of (4.6) comes from the leading term of

$$\frac{(t^2 - 1)^3}{t^2} = t^4 - 3t^2 + 3 - \frac{1}{t^2}.$$

Thus (4.6) reduces to (5.4).  $\square$

## 6. THE SYMPLECTIC VOLUME OF THE MODULI SPACE AND THE KONTSEVICH CONSTANTS

Suppose the  $i$ -th face of a metric ribbon graph  $\Gamma \in RG_{g,n}(\mathbf{p})$  consists of edges labeled by  $1, 2, \dots, k$  in this cyclic order. (Here again we are abusing the notation to indicate a metric ribbon graph by the same letter  $\Gamma$ .) If an edge appears twice in this list, then we count it repetitively. Denote by  $\ell_\alpha$  the length of edge  $\alpha$ . They satisfy the relation  $\ell_1 + \dots + \ell_k = p_i$ . Note that the collection of edge lengths forms an orbifold coordinate system on  $RG_{g,n}$  at each point  $\Gamma$ . Kontsevich [28] defines a 2-form on  $RG_{g,n}$  by

$$(6.1) \quad \omega_K(\mathbf{p}) = \sum_{i=1}^n p_i^2 \omega_i, \quad \omega_i = \sum_{\alpha < \beta} d\left(\frac{\ell_\alpha}{p_i}\right) \wedge d\left(\frac{\ell_\beta}{p_i}\right) \quad \text{on face } i.$$

If we change the cyclic order from  $(1, 2, \dots, k)$  to  $(2, 3, \dots, k, 1)$  and define the form  $\omega'_i$  in the same manner, then we have

$$\omega_i - \omega'_i = 2d\left(\frac{\ell_1}{p_i}\right) \wedge d\left(\frac{\ell_2 + \dots + \ell_k}{p_i}\right) = 0.$$

Therefore, each  $\omega_i$  and  $\omega_K(\mathbf{p})$  are well defined as genuine 2-forms on  $RG_{g,n}$ . The restriction of the 2-form  $\omega_K(\mathbf{p})$  defines a symplectic structure on  $RG_{g,n}(\mathbf{p}) \cong \mathcal{M}_{g,n}$  for each  $\mathbf{p} \in \mathbb{R}_+^n$ .

To see the non-degeneracy of  $\omega_K(\mathbf{p})$ , let us analyze the perimeter map  $\pi$  locally around a trivalent ribbon graph  $\Gamma$ . As in Section 2 we give a name to all edges of  $\Gamma$ , this time without repetition, indexed by  $\{0, 1, 2, \dots, e(\Gamma) - 1\}$ . Faces of  $\Gamma$  are indexed by  $N = \{1, 2, \dots, n\}$ . The edge-face incidence matrix  $A_\Gamma$  of (2.2) gives the differential of the perimeter map

$$A_\Gamma = d\pi_\Gamma$$

at the metric ribbon graph  $\Gamma$  if it is trivalent. To set notations simple, we assume that faces 1 through 4 and edges 0 through 4 are arranged as in Figure 6.1.




FIGURE 6.1. The vector field  $X_0$ .

Define the vector field

$$(6.2) \quad X_0 = -\frac{\partial}{\partial \ell_1} + \frac{\partial}{\partial \ell_2} - \frac{\partial}{\partial \ell_3} + \frac{\partial}{\partial \ell_4}.$$



We then have

$$\begin{aligned}\iota_{X_0}(p_1^2\omega_1)|_{RG_{g,n}(\mathbf{p})} &= -dl_0 - dl_1 - dl_4 \\ \iota_{X_0}(p_2^2\omega_2)|_{RG_{g,n}(\mathbf{p})} &= dl_1 + dl_2 \\ \iota_{X_0}(p_3^2\omega_3)|_{RG_{g,n}(\mathbf{p})} &= -dl_0 - dl_2 - dl_3 \\ \iota_{X_0}(p_4^2\omega_4)|_{RG_{g,n}(\mathbf{p})} &= dl_3 + dl_4.\end{aligned}$$

Therefore,

$$\iota_{X_0}\omega_K(\mathbf{p})|_{RG_{g,n}(\mathbf{p})} = -2dl_0$$

on the tangent space  $T_\Gamma RG_{g,n}(\mathbf{p})$ . This shows that the 2-form  $\omega_K(\mathbf{p})$  restricted on  $\text{Ker}(d\pi_\Gamma)$  is a linear isomorphism. We refer to [2] for more detail.

Alternatively, we can introduce the symplectic structure on  $RG_{r,n}(\mathbf{p})$  through symplectic reduction. The ribbon graph complex  $RG_{g,n}$  comes with a natural fibration on it, the *tautological torus bundle*

$$(6.3) \quad \begin{array}{ccc} \mathbb{T} & \xrightarrow{\mu} & \mathbb{R}_+^n \\ \tau \downarrow & & \cdot \\ & & RG_{g,n} \end{array}$$

The fiber of  $\tau$  at a metric ribbon graph  $\Gamma$  is the cartesian product of the boundary of the  $n$  faces of  $\Gamma$ , which is identified with the collection of  $n$  polygons. Topologically each fiber of  $\mathbb{T}$  is an  $n$ -dimensional torus  $T^n = (S^1)^n$ . We use the same letter  $\mathbb{T}$  for the total space of this torus bundle, whose dimension  $2(3g - 3 + 2n)$  is always even.

The identification of the  $i$ -th face of  $\Gamma \in RG_{r,n}(\mathbf{p})$  and the circle  $S^1 = \mathbb{R}/\mathbb{Z}$  is given as follows. First we choose a vertex on the  $i$ -th polygon, and name the edges on the  $i$ -th face as  $1, 2, \dots, k$  in this cyclic order such that the chosen vertex is the beginning point of edge 1 and the end point of edge  $k$ . Let  $\ell_\alpha$  be the length of edge  $\alpha$  as before. We choose a parameter  $\phi_i$  subject to  $0 \leq \phi_i \leq \ell_1$ . Under the re-naming of the edges  $(1, 2, \dots, k) \mapsto (2, 3, \dots, k, 1)$ ,  $\phi_i$  changes to  $\phi'_i = \phi_i + \ell_1$ . The choice of the vertex and  $\phi_i$  is identified with an element of  $S^1$ , and also determines the torus action on the fibration  $\mathbb{T}$ .

Define a 2-form  $\Omega$  by

$$(6.4) \quad \begin{aligned}\Omega &= \sum_{i=1}^n \hat{\omega}_i \\ \hat{\omega}_i &= \sum_{\alpha < \beta} dl_\alpha \wedge dl_\beta + d\left(\frac{\phi_i}{p_i}\right) \wedge d(p_i^2).\end{aligned}$$

The cyclic re-naming of edges changes  $\hat{\omega}_i$  to

$$\hat{\omega}'_i = \sum_{2 \leq \alpha < \beta} dl_\alpha \wedge dl_\beta + \sum_{2 \leq \alpha} dl_\alpha \wedge dl_1 + d\left(\frac{\phi_i + \ell_1}{p_i}\right) \wedge d(p_i^2).$$

Therefore,

$$\hat{\omega}_i - \hat{\omega}'_i = \sum_{2 \leq \beta} dl_1 \wedge dl_\beta - \sum_{2 \leq \alpha} dl_\alpha \wedge dl_1 + 2dl_1 \wedge dp_i = 0,$$

and hence  $\Omega$  is a globally well-defined 2-form on the total space  $\mathbb{T}$ . The moment map of the torus action on  $\mathbb{T}$  is the assignment

$$\mu : \mathbb{T} \ni (\Gamma, \phi_1, \dots, \phi_n) \mapsto (p_1^2, \dots, p_n^2) \in \mathbb{R}_+^n.$$

The symplectic quotient  $\mu^{-1}(L)//T^n$  of  $\mathbb{T}$  by this torus action is  $(RG_{g,n}(\mathbf{p}), \omega_K(\mathbf{p}))$  of (6.1).

Now we define the symplectic volume of the moduli space  $\mathcal{M}_{g,n} \cong RG_{g,n}(\mathbf{p})$  by

$$(6.5) \quad v_{g,n}^S(\mathbf{p}) = \int_{RG_{g,n}(\mathbf{p})} \exp(\omega_K(\mathbf{p})).$$

Applying the recursion argument similar to our proof of Theorem 3.3 to the symplectic reduction of  $RG_{g,n}$  by the torus action, the following theorem was established in [2].

**Theorem 6.1** ([2]). *The symplectic volume satisfies the following topological recursion.*

$$(6.6) \quad \begin{aligned} p_1 v_{g,n}^S(p_N) = & \sum_{j=2}^n \left[ \int_0^{p_1+p_j} q(p_1+p_j-q) v_{g,n-1}^S(q, p_{N \setminus \{1,j\}}) dq \right. \\ & + H(p_1-p_j) \int_0^{p_1-p_j} q(p_1-p_j-q) v_{g,n-1}^S(q, p_{N \setminus \{1,j\}}) dq \\ & \left. - H(p_j-p_1) \int_0^{p_j-p_1} q(p_j-p_1-q) v_{g,n-1}^S(q, p_{N \setminus \{1,j\}}) dq \right] \\ & + 2 \iint_{0 \leq q_1+q_2 \leq p_1} q_1 q_2 (p_1 - q_1 - q_2) \left[ v_{g-1, n+1}^S(q_1, q_2, p_{N \setminus \{1\}}) \right. \\ & \left. + \sum_{\substack{\text{stable} \\ g_1+g_2=g \\ I \sqcup J = N \setminus \{1\}}} v_{g_1, |I|+1}^S(q_1, p_I) v_{g_2, |J|+1}^S(q_2, p_J) dq_1 dq_2 \right]. \end{aligned}$$

The initial values are easy to calculate. For the case of  $(g, n) = (0, 3)$ , since the perimeter  $(p_1, p_2, p_3) \in \mathbb{R}_+^3$  determines the length of each edge, the symplectic form is 1 on a single point. Thus we have

$$(6.7) \quad v_{0,3}^S(p_1, p_2, p_3) = 1.$$

The unique trivalent graph of type  $(g, n) = (1, 1)$  is given in Figure 6.2, which has the automorphism group  $\mathbb{Z}/6\mathbb{Z}$ . The perimeter map is given by  $p = 2(\ell_1 + \ell_2 + \ell_3)$ . The restriction of  $\omega_K(p)$  on  $RG_{1,1}(p)$  is  $2d\ell_1 \wedge d\ell_2$ . Therefore, we have

$$(6.8) \quad v_{1,1}^S(p) = \frac{1}{6} \int_{0 \leq \ell_1 + \ell_2 \leq \frac{p}{2}} 2d\ell_1 \wedge d\ell_2 = \frac{1}{24} p^2.$$




FIGURE 6.2. The trivalent ribbon graph of type  $(1, 1)$ .

We now consider the Laplace transform of the symplectic volume  $v_{g,n}^S(\mathbf{p})$ .

**Theorem 6.2.** *The symmetric function  $V_{g,n}^S(t_N)$  defined by the Laplace transform*

$$(6.9) \quad V_{g,n}^S(t_1, \dots, t_n) dt_1 \otimes \dots \otimes dt_n \stackrel{\text{def}}{=} d_1 \otimes \dots \otimes d_n \int_{\mathbb{R}_+^n} v_{g,n}^S(\mathbf{p}) e^{-\langle w, \mathbf{p} \rangle} dp_1 \dots dp_n$$

and the coordinate change

$$(6.10) \quad w_j = -\frac{2}{t_j}$$

satisfies the topological recursion

$$(6.11) \quad V_{g,n}^S(t_N) = -\frac{1}{4} \sum_{j=2}^{\infty} \frac{\partial}{\partial t_j} \left[ \frac{t_j}{t_1^2 - t_j^2} \left( t_1^4 V_{g,n-1}^S(w_{N \setminus \{j\}}) - t_j^4 V_{g,n-1}^S(w_{N \setminus \{1\}}) \right) \right] \\ - \frac{1}{4} t_1^4 \left( V_{g-1,n+1}^S(t_1, t_1, t_{N \setminus \{1\}}) + \sum_{\substack{g_1+g_2=g, \\ I \sqcup J = N \setminus \{1\}}} V_{g_1,|I|+1}^S(t_1, t_I) V_{g_2,|J|+1}^S(t_1, t_J) \right).$$

The proof of this theorem is given in Appendix A. The very reason that Kontsevich was interested in the symplectic volume of the moduli space is that it gives the generating function of the intersection numbers (1.2)

$$(6.12) \quad V_{g,n}^S(t_N) = (-1)^n \sum_{\substack{d_1+\dots+d_n \\ =3g-3+n}} \langle \tau_{d_1} \dots \tau_{d_n} \rangle_{g,n} \prod_{j=1}^n (2d_j + 1)!! \left( \frac{t_j}{2} \right)^{2d_j}.$$

The topological recursion (6.11) produces a relation among the coefficients, which is known as the DVV formula of [9], and is equivalent to the Virasoro constraint condition of [48].

Since the volume is for  $\mathcal{M}_{g,n}$  and the intersection numbers are for  $\overline{\mathcal{M}}_{g,n}$ , it is not obvious why they are the same thing. From the deep theory of Mirzakhani [32, 33], it becomes obvious why and how they are related.

We are now ready to calculate the Kontsevich constants.

**Theorem 6.3.** *The ratio of the two volume polynomials  $V_{g,n}^S(t_N)$  and  $V_{g,n}^E(t_N)$  is a constant depending only on  $g$  and  $n$ :*

$$(6.13) \quad \rho_{g,n}(t) \stackrel{\text{def}}{=} \frac{V_{g,n}^S(t_N)}{V_{g,n}^E(t_N)} = 2^{5g-5+2n}.$$

*Proof.* We use induction on  $2g - 2 + n$ . From (6.7), (6.8), (B.3) and (B.6), we have

$$(6.14) \quad \begin{cases} V_{0,3}^E(t_1, t_2, t_3) = -\frac{1}{16} \\ V_{0,3}^S(t_1, t_2, t_3) = -\frac{1}{8} \end{cases} \quad \text{and} \quad \begin{cases} V_{1,1}^E(t) = -\frac{1}{128} t^2 \\ V_{1,1}^S(t) = -\frac{1}{32} t^2 \end{cases}.$$

Thus the initial values satisfy (6.13). We observe that the recursion formulas (5.4) and (6.11) are the same except for the constant factors on the first and the second lines of the right-hand side. Therefore, if we changed  $V_{g,n}^E(t_N)$  to  $2^{5g-5+2n} \cdot V_{g,n}^E(t_N)$  in (5.4), then its recursion formula would become identical to (6.11). Since the recursion uniquely determines all values for  $(g, n)$  subject to  $2g - 2 + n > 0$  from the initial values (6.14), we establish (6.13). This completes the proof.  $\square$

7. THE EYNARD-ORANTIN THEORY ON  $\mathbb{P}^1$ 

The number of integral ribbon graphs  $N_{g,n}(\mathbf{p})$  is a difficult function to deal with because it is not given by a single formula. As we have noted, it behaves like a polynomial for large  $\mathbf{p} \in \mathbb{Z}_+^n$ , while it takes value 0 whenever  $p_1 + \cdots + p_n$  is odd. Compared to this, the Laplace transformed function such as  $\mathcal{L}_{g,n}(t_N)$  is a far nicer object. Indeed  $\mathcal{L}_{g,n}(t_N)$  is a Laurent polynomial and satisfies a simple differential recursion formula (4.6). We also note that the recursion formulas (4.6), (5.4), and (6.11) take a very similar shape. Over the years several authors (including [4, 5, 8, 13, 14, 15, 16, 28, 31, 36, 42, 50, 51]) have noticed that many different combinatorial structures (on the A-model side of a topological string theory) can be uniformly treated on the B-model side, after taking the Laplace transform. The importance of the Laplace transform as the mirror map was noted in [14]. This uniform structure after the Laplace transform is the manifestation of the Eynard-Orantin theory. We will show in this section that the recursions (4.6), (5.4), and (6.11) become identical under the formalism proposed in [15].

We are not in the place to formally present the Eynard-Orantin formalism in an axiomatic way. Instead of giving the full account, we are satisfied with explaining a limited case when the *spectral curve* of the theory is  $\mathbb{P}^1$ . The word ‘‘spectral curve’’ was used in [15] because of the analogy of the spectral curves appearing in the Lax formalism of integrable systems.

We start with the spectral curve  $C = \mathbb{P}^1 \setminus S$ , where  $S \subset \mathbb{P}^1$  is a finite set. We also need two generic elements  $x$  and  $y$  of  $H^0(C, \mathcal{O}_C)$ , where  $\mathcal{O}_C$  denotes the sheaf of holomorphic functions on  $C$ . The condition we impose on  $x$  and  $y$  is that the holomorphic maps

$$(7.1) \quad x : C \longrightarrow \mathbb{C} \quad \text{and} \quad y : C \longrightarrow \mathbb{C}$$

have only simple ramification points, i.e., their derivatives  $dx$  and  $dy$  have simple zeros, and that

$$(7.2) \quad (x, y) : C \ni t \longmapsto (x(t), y(t)) \in \mathbb{C}^2$$

is an immersion. Let  $\Lambda^1(C)$  denote the sheaf of meromorphic 1-forms on  $C$ , and

$$(7.3) \quad H^n = H^0(C^n, \text{Sym}^n(\Lambda^1(C)))$$

the space of meromorphic symmetric differentials of degree  $n$ . The Cauchy differentiation kernel is an example of such differentials:

$$(7.4) \quad W_{0,2}(t_1, t_2) = \frac{dt_1 \otimes dt_2}{(t_1 - t_2)^2} \in H^2.$$

In the literatures starting from [15], the Cauchy differentiation kernel has been called the *Bergman kernel*, even though it has nothing to do with the Bergman kernel in complex analysis. A bilinear operator

$$(7.5) \quad K : H \otimes H \longrightarrow H$$

naturally extends to

$$\begin{aligned} K : H^{n_1+1} \otimes H^{n_2+1} &\ni (f_0, f_1, \dots, f_{n_1}) \otimes (h_0, h_1, \dots, h_{n_2}) \\ &\longmapsto (K(f_0, h_0), f_1, \dots, f_{n_1}, h_1, \dots, h_{n_2}) \in H^{n_1+n_2+1} \end{aligned}$$

$$K : H^{n+1} \ni (f_0, f_1, \dots, f_n) \longmapsto (K(f_0, f_1), f_2, \dots, f_n) \in H^n.$$

Suppose we are given an infinite sequence  $\{W_{g,n}\}$  of differentials  $W_{g,n} \in H^n$  for all  $(g, n)$  subject to the stability condition  $2g - 2 + n > 0$ . We say *this sequence satisfies a topological*

recursion with respect to the kernel  $K$  if

$$(7.6) \quad W_{g,n} = K(W_{g,n-1}, W_{0,2}) + K(W_{g-1,n+1}) + \frac{1}{2} \sum_{\substack{\text{stable} \\ g_1+g_2=g \\ I \sqcup J = N \setminus \{1\}}} K(W_{g_1,|I|+1}, W_{g_2,|J|+1}).$$

The characteristic of the Eynard-Orantin theory lies in the particular choice of the *Eynard kernel* that reflects the parametrization (7.2) and the ramified coverings (7.1). Let  $\mathcal{A} = \{a_1, \dots, a_r\} \subset C$  be the set of simple ramification points of the  $x$ -projection map. Since locally at each  $a_\lambda$  the  $x$ -projection is a double-sheeted covering, we can choose the deck transformation map

$$(7.7) \quad s_\lambda : U_\lambda \xrightarrow{\sim} U_\lambda,$$

where  $U_\lambda \subset C$  is an appropriately chosen simply connected neighborhood of  $a_\lambda$ .

**Definition 7.1.** The *Eynard kernel* is the linear map  $H \otimes H \rightarrow H$  defined by

$$(7.8) \quad K(f_1(t_1)dt_1, f_2(t_2)dt_2) = \frac{1}{2\pi i} \sum_{\lambda=1}^r \oint_{|t-a_\lambda|<\epsilon} K_\lambda(t, t_1) \left( f_1(t)dt \otimes f_2(s_\lambda(t))ds_\lambda(t) + f_2(t)dt \otimes f_1(s_\lambda(t))ds_\lambda(t) \right),$$

where

$$(7.9) \quad K_\lambda(t, t_1) = \frac{1}{2} \left( \int_t^{s_\lambda(t)} W_{0,2}(t, t_1) dt \right) \otimes dt_1 \cdot \frac{1}{\left( y(t) - y(s_\lambda(t)) \right) dx(t)},$$

and  $\frac{1}{dx(t)}$  is the contraction operator with respect to the vector field

$$\left( \frac{dx}{dt} \right)^{-1} \frac{\partial}{\partial t}.$$

The integration is taken with respect to the  $t$ -variable along a small loop around  $a_\lambda$  that contains no singularities other than  $t = a_\lambda$ . A topological recursion with respect to the Eynard kernel is what we call the *Eynard-Orantin recursion* in this paper.

To convert (4.6) to the Eynard-Orantin formalism, we need to identify the spectral curve of the theory and the unstable case  $\mathcal{L}_{0,2}(t_1, t_2)$ . The spectral curve is a plane algebraic curve

$$(7.10) \quad C = \{ (x, y) \in \mathbb{C}^2 \mid xy = y^2 + 1 \},$$

which is the same curve considered in [39]. Here we introduce a different parametrization

$$(7.11) \quad \begin{aligned} x(t) &= \frac{t+1}{t-1} + \frac{t-1}{t+1} = 2 + \frac{4}{t^2-1} \\ y(t) &= \frac{t+1}{t-1} \end{aligned}$$

with a parameter  $t \in \mathbb{P}^1 \setminus \{1, -1\}$  so that the resulting differentials become Laurent polynomials. This use of the parametrization is similar to that of [14, 36]. The  $x$ -projection

$$(7.12) \quad \pi : C \ni t \longmapsto x(t) \in \mathbb{C}$$

has simple ramification points at  $t = 0$  and  $t = \infty$ , since

$$dx = -\frac{8t}{(t^2-1)^2} dt.$$

We note that since the map  $\pi$  is globally a branched double-sheeted covering, its covering transformation is globally defined and is given by

$$(7.13) \quad s : C \ni t \longmapsto s(t) = -t \in C.$$




FIGURE 7.1. The spectral curve  $x = y + \frac{1}{y}$ .

The unstable  $(0, 2)$  case is calculated in Appendix B, (B.9). The result is

$$\mathcal{L}_{0,2}(t_1, t_2) dt_1 \otimes dt_2 = \frac{dt_1 \otimes dt_2}{(t_1 + t_2)^2} = \frac{dt_1 \otimes dt_2}{(t_1 - s(t_2))^2}.$$

This quadratic differential form plays the role of the Cauchy differentiation kernel. For every holomorphic differential  $f(t)dt$  on  $C$ , we have

$$(7.14) \quad -\frac{1}{2\pi i} \oint \frac{1}{dt} \left[ f(s(t)) ds(t) \mathcal{L}_{0,2}(t, t_1) dt \otimes dt_1 + f(t) dt \mathcal{L}_{0,2}(s(t), t_1) ds(t) \otimes dt_1 \right] \\ = \left( \frac{1}{2\pi i} \oint \frac{f(-t)}{(t + t_1)^2} dt \right) \otimes dt_1 + \left( \frac{1}{2\pi i} \oint \frac{f(t)}{(t - t_1)^2} dt \right) \otimes dt_1 = 2f'(t_1) dt_1,$$

where the operation  $\frac{1}{dt}$  is the contraction by the vector field  $\frac{\partial}{\partial t}$ , and the integration is taken with respect to  $t$  along a positively oriented simple loop that contains both  $t_1$  and  $s(t_1)$ . Actually, the contour integral should be considered as the residue calculation at  $t = \infty$  with respect to the opposite orientation. This explains the minus sign in (7.14).

**Theorem 7.2.** *The topological recursion (4.6) is equivalent to the Eynard-Orantin recursion of [15]:*

$$(7.15) \quad \mathcal{L}_{g,n}(t_N) dt_N \\ = \frac{1}{2\pi i} \int_{\Gamma} K(t, t_1) \left[ \sum_{j=2}^n \left( \mathcal{L}_{g,n-1}(t, t_{N \setminus \{1,j\}}) dt \otimes dt_{N \setminus \{1,j\}} \otimes \mathcal{L}_{0,2}(s(t), t_j) ds(t) \otimes dt_j \right. \right. \\ \left. \left. + \mathcal{L}_{g,n-1}(s(t), t_{N \setminus \{1,j\}}) ds(t) \otimes dt_{N \setminus \{1,j\}} \otimes \mathcal{L}_{0,2}(t, t_j) dt \otimes dt_j \right) \right. \\ \left. + \mathcal{L}_{g-1,n+1}(t, s(t), t_{N \setminus \{1\}}) dt \otimes ds(t) \otimes dt_{N \setminus \{1\}} \right. \\ \left. + \sum_{\substack{\text{stable} \\ g_1 + g_2 = g \\ I \sqcup J = N \setminus \{1\}}} \left( \mathcal{L}_{g_1, |I|+1}(t, t_I) dt \otimes dt_I \right) \otimes \left( \mathcal{L}_{g_2, |J|+1}(s(t), t_J) ds(t) \otimes dt_J \right) \right].$$

Here the contour integration is taken with respect to  $t$  along a curve  $\Gamma$  that consists of a large circle of the negative orientation centered at the origin with radius  $r > \max_{j \in N} |t_j|$ ,

and a small circle around the origin of the positive orientation. We use a simplified notation  $dt_I = \otimes_{i \in I} dt_i$  for  $I \subset N$ .




FIGURE 7.2. The integration contour  $\Gamma$ . This contour encloses an annulus bounded by two concentric circles centered at the origin. The outer one has a large radius  $r > \max_{j \in N} |t_j|$  and the negative orientation, and the inner one has an infinitesimally small radius with the positive orientation.

- Remark 7.3.** (1) The contour integral (7.15) can be phrased as the sum of the *residues* of the integrand at the ramification points of the spectral curve  $t = 0$  and  $t = \infty$ , which is the language used in [15].
- (2) The first and the second lines of the right-hand side of (7.15) are unstable  $(0, 2)$  cases of the fourth line when we have  $(g_1, I) = (0, \{j\})$  or  $(g_2, J) = (0, \{j\})$ .
- (3) In terms of the Cauchy differentiation kernel  $W_{0,2}(t, t_1)$  of (7.4), we have

$$\mathcal{L}_{0,2}(t, t_1) dt \otimes dt_1 = W_{0,2}(t, t_1) - \pi^* W_{0,2}(x, x_1)$$

as proved in Appendix B, (B.10). Since  $\pi^* W_{0,2}(x, x_1)$  is invariant under the deck transformation  $s : C \rightarrow C$  applied to the entry  $x$ , and since  $\mathcal{L}_{g,n}(t_N)$  is an even function by Theorem 4.1, we can replace  $\mathcal{L}_{0,2}(t, t_1) dt \otimes dt_1$  with  $W_{0,2}(t, t_1)$  in (7.15).

*Proof.* The Eynard kernel of our setting is

$$\begin{aligned} K(t, t_1) &= \frac{1}{2} \left( \int_t^{s(t)} \mathcal{L}_{0,2}(t, t_1) dt \right) \otimes dt_1 \cdot \frac{1}{\left( y(t) - y(s(t)) \right) dx(t)} \\ &= \frac{1}{2} \left( \int_t^{s(t)} \frac{dt}{(t + t_1)^2} \right) \otimes dt_1 \cdot \frac{1}{\left( \frac{t+1}{t-1} - \frac{-t+1}{-t-1} \right) \frac{\partial}{\partial t} \left( \frac{t+1}{t-1} + \frac{t-1}{t+1} \right)} \cdot \frac{1}{dt} \\ &= -\frac{1}{2} \left( \frac{1}{t - t_1} + \frac{1}{t + t_1} \right) \frac{1}{32} \frac{(t^2 - 1)^3}{t^2} \cdot \frac{1}{dt} \otimes dt_1. \end{aligned}$$

Thus for any symmetric Laurent polynomial  $f(t, s)$  in  $t^2$  and  $s^2$ , we have

$$\frac{1}{2\pi i} \int_{\Gamma} K(t, t_1) f(t, s(t)) dt \otimes ds(t) = -f(t_1, t_1) \frac{1}{32} \frac{(t_1^2 - 1)^3}{t_1^2} dt_1,$$

since  $s(t) = -t$ . Therefore, the third and the fourth lines of the right-hand side of (7.15) becomes

$$-\frac{1}{32} \frac{(t_1^2 - 1)^3}{t_1^2} \left[ \mathcal{L}_{g-1, n+1}(t_1, t_1, t_{N \setminus \{1\}}) \right]$$

$$+ \left. \sum_{\substack{\text{stable} \\ g_1+g_2=g \\ I \sqcup J = N \setminus \{1\}}} \mathcal{L}_{g_1,|I|+1}(t_1, t_I) \mathcal{L}_{g_2,|J|+1}(t_1, t_J) \right] \otimes dt_N.$$

This is because  $\mathcal{L}_{g,n}(t_N)$  for  $(g, n)$  in the stable range is a Laurent polynomial in  $t_1^2, \dots, t_n^2$ , hence the only simple poles in the complex  $t$ -plane within the contour  $\Gamma$  of (7.15) that appear in the third and fourth lines are located at  $t = t_1$  and  $t = -t_1$ .

Even though the first and the second lines of the right-hand side of (7.15) are somewhat a degenerate case of the fourth line as remarked above, the analysis becomes different because  $\mathcal{L}_{0,2}(t, t_j)$  contributes new poles in the  $t$ -plane. First we note that

$$\begin{aligned} (7.16) \quad & \mathcal{L}_{g,n-1}(t, t_{N \setminus \{1,j\}}) dt \otimes dt_{N \setminus \{1,j\}} \otimes \mathcal{L}_{0,2}(s(t), t_j) ds(t) \otimes dt_j \\ & + \mathcal{L}_{g,n-1}(s(t), t_{N \setminus \{1,j\}}) ds(t) \otimes dt_{N \setminus \{1,j\}} \otimes \mathcal{L}_{0,2}(t, t_j) dt \otimes dt_j \\ & = -\mathcal{L}_{g,n-1}(t, t_{N \setminus \{1,j\}}) \left( \frac{1}{(-t+t_j)^2} + \frac{1}{(t+t_j)^2} \right) dt^{\otimes 2} \otimes dt_{N \setminus \{1\}} \\ & = -2\mathcal{L}_{g,n-1}(t, t_{N \setminus \{1,j\}}) \frac{\partial}{\partial t_j} \frac{t_j}{t^2 - t_j^2} dt^{\otimes 2} \otimes dt_{N \setminus \{1\}}. \end{aligned}$$

Apply the operation  $\frac{1}{2\pi i} \int_{\Gamma} K(t, t_1)$  to (7.16) and collect the residues at  $t = t_1$  and  $t = -t_1$ . We then obtain

$$-\frac{1}{16} \frac{\partial}{\partial t_j} \left[ \frac{t_j}{t_1^2 - t_j^2} \frac{(t_1^2 - 1)^3}{t_1^2} \mathcal{L}_{g,n-1}(t_{N \setminus \{j\}}) \right] dt_N.$$

When  $t \sim t_j$  or  $t \sim -t_j$ , we use (7.14) to derive

$$\begin{aligned} & -\frac{1}{2\pi i} \int_{\Gamma} \frac{t}{t^2 - t_1^2} \frac{1}{32} \frac{(t^2 - 1)^3}{t^2} \cdot \frac{1}{dt} \otimes dt_1 \\ & \otimes (-1) \mathcal{L}_{g,n-1}(t, t_{N \setminus \{1,j\}}) \left( \frac{1}{(-t+t_j)^2} + \frac{1}{(t+t_j)^2} \right) dt^{\otimes 2} \otimes dt_{N \setminus \{1\}} \\ & = -\frac{1}{32} \frac{\partial}{\partial t_j} \left[ \frac{t_j}{t_j^2 - t_1^2} \frac{(t_j^2 - 1)^3}{t_j^2} \mathcal{L}_{g,n-1}(t_j, t_{N \setminus \{1,j\}}) \right] dt_N \\ & - \frac{1}{32} \left( -\frac{\partial}{\partial t_j} \right) \left[ \frac{-t_j}{t_j^2 - t_1^2} \frac{(t_j^2 - 1)^3}{t_j^2} \mathcal{L}_{g,n-1}(-t_j, t_{N \setminus \{1,j\}}) \right] dt_N \\ & = \frac{1}{16} \frac{\partial}{\partial t_j} \left[ \frac{t_j}{t_1^2 - t_j^2} \frac{(t_j^2 - 1)^3}{t_j^2} \mathcal{L}_{g,n-1}(t_j, t_{N \setminus \{1,j\}}) \right] dt_N. \end{aligned}$$

This completes the proof.  $\square$

Here we note that the spectral curve (7.11), and hence the topological recursion theory of our case, has a non-trivial automorphism. It is given by the transformation

$$(7.17) \quad t \longmapsto \frac{1}{t},$$

which induces an automorphism

$$(7.18) \quad C \ni (x, y) \longmapsto (-x, -y) \in C$$



of the spectral curve. It interchanges the two ramification points of Figure 7.1. Let

$$u = \frac{1}{t}, \quad u_j = \frac{1}{t_j} \quad \text{for } j = 1, 2, \dots, n.$$

Then we have

$$\mathcal{L}_{0,2}(t, t_j) dt \otimes dt_j = \mathcal{L}_{0,2}(u, u_j) du \otimes du_j,$$

and  $ydx = (-y)d(-x)$ . It follows that  $K(t, t_1) = K(u, u_1)$ , and we have  $\mathbb{Z}/2\mathbb{Z}$  as the automorphism group of the theory. Reflecting this automorphism, the function  $\mathcal{L}_{g,n}(t_N)$  exhibits the following transformation property:

$$(7.19) \quad \mathcal{L}_{g,n} \left( \frac{1}{t_1}, \dots, \frac{1}{t_n} \right) = (-1)^n \mathcal{L}_{g,n}(t_1, \dots, t_n) t_1^2 \cdots t_n^2.$$

The reason that we choose  $t$  as our preferred parameter rather than  $1/t$  in (4.2) is to extract the polynomial behavior of the Laplace transform of the Euclidean volume. As  $t \rightarrow \infty$  the spectral curve  $C$  degenerates to a parabola, and the theory changes from counting the integral ribbon graphs to calculating the Euclidean volume, as we shall see below. By the symmetry argument, the  $t \rightarrow 0$  limit also deforms  $C$  to a parabola. We can see from (4.2) that

$$t \rightarrow \infty \quad \iff \quad e^{-w} = \frac{t+1}{t-1} \rightarrow 1 \quad \iff \quad w \rightarrow 0,$$

and the  $w \rightarrow 0$  behavior of the Laplace transform represents the Euclidean volume function, as explained in Section 5. Even though there is a symmetry in the  $t$ -variables, in terms of  $w$ , we have

$$t \rightarrow 0 \quad \iff \quad e^{-w} = \frac{t+1}{t-1} \rightarrow -1,$$

and this limit does not correspond to bringing the mesh of the lattice to 0.

By restricting (7.15) to the top degree terms using

$$\frac{(t^2 - 1)^3}{t^2} = t^4 - 3t^2 + 3 - \frac{1}{t^2},$$

we obtain the recursion for the Euclidean volume.

**Theorem 7.4.** *Define the Eynard kernel for the Euclidean volume by*

$$(7.20) \quad K_E(t, t_1) = -\frac{1}{2} \left( \frac{1}{t-t_1} + \frac{1}{t+t_1} \right) \frac{1}{32} t^4 \cdot \frac{1}{dt} \otimes dt_1$$

on the spectral curve  $C_E$  defined by the parametrization

$$(7.21) \quad \begin{cases} x - 2 = \frac{4}{t^2} \\ y - 1 = \frac{2}{t} \end{cases}.$$

Then the Laplace transformed Euclidean volume function  $V_{g,n}^E(t_N)$  satisfies an Eynard-Orantin type recursion

$$(7.22) \quad \begin{aligned} V_{g,n}^E(t_N) &= -\frac{1}{2\pi i} \oint K_E(t, t_1) \left[ \sum_{j=2}^n \left( V_{g,n-1}^E(t, t_{N \setminus \{1,j\}}) dt \otimes dt_{N \setminus \{1,j\}} \otimes \frac{ds(t) \otimes dt_j}{(s(t) + t_j)^2} \right. \right. \\ &\quad \left. \left. + V_{g,n-1}^E(s(t), t_{N \setminus \{1,j\}}) ds(t) \otimes dt_{N \setminus \{1,j\}} \otimes \frac{dt \otimes dt_j}{(t + t_j)^2} \right) \right] \end{aligned}$$

$$\begin{aligned}
& + V_{g-1, n+1}^E(t, s(t), t_{N \setminus \{1\}}) dt \otimes ds(t) \otimes dt_{N \setminus \{1\}} \\
& + \sum_{\substack{\text{stable} \\ g_1 + g_2 = g \\ I \sqcup J = N \setminus \{1\}}} \left( V_{g_1, |I|+1}^E(t, t_I) dt \otimes dt_I \right) \otimes \left( V_{g_2, |J|+1}^E(s(t), t_J) ds(t) \otimes dt_J \right) \Big].
\end{aligned}$$

Here the integration contour is a positively oriented circle of large radius.

The geometry behind the recursion formula (7.22) is the following. The Euclidean volume is obtained by extracting the asymptotic behavior of  $\mathcal{L}_{g,n}(t_N)$  as  $t \rightarrow \infty$ . The parameterization

$$\begin{cases} x = 2 + \frac{4}{t^2-1} \\ y = 1 + \frac{2}{t-1} \end{cases}$$

of the spectral curve (7.11) of Figure 7.1 near  $t \sim \infty$  gives a neighborhood of one of the critical points  $(x, y) = (2, 1)$ . Thus we *define* a new spectral curve  $C_E$  by the parametrization (7.21), which is simply a parabola  $x - 2 = (y - 1)^2$ . The deck-transformation of the  $x$ -projection of the parabola  $C_E$  is still given by  $t \mapsto s(t) = -t$ . The recipe of (7.9) then gives (7.20), provided that the unstable  $(0, 2)$  geometry still gives the same kernel

$$(7.23) \quad V_{0,2}^E(t_1, t_2) = \frac{dt_1 \otimes dt_2}{(t_1 + t_2)^2}.$$

The continuum limit of (B.7) is

$$v_{0,2}^E(p_1, p_2) = \frac{1}{p_1} \delta(p_1 - p_2).$$

We thus calculate

$$(7.24) \quad \left( \int_0^\infty \int_0^\infty p_1 p_2 v_{0,2}^E(p_1, p_2) e^{-(p_1 w_1 + p_2 w_2)} dp_1 dp_2 \right) dw_1 \otimes dw_2 = \frac{dw_1 \otimes dw_2}{(w_1 + w_2)^2}.$$

The coordinate change (4.2) near  $t \sim \infty$  becomes

$$e^{-w} = \frac{t+1}{t-1} \mapsto 1 - w = 1 + \frac{2}{t},$$

i.e.,  $w = -\frac{2}{t}$ . Under this change, which is an automorphism of  $\mathbb{P}^1$ , (7.24) remains the same, and we obtain (7.23). The  $x$ -projection of the spectral curve  $C_E$  defined by the parametrization (7.21) now has only one ramification point at  $t = \infty$ . Thus the integration contour  $\Gamma$  of (7.15) has changed into a single large circle in (7.22).

The Eynard-Orantin recursion for the symplectic volume is given by the choice of the spectral curve  $C_S$  parametrized by

$$(7.25) \quad \begin{cases} x = \frac{1}{t^2} \\ y = \frac{1}{t} \end{cases}.$$

Since the curve is isomorphic to  $\mathbb{C}$ , we use the same Cauchy differentiation kernel  $W_{02}(t_1, t_2)$  of (7.4) in place of  $V_{0,2}^S(t_1, t_2)$ . The Eynard kernel (7.9) for this case is

$$(7.26) \quad K(t, t_1) = -\frac{1}{2} \left( -\frac{1}{s(t) + t_1} + \frac{1}{t + t_1} \right) \frac{t^4}{4} \cdot \frac{1}{dt} \otimes dt_1.$$

Then the recursion takes exactly the same form of (7.22).

## APPENDIX A. CALCULATION OF THE LAPLACE TRANSFORMS

In this Appendix we prove the Laplace transform formulas used in the main text. We first derive the topological recursion for

$$(A.1) \quad \widehat{L}_{g,n}(w_1, \dots, w_n) = \sum_{\mathbf{p} \in \mathbb{Z}_{\geq 0}^n} p_1 p_2 \cdots p_n N_{g,n}(\mathbf{p}) e^{-\langle p, w \rangle}.$$

Since we multiply the number of integral ribbon graphs by  $p_1 \cdots p_n$ , we can allow all non-negative integers  $p_j$  in the summation, which makes our calculations simpler.

**Proposition A.1.** *The Laplace transform  $\widehat{L}_{g,n}(w_N)$  satisfies the following topological recursion.*

$$(A.2) \quad \begin{aligned} & \widehat{L}_{g,n}(w_N) \\ &= \sum_{j=2}^n \frac{\partial}{\partial w_j} \left[ \left( \frac{e^{w_1}}{e^{w_1} - e^{w_j}} - \frac{e^{w_1+w_j}}{e^{w_1+w_j} - 1} \right) \left( \frac{\widehat{L}_{g,n-1}(w_{N \setminus \{j\}})}{(e^{w_1} - e^{-w_1})^2} - \frac{\widehat{L}_{g,n-1}(w_{N \setminus \{1\}})}{(e^{w_j} - e^{-w_j})^2} \right) \right] \\ & \quad + \frac{1}{(e^{w_1} - e^{-w_1})^2} \left[ \widehat{L}_{g-1,n+1}(w_1, w_1, w_{N \setminus \{1\}}) \right. \\ & \quad \left. + \sum_{\substack{\text{stable} \\ g_1+g_2=g \\ I \sqcup J = N \setminus \{1\}}} \widehat{L}_{g_1,|I|+1}(w_1, w_I) \widehat{L}_{g_2,|J|+1}(w_1, w_J) \right]. \end{aligned}$$

*Proof.* First we multiply both sides of (3.3) by  $p_2 p_3 \cdots p_n$  and compute its Laplace transform. The left-hand side gives  $\widehat{L}_{g,n}(w_N)$ .

The first line of the right-hand side is

$$\begin{aligned} & \sum_{j=2}^n \sum_{\mathbf{p} \in \mathbb{Z}_{\geq 0}^n} \sum_{q=0}^{p_1+p_j} p_j \frac{p_1+p_j-q}{2} [qp_2 \cdots \widehat{p}_j \cdots p_n N_{g,n-1}(q, p_{N \setminus \{1,j\}})] e^{-\langle p, w \rangle} \\ &= \sum_{j=2}^n \sum_{q=0}^{\infty} \sum_{p_{N \setminus \{1,j\}} \in \mathbb{Z}_{\geq 0}^{n-2}} [qp_2 \cdots \widehat{p}_j \cdots p_n N_{g,n-1}(q, p_{N \setminus \{1,j\}})] e^{-\langle p_{N \setminus \{1,j\}}, w_{N \setminus \{1,j\}} \rangle} e^{-qw_1} \\ & \quad \times \sum_{\ell=0}^{\infty} \ell e^{-2\ell w_1} \sum_{p_j=0}^{q+2\ell} p_j e^{p_j(w_1-w_j)}, \end{aligned}$$

where the symbol  $\widehat{\phantom{x}}$  indicates omission of the variable, and we set  $p_1 + p_j - q = 2\ell$ . Note that  $N_{g,n}(p_N) = 0$  unless  $p_1 + \cdots + p_n$  is even. Therefore, in the Laplace transform we are summing over all  $p_N \in \mathbb{Z}_{\geq 0}^n$  such that  $p_1 + \cdots + p_n \equiv 0 \pmod{2}$ . Since  $N_{g,n-1}(q, p_{N \setminus \{1,j\}}) = 0$  unless  $q + p_2 + \cdots + \widehat{p}_j + \cdots + p_n \equiv 0 \pmod{2}$ , only those  $p_1, p_j$  and  $q$  satisfying  $p_1 + p_j - q \equiv 0 \pmod{2}$  contribute in the summation. We now calculate from the last factor (the  $p_j$ -summation)

$$\begin{aligned} & \sum_{p_j=0}^{q+2\ell} e^{-qw_1} \ell e^{-2\ell w_1} p_j e^{p_j(w_1-w_j)} = e^{-qw_1} \ell e^{-2\ell w_1} \frac{\partial}{\partial w_1} \frac{e^{w_j} - e^{w_j} e^{(1+q+2\ell)(w_1-w_j)}}{e^{w_j} - e^{w_1}} \\ &= \frac{e^{-qw_1} \ell e^{-2\ell w_1}}{(e^{w_1} - e^{w_j})^2} \left[ e^{w_1+w_j} + (q+2\ell) e^{2w_1} e^{(q+2\ell)(w_1-w_j)} - (1+q+2\ell) e^{w_1+w_j} e^{(q+2\ell)(w_1-w_j)} \right] \end{aligned}$$

$$= \frac{1}{(e^{w_1} - e^{w_j})^2} \left[ e^{w_1+w_j} e^{-qw_1} \ell e^{-2\ell w_1} + \ell(q+2\ell) e^{2w_1} e^{-qw_j} e^{-2\ell w_j} \right. \\ \left. - \ell(1+q+2\ell) e^{w_1+w_j} e^{-qw_j} e^{-2\ell w_j} \right]$$

followed by the  $\ell$ -summation and then the  $q$ -summation. We obtain

$$= \sum_{j=2}^n \left[ \frac{e^{w_1+w_j}}{(e^{w_1} - e^{w_j})^2} \left( \frac{\widehat{L}_{g,n-1}(w_{N \setminus \{j\}})}{(e^{w_1} - e^{-w_1})^2} - \frac{\widehat{L}_{g,n-1}(w_{N \setminus \{1\}})}{(e^{w_j} - e^{-w_j})^2} \right) \right. \\ \left. - \frac{e^{w_1}}{e^{w_1} - e^{w_j}} \frac{\partial}{\partial w_j} \frac{\widehat{L}_{g,n-1}(w_{N \setminus \{1\}})}{(e^{w_j} - e^{-w_j})^2} \right].$$

The second line of (3.3) contributes

$$\sum_{j=2}^n \sum_{\mathbf{p} \in \mathbb{Z}_{\geq 0}^n} H(p_1 - p_j) \sum_{q=0}^{p_1-p_j} p_j \frac{p_1 - p_j - q}{2} [qp_2 \cdots \widehat{p}_j \cdots p_n N_{g,n-1}(q, p_{N \setminus \{1,j\}})] e^{-\langle \mathbf{p}, w \rangle} \\ = \sum_{j=2}^n \sum_{\ell=0}^{\infty} \ell e^{-2\ell w_1} \sum_{p_j=0}^{\infty} p_j e^{-p_j(w_1+w_j)} \\ \times \sum_{q=0}^{\infty} e^{-qw_1} \sum_{p_{N \setminus \{1,j\}} \in \mathbb{Z}_{\geq 0}^{n-2}} [qp_2 \cdots \widehat{p}_j \cdots p_n N_{g,n-1}(q, p_{N \setminus \{1,j\}})] e^{-\langle p_{N \setminus \{1,j\}}, w_{N \setminus \{1,j\}} \rangle} \\ = \sum_{j=2}^n \frac{e^{w_1+w_j}}{(1 - e^{w_1+w_j})^2} \frac{\widehat{L}_{g,n-1}(w_{N \setminus \{j\}})}{(e^{w_1} - e^{-w_1})^2}.$$

In this calculation we set  $p_1 - p_j - q = 2\ell$ . Similarly, after putting  $p_j - p_1 - q = 2\ell$ , the third line of (3.3) yields

$$- \sum_{j=2}^n \sum_{\mathbf{p} \in \mathbb{Z}_{\geq 0}^n} H(p_j - p_1) \sum_{q=0}^{p_j-p_1} p_j \frac{p_j - p_1 - q}{2} [qp_2 \cdots \widehat{p}_j \cdots p_n N_{g,n-1}(q, p_{N \setminus \{1,j\}})] e^{-\langle \mathbf{p}, w \rangle} \\ = - \sum_{j=2}^n \sum_{q=0}^{\infty} \sum_{\ell=0}^{\infty} \sum_{p_1=0}^{\infty} (p_1 + q + 2\ell) \ell e^{-p_1(w_1+w_j)} e^{-2\ell w_j} e^{-qw_j} \\ \times \sum_{p_{N \setminus \{1,j\}} \in \mathbb{Z}_{\geq 0}^{n-2}} [qp_2 \cdots \widehat{p}_j \cdots p_n N_{g,n-1}(q, p_{N \setminus \{1,j\}})] e^{-\langle p_{N \setminus \{1,j\}}, w_{N \setminus \{1,j\}} \rangle} \\ = - \sum_{j=2}^n \frac{e^{w_1+w_j}}{(1 - e^{w_1+w_j})^2} \frac{\widehat{L}_{g,n-1}(w_{N \setminus \{1\}})}{(e^{w_j} - e^{-w_j})^2} + \sum_{j=2}^n \frac{e^{w_1}}{e^{w_1} - e^{-w_j}} \frac{\partial}{\partial w_j} \frac{\widehat{L}_{g,n-1}(w_{N \setminus \{1\}})}{(e^{w_j} - e^{-w_j})^2}.$$

Summing all contributions, we obtain

$$\sum_{j=2}^n \left[ \frac{e^{w_1+w_j}}{(e^{w_1} - e^{w_j})^2} \left( \frac{\widehat{L}_{g,n-1}(w_{N \setminus \{j\}})}{(e^{w_1} - e^{-w_1})^2} - \frac{\widehat{L}_{g,n-1}(w_{N \setminus \{1\}})}{(e^{w_j} - e^{-w_j})^2} \right) \right.$$

$$\begin{aligned}
& \left. - \frac{e^{w_1}}{e^{w_1} - e^{w_j}} \frac{\partial}{\partial w_j} \frac{\widehat{L}_{g,n-1}(w_{N \setminus \{1\}})}{(e^{w_j} - e^{-w_j})^2} \right] \\
& + \sum_{j=2}^n \frac{e^{w_1+w_j}}{(1 - e^{w_1+w_j})^2} \frac{\widehat{L}_{g,n-1}(w_{N \setminus \{j\}})}{(e^{w_1} - e^{-w_1})^2} \\
& - \sum_{j=2}^n \frac{e^{w_1+w_j}}{(1 - e^{w_1+w_j})^2} \frac{\widehat{L}_{g,n-1}(w_{N \setminus \{1\}})}{(e^{w_j} - e^{-w_j})^2} + \sum_{j=2}^n \frac{e^{w_1}}{e^{w_1} - e^{-w_j}} \frac{\partial}{\partial w_j} \frac{\widehat{L}_{g,n-1}(w_{N \setminus \{1\}})}{(e^{w_j} - e^{-w_j})^2} \\
& = \sum_{j=2}^n \frac{\partial}{\partial w_j} \left[ \left( \frac{e^{w_1}}{e^{w_1} - e^{w_j}} - \frac{e^{w_1+w_j}}{e^{w_1+w_j} - 1} \right) \left( \frac{\widehat{L}_{g,n-1}(w_{N \setminus \{j\}})}{(e^{w_1} - e^{-w_1})^2} - \frac{\widehat{L}_{g,n-1}(w_{N \setminus \{1\}})}{(e^{w_j} - e^{-w_j})^2} \right) \right].
\end{aligned}$$

To compute the Laplace transform of the fourth line of (3.3), we note that

$$\begin{aligned}
\frac{1}{2} \sum_{p_1=0}^{\infty} \sum_{0 \leq q_1+q_2 \leq p_1} q_1 q_2 (p_1 - q_1 - q_2) e^{-p_1 w_1} f(q_1, q_2) \\
= \frac{1}{2} \sum_{q_1=0}^{\infty} \sum_{q_2=0}^{\infty} \sum_{\ell=0}^{\infty} 2\ell e^{-2\ell w_1} e^{-(q_1+q_2)w_1} q_1 q_2 f(q_1, q_2) \\
= \frac{1}{(e^{w_1} - e^{-w_1})^2} \widehat{f}(w_1, w_1),
\end{aligned}$$

where we set  $p_1 - q_1 - q_2 = 2\ell$ , and

$$\widehat{f}(w_1, w_2) = \sum_{q_1=0}^{\infty} \sum_{q_2=0}^{\infty} q_1 q_2 f(q_1, q_2) e^{-(q_1 w_1 + q_2 w_2)}.$$

The reason that  $p_1 - q_1 - q_2$  is even comes from the fact that we are summing over  $p_N \in \mathbb{Z}_{\geq 0}^n$  subject to  $p_1 + \dots + p_n \equiv 0 \pmod{2}$ , while on the fourth line of (3.3) contributions vanish unless  $q_1 + q_2 + p_2 + \dots + p_n \equiv 0 \pmod{2}$ . Therefore, we can restrict the summation over those  $p_1, q_1$  and  $q_2$  subject to  $p_1 \equiv q_1 + q_2 \pmod{2}$ . The calculation of the Laplace transform then becomes straightforward, and the contribution is as in (A.2).

To change from the  $w$ -coordinates to the  $t$ -coordinates, we use (4.2) to find

$$dw_j = \frac{2}{t_j^2 - 1} dt_j, \quad \frac{\partial}{\partial w_j} = \frac{t_j^2 - 1}{2} \frac{\partial}{\partial t_j}.$$

Each factor changes as follows:

$$\begin{aligned}
\frac{1}{(e^{w_j} - e^{-w_j})^2} &= \frac{1}{16} \frac{(t_j^2 - 1)^2}{t_j^2} \\
\frac{e^{w_1}}{e^{w_1} - e^{w_j}} - \frac{e^{w_1+w_j}}{e^{w_1+w_j} - 1} &= \frac{t_j(t_1^2 - 1)}{t_1^2 - t_j^2} \\
\widehat{L}_{g,n}(w_1(t), \dots, w_n(t)) &= (-1)^n 2^{-n} \mathcal{L}_{g,n}(t_1, \dots, t_n) (t_1^2 - 1) \cdots (t_n^2 - 1).
\end{aligned}$$

We can now convert (A.2) to (4.6) by a straightforward calculation.  $\square$

We now prove Theorem 6.2.

**Theorem A.2.** *The symmetric function  $\widehat{V}_{g,n}^S(w_N)$  defined by the Laplace transform*

$$\widehat{V}_{g,n}^S(w_1, \dots, w_n) dw_1 \otimes \dots \otimes dw_n = d_1 \otimes \dots \otimes d_n \int_{\mathbb{R}_+^n} v_{g,n}^S(\mathbf{p}) e^{-\langle w, \mathbf{p} \rangle} dp_1 \dots dp_n$$

*satisfies the topological recursion*

$$(A.3) \quad \widehat{V}_{g,n}^S(w_N) = -2 \sum_{j=2}^{\infty} \frac{\partial}{\partial w_j} \left[ \frac{w_j}{w_1^2 - w_j^2} \left( \frac{\widehat{V}_{g,n-1}^S(w_{N \setminus \{j\}})}{w_1^2} - \frac{\widehat{V}_{g,n-1}^S(w_{N \setminus \{1\}})}{w_j^2} \right) \right] \\ - \frac{2}{w_1^2} \left( \widehat{V}_{g-1,n+1}^S(w_1, w_1, w_{N \setminus \{1\}}) + \sum_{\substack{g_1+g_2=g, \\ I \sqcup J = N \setminus \{1\}}} \widehat{V}_{g_1,|I|+1}^S(w_1, w_I) \widehat{V}_{g_2,|J|+1}^S(w_1, w_J) \right).$$

*Proof.* Since

$$\widehat{V}_{g,n}^S(w_N) = (-1)^n \int_{\mathbb{R}_+^n} p_1 \dots p_n v_{g,n}^S(\mathbf{p}) e^{-\langle w, \mathbf{p} \rangle} dp_1 \dots dp_n,$$

we multiply both sides of (6.6) by  $(-1)^n p_2 \dots p_n$  and take the Laplace transform. The left-hand side gives  $\widehat{V}_{g,n}^S(w_N)$ .

For a continuous function  $f(q)$ , by putting  $p_1 + p_j - q = \ell$ , we have

$$(A.4) \quad \int_0^{\infty} dp_1 \int_0^{\infty} dp_j \int_0^{p_1+p_j} dq p_j q (p_1 + p_j - q) f(q) e^{-(p_1 w_1 + p_j w_j)} \\ = \int_0^{\infty} dq \int_0^{\infty} d\ell \int_0^{q+\ell} dp_j q \ell f(q) e^{-q w_1} e^{-\ell w_1} p_j e^{p_j (w_1 - w_j)} \\ = \frac{1}{(w_1 - w_j)^2} \int_0^{\infty} dq \int_0^{\infty} d\ell q \ell f(q) \left[ e^{-(q+\ell)w_1} - e^{-(q+\ell)w_j} + (q+\ell)(w_1 - w_j) e^{-(q+\ell)w_j} \right] \\ = \frac{1}{(w_1 - w_j)^2} \left( \frac{\widehat{f}(w_1)}{w_1^2} - \frac{\widehat{f}(w_j)}{w_j^2} \right) - \frac{1}{w_1 - w_j} \frac{\partial}{\partial w_j} \left( \frac{\widehat{f}(w_j)}{w_j^2} \right) \\ = \frac{\partial}{\partial w_j} \left[ \frac{1}{w_1 - w_j} \left( \frac{\widehat{f}(w_1)}{w_1^2} - \frac{\widehat{f}(w_j)}{w_j^2} \right) \right],$$

where  $\widehat{f}(w) = \int_0^{\infty} q f(q) e^{-qw} dq$ . By setting  $p_1 - p_j - q = \ell$  we calculate

$$(A.5) \quad \int_0^{\infty} dp_1 \int_0^{\infty} dp_j H(p_1 - p_j) \int_0^{p_1 - p_j} dq p_j q (p_1 - p_j - q) f(q) e^{-(p_1 w_1 + p_j w_j)} \\ = \int_0^{\infty} dq \int_0^{\infty} d\ell \int_0^{\infty} dp_j q \ell f(q) e^{-q w_1} e^{-\ell w_1} p_j e^{-p_j (w_1 + w_j)} = \frac{1}{(w_1 + w_j)^2} \frac{\widehat{f}(w_1)}{w_1^2},$$

and similarly,

$$(A.6) \quad - \int_0^{\infty} dp_1 \int_0^{\infty} dp_j H(p_j - p_1) \int_0^{p_j - p_1} dq p_j q (p_j - p_1 - q) f(q) e^{-(p_1 w_1 + p_j w_j)} \\ = - \int_0^{\infty} dq \int_0^{\infty} d\ell \int_0^{\infty} dp_1 q \ell f(q) e^{-q w_j} e^{-\ell w_j} (p_1 + q + \ell) e^{-p_1 (w_1 + w_j)} \\ = - \int_0^{\infty} dq \int_0^{\infty} d\ell q \ell f(q) e^{-q w_j} e^{-\ell w_j} \left[ \frac{1}{(w_1 + w_j)^2} + \frac{q + \ell}{w_1 + w_j} \right]$$

$$= -\frac{1}{(w_1 + w_j)^2} \frac{\widehat{f}(w_j)}{w_j^2} + \frac{1}{w_1 + w_j} \frac{\partial}{\partial w_j} \left( \frac{\widehat{f}(w_j)}{w_j^2} \right).$$

Adding (A.5) and (A.6) we obtain

$$\begin{aligned} & \int_0^\infty dp_1 \int_0^\infty dp_j H(p_1 - p_j) \int_0^{p_1 - p_j} dq p_j q (p_1 - p_j - q) f(q) e^{-(p_1 w_1 + p_j w_j)} \\ & - \int_0^\infty dp_1 \int_0^\infty dp_j H(p_j - p_1) \int_0^{p_j - p_1} dq p_j q (p_j - p_1 - q) f(q) e^{-(p_1 w_1 + p_j w_j)} \\ & = \frac{1}{(w_1 + w_j)^2} \left( \frac{\widehat{f}(w_1)}{w_1^2} - \frac{\widehat{f}(w_j)}{w_j^2} \right) + \frac{1}{w_1 + w_j} \frac{\partial}{\partial w_j} \left( \frac{\widehat{f}(w_j)}{w_j^2} \right) \\ & = -\frac{\partial}{\partial w_j} \left[ \frac{1}{w_1 + w_j} \left( \frac{\widehat{f}(w_1)}{w_1^2} - \frac{\widehat{f}(w_j)}{w_j^2} \right) \right]. \end{aligned}$$

The sum of the right-hand sides of (A.4)-(A.6) thus becomes

$$\frac{\partial}{\partial w_j} \left[ \left( \frac{1}{w_1 - w_j} - \frac{1}{w_1 + w_j} \right) \left( \frac{\widehat{f}(w_1)}{w_1^2} - \frac{\widehat{f}(w_j)}{w_j^2} \right) \right].$$

Therefore, the first three lines of (6.6) yield

$$-2 \sum_{j=2}^{\infty} \frac{\partial}{\partial w_j} \left[ \frac{w_j}{w_1^2 - w_j^2} \left( \frac{\widehat{V}_{g,n-1}^S(w_{N \setminus \{j\}})}{w_1^2} - \frac{\widehat{V}_{g,n-1}^S(w_{N \setminus \{1\}})}{w_j^2} \right) \right].$$

For a continuous function  $f(q_1, q_2)$ , we have

$$\begin{aligned} & \int_0^\infty dp_1 \iint_{0 \leq q_1 + q_2 \leq p_1} q_1 q_2 (p_1 - q_1 - q_2) f(q_1, q_2) e^{-p_1 w_1} dq_1 dq_2 \\ & = \int_0^\infty dq_1 \int_0^\infty dq_2 \int_0^\infty d\ell q_1 q_2 \ell f(q_1, q_2) e^{-\ell w_1} e^{-(q_1 + q_2) w_1} = \frac{\widehat{f}(w_1, w_1)}{w_1^2}, \end{aligned}$$

where  $\widehat{f}(w_1, w_2) = \int_{\mathbb{R}_+^2} p_1 p_2 f(p_1, p_2) e^{-(p_1 w_1 + p_2 w_2)} dp_1 dp_2$ . Thus the last two lines of (6.6) give

$$-\frac{2}{w_1^2} \left( \widehat{V}_{g-1, n+1}^S(w_1, w_1, w_{N \setminus \{1\}}) + \sum_{\substack{g_1 + g_2 = g, \\ I \sqcup J = N \setminus \{1\}}} \widehat{V}_{g_1, |I|+1}^S(w_1, w_I) \widehat{V}_{g_2, |J|+1}^S(w_1, w_J) \right).$$

This completes the proof of (A.3).  $\square$

Let us now change the coordinates from  $w_j$ 's to  $t_j$ 's that are given by

$$w_j = -\frac{2}{t_j}$$

this time. This change of coordinate gives

$$dw_j = \frac{2}{t_j^2} dt_j, \quad \frac{\partial}{\partial w_j} = \frac{t_j^2}{2} \frac{\partial}{\partial t_j}.$$

Thus the relation in terms of symmetric differential form is

$$V_{g,n}^S(t_N)dt_N = \widehat{V}_{g,n}^S(w_N)dw_N,$$

or

$$V_{g,n}^S(t_N) = 2^n \frac{\widehat{V}_{g,n}^S(w_N)}{t_1^2 \cdots t_n^2}.$$

So we multiply both sides of (A.3) by  $\frac{2^n}{t_1^2 \cdots t_n^2}$ . From the first term of the first line we obtain

$$\begin{aligned} & -2 \frac{2^n}{t_1^2 \cdots t_n^2} \sum_{j=2}^{\infty} \frac{\partial}{\partial w_j} \left[ \frac{w_j}{w_1^2 - w_j^2} \frac{\widehat{V}_{g,n-1}^S(w_{N \setminus \{j\}})}{w_1^2} \right] \\ &= -2 \sum_{j=2}^{\infty} \frac{\partial}{\partial t_j} \left[ \frac{1}{2} \frac{t_1^2 t_j}{t_1^2 - t_j^2} \frac{t_1^2}{4} V_{g,n-1}^S(t_{N \setminus \{j\}}) \right] = -\frac{1}{4} \sum_{j=2}^{\infty} \frac{\partial}{\partial t_j} \left[ \frac{t_j}{t_1^2 - t_j^2} t_1^4 V_{g,n-1}^S(t_{N \setminus \{j\}}) \right]. \end{aligned}$$

Similarly, the second term of the first line becomes

$$\begin{aligned} & 2 \frac{2^n}{t_1^2 \cdots t_n^2} \sum_{j=2}^{\infty} \frac{\partial}{\partial w_j} \left[ \frac{w_j}{w_1^2 - w_j^2} \frac{\widehat{V}_{g,n-1}^S(w_{N \setminus \{1\}})}{w_j^2} \right] \\ &= 2 \sum_{j=2}^{\infty} \frac{\partial}{\partial t_j} \left[ \frac{1}{2} \frac{t_1^2 t_j}{t_1^2 - t_j^2} \frac{t_j^2}{4} \frac{t_j^2 V_{g,n-1}^S(t_{N \setminus \{1\}})}{t_1^2} \right] = \frac{1}{4} \sum_{j=2}^{\infty} \frac{\partial}{\partial t_j} \left[ \frac{t_j}{t_1^2 - t_j^2} t_j^4 V_{g,n-1}^S(t_{N \setminus \{1\}}) \right]. \end{aligned}$$

The second line of (A.3) is easy to convert. This completes the proof of Theorem 6.2.

#### APPENDIX B. EXAMPLES

For  $(g, n) = (0, 3)$ , there are three topological shapes of ribbon graphs listed in Figure B.1. Cyclic permutations of  $(p_1, p_2, p_3)$  produce different graphs.





FIGURE B.1. Three ribbon graphs for  $(g, n) = (0, 3)$ .

Which ribbon graph corresponds to a point  $(p_1, p_2, p_3) \in \mathbb{Z}_+^3$  depends on which inequality these three numbers satisfy. If  $p_1 > p_2 + p_3$ , then the dumbbell shape (Figure B.1, bottom left) corresponds to this point. If  $p_1 = p_2 + p_3$ , then the shape of  $\infty$  (Figure B.1, bottom right) corresponds, and if no coordinate is greater than the sum of the other two, then the double circle graph (Figure B.1, top) corresponds. These inequalities divide  $\mathbb{Z}_+^3$  into four regions as in Figure B.2.



FIGURE B.2. The partition of  $\mathbb{Z}_+^3$ .

Thus we conclude

$$(B.1) \quad N_{0,3}(p_1, p_2, p_3) = \begin{cases} 1 & p_1 + p_2 + p_3 \equiv 0 \pmod{2}, \\ 0 & \text{otherwise.} \end{cases}$$

The even parity condition can be met if all three are even or only one of them is even. Let us substitute  $p_j = 2q_j$  when it is even and  $p_j = 2q_j - 1$  if it is odd. Thus the Laplace transform can be calculated by

$$\begin{aligned} L_{0,3}(w_1, w_2, w_3) &= \sum_{(p_1, p_2, p_3) \in \mathbb{Z}_+^3} N_{0,3}(p_1, p_2, p_3) e^{-(p_1 w_1 + p_2 w_2 + p_3 w_3)} \\ &= \sum_{(q_1, q_2, q_3) \in \mathbb{Z}_+^3} \left( 1 + e^{w_1 + w_2} + e^{w_2 + w_3} + e^{w_3 + w_1} \right) e^{-2(q_1 w_1 + q_2 w_2 + q_3 w_3)} \\ &= \left( 1 + e^{w_1 + w_2} + e^{w_2 + w_3} + e^{w_3 + w_1} \right) \frac{e^{-(w_1 + w_2 + w_3)}}{(e^{w_1} - e^{-w_1})(e^{w_2} - e^{-w_2})(e^{w_3} - e^{-w_3})}. \end{aligned}$$

Using  $e^{-w_j} = \frac{t_j + 1}{t_j - 1}$ , we obtain

$$(B.2) \quad L_{0,3}(w(t_1), w(t_2), w(t_3)) = -\frac{1}{16} (t_1 + 1)(t_2 + 1)(t_3 + 1) \left( 1 + \frac{1}{t_1 t_2 t_3} \right)$$

and

$$(B.3) \quad \mathcal{L}_{0,3}(t_1, t_2, t_3) = \frac{\partial^3}{\partial t_1 \partial t_2 \partial t_3} L_{0,3}(w(t_1), w(t_2), w(t_3)) = -\frac{1}{16} \left( 1 - \frac{1}{t_1^2 t_2^2 t_3^2} \right).$$

For  $(g, n) = (1, 1)$ , there are two ribbon graphs (see Figure B.3) corresponding to a hexagonal and a square tiling of the plane. The hexagonal tiling gives a ribbon graph on the left, and the square one on the right is a degeneration obtained by shrinking the horizontal edge to 0. The automorphism group is  $\mathbb{Z}/6\mathbb{Z}$  for the degree 3 graph, and  $\mathbb{Z}/4\mathbb{Z}$  for the degree 4 graph.

The number of integral ribbon graphs in this case is the number of partitions of the half of the given perimeter length  $p = 2q \in 2\mathbb{Z}_+$  into two or three positive integers corresponding to edge lengths. Taking the automorphism factors into account, we calculate

$$N_{1,1}(2q) = \frac{1}{4}(q-1) + \frac{1}{6} \sum_{r=1}^{q-1} (r-1) = \frac{1}{12} (q^2 - 1).$$

FIGURE B.3. Two ribbon graphs of type  $(1, 1)$ .

Therefore,

$$(B.4) \quad N_{1,1}(p) = \begin{cases} \frac{1}{48}(p^2 - 4) & p \equiv 0 \pmod{2}, \\ 0 & \text{otherwise.} \end{cases}$$

The Laplace transform can be calculated immediately:

$$L_{1,1}(w) = \sum_{p=2}^{\infty} N_{1,1}(p)e^{-pw} = \frac{1}{12} \sum_{q=1}^{\infty} (q^2 - 1)e^{-2qw} = \frac{3e^{2w} - 1}{12(e^{2w} - 1)^3}.$$


We thus obtain

$$(B.5) \quad L_{1,1}(w(t)) = -\frac{1}{384} \frac{(t+1)^4}{t^2} \left( t - 4 + \frac{1}{t} \right)$$

and

$$(B.6) \quad \mathcal{L}_{1,1}(t) = -\frac{1}{2^7} \frac{(t^2 - 1)^3}{t^4}.$$

The values of  $(g, n)$  corresponding to genus 0 unstable geometries  $(0, 1)$ ,  $(0, 2)$  play an important role in topological recursion. We have seen this phenomena in Hurwitz theory [14, 36]. Let us consider the unstable  $(0, 2)$  case for the integral ribbon graph counting. Although we have restricted our ribbon graphs to have vertices of degree 3 or more, it is indeed more consistent to allow vertices of degree 2. Actually, a *metric* ribbon graph with integer edge lengths is a plain ribbon graph whose vertices have degree 2 or more. For such a ribbon graph, we assign length 1 to every edge. We recover a metric ribbon graph with integer edge length by disregarding all vertices of degree 2. This point of view is natural when we assign a Belyi morphism [1, 34, 43] to a ribbon graph.

FIGURE B.4. A ribbon graph of type  $(0, 2)$ .

Once we allow degree 2 vertices, there is only one kind of ribbon graph of type  $(0, 2)$ , which is a circle (Figure B.4). If the graph has  $p$  edges, then the automorphism group of this graph is  $\mathbb{Z}/p\mathbb{Z}$ . Therefore, we have

$$(B.7) \quad N_{0,2}(p_1, p_2) = \frac{1}{p_1} \delta_{p_1 p_2},$$

and its Laplace transform becomes

$$L_{0,2}(w_1, w_2) = \sum_{p_1=1}^{\infty} \sum_{p_2=1}^{\infty} N_{0,2}(p_1, p_2) e^{-(p_1 w_1 + p_2 w_2)} = -\log(1 - e^{-w_1 - w_2}).$$

In terms of the  $t$ -coordinates we have

$$(B.8) \quad L_{0,2}(w(t_1), w(t_2)) = -\log\left(1 - \frac{t_1 + 1}{t_1 - 1} \cdot \frac{t_2 + 1}{t_2 - 1}\right),$$

which gives

$$(B.9) \quad \mathcal{L}_{0,2}(t_1, t_2) = \frac{1}{(t_1 + t_2)^2}.$$

Note that (B.9) is not a Laurent polynomial and exhibits an exception to the general statement of Theorem 4.1.

The parametrization of the spectral curve (7.11) defines the  $x$ -projection map

$$\pi : \mathbb{P}^1 \ni t \mapsto x = \frac{t+1}{t-1} + \frac{t-1}{t+1} \in \mathbb{P}^1.$$

We find that the difference of the Cauchy differentiation kernels of the curve upstairs and downstairs is  $\mathcal{L}_{0,2}(t_1, t_2)$ :

$$(B.10) \quad \mathcal{L}_{0,2}(t_1, t_2) dt_1 \otimes dt_2 = \frac{dt_1 \otimes dt_2}{(t_1 + t_2)^2} = \frac{dt_1 \otimes dt_2}{(t_1 - t_2)^2} - \pi^* \frac{dx_1 \otimes dx_2}{(x_1 - x_2)^2},$$

where  $\pi$  is the  $x$ -projection map (7.12). We note that this situation is exactly the same as the Hurwitz theory [14, Remark 4.5].

The other genus 0 unstable case  $(0, 1)$  is tricky because we need to allow degree 1 vertices. Since all possible trees can be included if we allow degree 1 vertices, we have to make a choice as to what kind of trees are allowed. If we allow arbitrary trees, then we are led to the Hurwitz theory of [42]. In the integral ribbon graph case, we allow only a chain of edges of finite length. A chain has automorphism group  $\mathbb{Z}/2\mathbb{Z}$ , and the perimeter is twice the length of the chain. Thus we obtain

$$(B.11) \quad N_{0,1}(p) = \begin{cases} \frac{1}{2} & p \equiv 0 \pmod{2} \\ 0 & \text{otherwise.} \end{cases}$$

We have

$$L_{0,1}(w) = \sum_{p=1}^{\infty} N_{0,1}(p) e^{-pw} = \frac{1}{2} \sum_{q=1}^{\infty} e^{-2qw} = \frac{1}{2} \frac{1}{e^{2w} - 1}.$$

It follows that

$$(B.12) \quad L_{0,1}(w(t)) = -\frac{1}{8} \frac{(t+1)^2}{t},$$

$$(B.13) \quad \mathcal{L}_{0,1}(t) = -\frac{1}{8} \frac{t^2 - 1}{t^2}.$$

Using the recursion formula (4.6), we can calculate  $\mathcal{L}_{g,n}(t_N)$  systematically. A few examples are listed below.

$$(B.14) \quad \mathcal{L}_{0,4}(t_1, t_2, t_3, t_4) = \frac{1}{28} \left[ 3 \sum_{j=1}^4 t_j^2 - 9 - \sum_{i < j} \frac{1}{t_i^2 t_j^2} - \frac{9}{t_1^2 t_2^2 t_3^2 t_4^2} + \frac{3}{t_1^2 t_2^2 t_3^2 t_4^2} \sum_{j=1}^4 \frac{1}{t_j^2} \right].$$

$$(B.15) \quad \mathcal{L}_{1,2}(t_1, t_2) = \frac{1}{2^{11}} \left[ 5(t_1^4 + t_2^4) + 3t_1^2 t_2^2 - 18(t_1^2 + t_2^2) + 27 - 4 \left( \frac{1}{t_1^2} + \frac{1}{t_2^2} \right) + \frac{27}{t_1^2 t_2^2} - \frac{18}{t_1^2 t_2^2} \left( \frac{1}{t_1^2} + \frac{1}{t_2^2} \right) + \frac{3}{t_1^4 t_2^4} + \frac{5}{t_1^2 t_2^2} \left( \frac{1}{t_1^4} + \frac{1}{t_2^4} \right) \right].$$

$$(B.16) \quad \mathcal{L}_{2,1}(t) = -\frac{21}{2^{19}} \frac{(t^2 - 1)^7}{t^8} \left( 5t^2 + 6 + \frac{5}{t^2} \right).$$

$$(B.17) \quad \mathcal{L}_{3,1}(t) = -\frac{11}{2^{30}} \frac{(t^2 - 1)^{11}}{t^{12}} \left( 2275t^4 + 4004t^2 + 4722 + \frac{4004}{t^2} + \frac{2275}{t^4} \right).$$

## REFERENCES

- [1] G. V. Belyi, *On galois extensions of a maximal cyclotomic fields*, Math. U.S.S.R. Izvestija **14**, 247–256 (1980).
- [2] J. Bennett, D. Cochran, B. Safnuk, and K. Woskoff, *Topological recursion for symplectic volumes of moduli spaces of curves*, in preparation.
- [3] G. Borot, B. Eynard, M. Mulase and B. Safnuk, *Hurwitz numbers, matrix models and topological recursion*, arXiv:0906.1206.
- [4] V. Bouchard, A. Klemm, M. Mariño, and S. Pasquetti, *Remodeling the B-model*, Commun. Math. Phys. **287**, 117–178 (2008).
- [5] V. Bouchard and M. Mariño, *Hurwitz numbers, matrix models and enumerative geometry*, Proc. Symposia Pure Math. **78**, 263–283 (2008).
- [6] A. Brini, *The local Gromov-Witten theory of  $\mathbb{C}P^1$  and integrable hierarchies*, arXiv:1002.0582v1 [math-ph].
- [7] L. Chen, *Bouchard-Klemm-Marino-Pasquetti Conjecture for  $\mathbb{C}^3$* , arXiv:0910.3739 (2009).
- [8] R. Dijkgraaf and C. Vafa, *Two Dimensional Kodaira-Spencer Theory and Three Dimensional Chern-Simons Gravity*, arXiv:0711.1932 [hep-th].
- [9] R. Dijkgraaf, E. Verlinde, and H. Verlinde, *Loop equations and Virasoro constraints in non-perturbative two-dimensional quantum gravity*, Nucl. Phys. **B348**, 435–456 (1991).
- [10] B. Dubrovin, *Geometry of 2D topological field theories*, in “Integrable systems and quantum groups,” Lecture Notes in Math. **1620**, 120–348 (1994).
- [11] B. Dubrovin and Y. Zhang, *Frobenius manifolds and Virasoro constraints*, Selecta Mathematica, New Ser. **5**, 423–466 (1999).
- [12] B. Eynard, *Topological expansion for the 1-hermitian matrix model correlation functions*, arXiv:hep-th/0407261.
- [13] B. Eynard, *Recursion between volumes of moduli spaces*, arXiv:0706.4403 [math-ph].
- [14] B. Eynard, M. Mulase and B. Safnuk, *The Laplace transform of the cut-and-join equation and the Bouchard-Mariño conjecture on Hurwitz numbers*, arXiv:0907.5224 math.AG (2009).
- [15] B. Eynard and N. Orantin, *Invariants of algebraic curves and topological expansion*, Commun. Number Theory Phys. **1**, 347–452 (2007).
- [16] B. Eynard and N. Orantin, *Weil-Petersson volume of moduli spaces, Mirzakhani’s recursion and matrix models*, arXiv:0705.3600 [math-ph].
- [17] H. Fan, T. J. Jarvis, Evan Merrell and Y. Ruan, *The Witten equation and its virtual fundamental cycle*, arXiv:0712.4025 (2007).
- [18] H. Fan, T. J. Jarvis and Y. Ruan, *Witten’s  $D_4$  Integrable Hierarchies Conjecture*, arXiv:1008.0927 (2010).
- [19] H. Fan, T. J. Jarvis and Y. Ruan, *The Witten equation and its virtual fundamental cycle*, arXiv:0712.4025 (2007).
- [20] I.P. Goulden and D.M. Jackson, *Transitive factorisations into transpositions and holomorphic mappings on the sphere*, Proc. A.M.S., **125**, 51–60 (1997).
- [21] J. L. Harer, *The cohomology of the moduli space of curves*, in Theory of Moduli, Montecatini Terme, 1985 (Edoardo Sernesi, ed.), Springer-Verlag, 1988, pp. 138–221.
- [22] J. L. Harer and D. Zagier, *The Euler characteristic of the moduli space of curves*, Inventiones Mathematicae **85**, 457–485 (1986).
- [23] Tamás Hausel and Michael Thaddeus, *Mirror symmetry, Langlands duality, and the Hitchin system*, Inventiones Math. **153**, 197–229 (2003).
- [24] L. Jeffrey and F. Kirwan, *Intersection theory on moduli spaces of holomorphic bundles of arbitrary rank on a Riemann surface*, Annals of Math. textbf148, 109–196 (1998).
- [25] L. Jeffrey and J. Weitsman, *Toric structures on the moduli space of flat connections on a Riemann surface: volumes and the moment map*, Advances Math. **106**, 151–168 (1994).
- [26] M. Kazarian, *KP hierarchy for Hodge integrals*, arXiv:0809.3263.

- [27] M. Kazarian, S. Lando, *An algebro-geometric proof of Witten's conjecture*, J. Amer. Math. Soc. **20**, 1079–1089 (2007).
- [28] M. Kontsevich, *Intersection theory on the moduli space of curves and the matrix Airy function*, Commun. Math. Phys. **147**, 1–23 (1992).
- [29] K. Liu, *Heat kernel and moduli space*, Math. Res. Letter **3**, 743–762 (1996).
- [30] K. Liu and H. Xu, *Recursion formulae of higher Weil-Petersson volumes*, International Mathematics Research Notices (2009), No. 5, pp. 835–859.
- [31] M. Mariño, *Open string amplitudes and large order behavior in topological string theory*, arXiv:hep-th/0612127 (2006–2008).
- [32] M. Mirzakhani, *Simple geodesics and Weil-Petersson volumes of moduli spaces of bordered Riemann surfaces*, Invent. Math. **167**, 179–222 (2007).
- [33] M. Mirzakhani, *Weil-Petersson volumes and intersection theory on the moduli space of curves*, J. Amer. Math. Soc. **20**, 1–23 (2007).
- [34] M. Mulase and M. Penkava, *Ribbon graphs, quadratic differentials on Riemann surfaces, and algebraic curves defined over  $\mathbb{Q}$* , The Asian Journal of Mathematics **2** (4), 875–920 (1998).
- [35] M. Mulase and B. Safnuk, *Mirzakhani's recursion relations, Virasoro constraints and the KdV hierarchy*, Indian J. Math. **50**, 189–228 (2008).
- [36] M. Mulase and N. Zhang, *Polynomial recursion formula for linear Hodge integrals*, Communications in Number Theory and Physics **4**, 267–294 (2010).
- [37] D. Mumford, *Towards an enumerative geometry of the moduli space of curves* (1983), in “Selected Papers of David Mumford,” 235–292 (2004).
- [38] P. Norbury, *Counting lattice points in the moduli space of curves*, Mathematics Research Letters, to appear (2010).
- [39] P. Norbury, *String and dilaton equations for counting lattice points in the moduli space of curves*, arXiv:0905.4141 (2009).
- [40] A. Okounkov, *Random matrices and random permutations*, International Mathematics Research Notices **2000**, 1043–1095 (2000).
- [41] A. Okounkov, *Toda equations for Hurwitz numbers*, Math. Res. Lett. **7**, 447 (2000) [arXiv:math.AG/0004128].
- [42] A. Okounkov and R. Pandharipande, *Gromov-Witten theory, Hurwitz numbers, and matrix models, I*, Proc. Symposia Pure Math. **80**, 325–414 (2009).
- [43] L. Schneps and P. Lochak, *Geometric Galois actions*, London Mathematical Society Lecture Notes Series **242**, 1997.
- [44] D. D. Sleator, R. E. Tarjan, and W. P. Thurston, *Rotation distance, triangulations, and hyperbolic geometry*, Journal of the American Mathematical Society **1**, 647–681 (1988).
- [45] Kurt Strebel, *Quadratic differentials*, Springer-Verlag, 1984.
- [46] A. Strominger, S. T. Yau, and E. Zaslow, *Mirror symmetry is T-duality*, Nuclear Phys. **B 479**, 243–259 (1996).
- [47] R. Vakil, Harvard Thesis 1997.
- [48] E. Witten, *Two dimensional gravity and intersection theory on moduli space*, Surveys in Differential Geometry **1**, 243–310 (1991).
- [49] E. Witten, *On quantum gauge theories in two dimensions*, Commun. Math. Physics **141**, 153–209 (1991).
- [50] J. Zhou, *Local Mirror Symmetry for One-Legged Topological Vertex*, arXiv:0910.4320 (2009).
- [51] J. Zhou, *Local Mirror Symmetry for the Topological Vertex* arXiv:0911.2343 (2009).

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, DAVIS, CA 95616–8633, U.S.A.  
*E-mail address:* `kmchapman@math.ucdavis.edu`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, DAVIS, CA 95616–8633, U.S.A.  
*E-mail address:* `mulase@math.ucdavis.edu`

DEPARTMENT OF MATHEMATICS, CENTRAL MICHIGAN UNIVERSITY, MOUNT PLEASANT, MI 48859, U.S.A.  
*E-mail address:* `brad.safnuk@cmich.edu`