

MAT 271: Applied & Computational Harmonic Analysis Supplementary Notes III by Naoki Saito

A Brief History of the Convergence of the Fourier Series

Theorem 1 (Dirichlet, 1829) Suppose f is 1-periodic, piecewise smooth on \mathbb{R} . Then, n th partial sum, $S_n[f](x) \triangleq \sum_{-n}^n c_k e^{2\pi i k x}$, satisfies

$$\lim_{n \rightarrow \infty} S_n[f](x) = \frac{1}{2} [f(x+) + f(x-)].$$

In particular, if x is a point of continuity, then $\lim_{n \rightarrow \infty} S_n[f](x) = f(x)$.

Theorem 2 (du Bois Reymond, 1876) There exists $f \in C(I)$ such that $\{S_n[f](0)\}$ diverges, where I is an interval of unit length.

Theorem 3 (A weak version of Fejér's Theorem) If f is 1-periodic, *continuous*, and piecewise smooth on \mathbb{R} , then the Fourier series of f converges to f *absolutely and uniformly*.

Definition: Suppose a series of functions $\sum_1^\infty g_n(x)$ converges to $g(x)$ on a set $x \in I$. Then, the convergence is called *absolute* if $\sum_1^\infty |g_n(x)|$ also converges for $x \in I$.

If we have $\sup_{x \in I} \left| g(x) - \sum_1^N g_n(x) \right| \rightarrow 0$ as $N \rightarrow \infty$, then we call this a *uniform convergence*.

Theorem 4 (Fejér 1904) If $f \in C(I)$, then the Cesàro means of $S_n[f]$ converge *uniformly* to f .

Definition: The m th *Cesàro mean* of partial sums is the mean of the first $m + 1$ partial sums, i.e., $\sigma_m[f](x) \triangleq \frac{1}{m+1} \sum_{n=0}^m S_n[f](x)$.

Theorem 5 (Size of the Fourier coefficients and the smoothness of the functions) Suppose f is 1-periodic. If $f \in C^{k-1}(\mathbb{R})$ and $f^{(k-1)}$ is piecewise smooth (i.e., $f^{(k)}$ exists and piecewise continuous), then the Fourier coefficients of f , c_n , satisfy $\sum_n |n^k c_n| < \infty$. In particular, $n^k c_n \rightarrow 0$. On the other hand, suppose $c_n, n \neq 0$, satisfy $|c_n| \leq C|n|^{-(k+\gamma)}$ for some $C > 0$ and $\gamma > 1$. Then $f \in C^k(\mathbb{R})$.

Theorem 6 (Kolmogorov, 1926) There exists $f \in L^1(I)$ such that $\{S_n[f](x)\}$ diverges for every x .

Theorem 7 (Carleson, 1966) If $f \in L^2(I)$, then $S_n[f](x)$ converges to $f(x)$ almost everywhere.

Theorem 8 (Hunt, 1967) If $f \in L^p(I), p > 1$, then $S_n[f](x)$ converges to $f(x)$ almost everywhere.

Mathematicians are still trying to simplify the proof of the Carlson-Hunt theorem as of today.

For the details of the above facts, see [1, Chap. 1,2], [2, Chap. 1], [3, Part 1], and [4, Chap. 1]. [5, Chap. 1].

References

- [1] J. M. ASH, ed., *Studies in Harmonic Analysis*, vol. 13 of MAA Studies in Mathematics, Math. Assoc. Amer., 1976.
- [2] H. DYM AND H. P. MCKEAN, *Fourier Series and Integrals*, Academic Press, 1972.
- [3] T. W. KÖRNER, *Fourier Analysis*, Cambridge Univ. Press, 1988.
- [4] S. G. KRANTZ, *A Panorama of Harmonic Analysis*, no. 27 in The Carus Mathematical Monographs, Math. Assoc. Amer., Washington, D.C., 1999.
- [5] M. A. PINSKY, *Introduction to Fourier Analysis and Wavelets*, Brooks/Cole, 2002.