



# *Well-posedness for the Classical Stefan Problem and the Zero Surface Tension Limit*

MAHIR HADŽIĆ, STEVE SHKOLLER

*Communicated by C. DAFERMOS*

## **Abstract**


We develop a framework for a *unified* treatment of well-posedness for the Stefan problem with or without surface tension. In the absence of surface tension, we establish well-posedness in Sobolev spaces for the classical Stefan problem. We introduce a new velocity variable which extends the velocity of the moving free-boundary into the interior domain. The equation satisfied by this velocity is used for the analysis in place of the heat equation satisfied by the temperature. Solutions to the classical Stefan problem are then constructed as the limit of solutions to a carefully chosen sequence of approximations to the velocity equation, in which the moving free-boundary is regularized and the boundary condition is modified in a such a way as to preserve the basic nonlinear structure of the original problem. With our methodology, we simultaneously find the required stability condition for well-posedness and obtain new estimates for the regularity of the moving free-boundary. Finally, we prove that solutions of the Stefan problem with positive surface tension  $\sigma$  converge to solutions of the classical Stefan problem as  $\sigma \rightarrow 0$ .

## **1. Introduction**

### *1.1. The Problem Formulation*

We consider the local well-posedness and boundary regularity of solutions to the classical one-phase Stefan problem, describing the evolving phase boundary of a freezing liquid. We also establish the limit of zero surface tension.

The temperature  $p(t, x)$  of a liquid inside of a time-dependent domain  $\Omega(t)$  and an *a priori unknown* moving boundary  $\Gamma(t)$  satisfies the following system of equations:



**Fig. 1.** The one-phase Stefan problem. Displayed on the *left side* of the figure is the reference domain  $\Omega$  and reference boundary  $\Gamma$ . The time-dependent domain  $\Omega(t)$  and the moving free-boundary  $\Gamma(t)$  is shown on the *right side* of the figure.

$$p_t - \Delta p = 0 \quad \text{in } \Omega(t), \quad (1.1a)$$

$$\partial_n p = V_{\Gamma(t)} \quad \text{on } \Gamma(t), \quad (1.1b)$$

$$p = \sigma \kappa_{\Gamma} \quad \text{on } \Gamma(t), \quad (1.1c)$$

$$p(0, \cdot) = p_0, \quad \Gamma(0) = \Gamma_0. \quad (1.1d)$$

The domain  $\Omega(t)$  is an evolving open subset of  $\mathbb{R}^d$  with  $d \geq 2$ . The set  $\Gamma(t)$  denotes the moving boundary (which may be a connected subset of  $\partial\Omega(t)$  if a part of the boundary of  $\Omega(t)$  is fixed). See Fig. 1.

Equation (1.1a) expresses the fact that heat diffuses in the bulk  $\Omega(t)$ , while the boundary condition (1.1b) states that the heat flux across the boundary governs the boundary evolution; that is,  $\partial_n p = \nabla p \cdot n$  is the normal derivative of  $p$  on  $\Gamma(t)$  where  $n$  stands for the outward pointing unit normal, and  $V_{\Gamma(t)}$  denotes the speed or the normal velocity of the hypersurface  $\Gamma(t)$ . In the case that  $\sigma = 0$ , (1.1c) is termed the *classical Stefan condition* and problem (1.1) is called the *classical Stefan problem*. In this case, freezing of the liquid occurs at a constant temperature  $p = 0$ . On the other hand, if  $\sigma > 0$  in (1.1c) then the boundary condition is called the *Gibbs–Thomson correction* to the classical Stefan condition, and the system (1.1) is then termed the *Stefan problem with surface tension*, whereby  $\sigma > 0$  is a given coefficient of surface tension and  $\kappa_{\Gamma(t)}$  stands for the mean curvature of the moving boundary  $\Gamma(t)$ . Finally, we equip the problem with suitable initial conditions (1.1d):  $p_0 : \Omega(0) \rightarrow \mathbb{R}$  and  $\Gamma_0$  are the prescribed initial temperature and boundary, respectively.

Problem (1.1) is an example of a free-boundary partial differential equation which requires the initial data to satisfy a *stability condition* in order to ensure well-posedness in Sobolev spaces; specifically, we shall require that

$$-\partial_n p_0 > 0 \quad \text{on } \Gamma(0),$$

which, by analogy to fluid dynamics, we shall refer to as the *Taylor sign condition* or the *Rayleigh–Taylor sign condition*. Below, we will explain how this Taylor sign condition naturally appears from our analysis.

### 1.2. The Reference Domain $\Omega$ and the Initial Domain $\Omega_0$

We will begin the analysis with motion in  $\mathbb{R}^2$ , and then describe the minor modifications needed to study motion in  $\mathbb{R}^3$ . To simplify our presentation, we will parameterize our initial free-boundary  $\Gamma_0$  as a graph over the one-dimensional torus  $\mathbb{T}^1$  which we identify with  $[0, 2\pi]$ ; we define

$$\Gamma_0 = \{\mathbf{x} \in \mathbb{T}^1 \times \mathbb{R}, \mathbf{x} = (x', h_0(x'))\}, \quad h_0 \in H^4(\mathbb{T}^1). \quad (1.2)$$

Without loss of generality we shall further assume that  $\Gamma_0$  is a small perturbation of the manifold  $\mathbb{T}^1 \times \{x^2 = 0\}$  in the sense that

$$\|h_0\|_{H^4(\mathbb{T}^1)} \leq \epsilon_0 \ll 1, \quad (1.3)$$

for some sufficiently small  $\epsilon_0$ . In Appendix A, we shall explain how to remove the assumption (1.3). The only reason for making this smallness assumption is that (1.3) and (1.2) allow us to use *one* global Cartesian coordinate system (rather than a collection of local coordinate charts). This is ideal for describing new identities that provide very natural estimates for the second-fundamental form of the evolving free-boundary  $\Gamma(t)$ . All of our results apply to general domains; however, in a general setting, we must employ a finite covering of  $\Omega$  by local coordinate charts, together with a partition-of-unity subordinate to that cover. In particular, the Stefan problem *localizes* to each chart and effectively reduces to the analysis on

$$\Omega = \mathbb{T}^1 \times (0, 1).$$

Again, we emphasize that the assumption (1.3) is not essential to our proof, and in Appendix A, we explain how to treat general  $H^4$  initial geometries.

We define the *initial domain*

$$\Omega_0 = \{(x_1, x_2) \in \mathbb{T}^1 \times \mathbb{R} \mid h_0(x^1) < x^2 < 1\},$$

while the *reference domain* is  $\Omega = \mathbb{T}^1 \times (0, 1)$ . The set

$$\Gamma = \mathbb{T}^1 \times \{x^2 = 0\}$$

is the *reference boundary* on which our parameterization  $(x', h(t, x'))$  will be defined. The *top boundary*  $\partial\Omega_{\text{top}} = \mathbb{T}^1 \times \{x^2 = 1\}$  is fixed in time, and

$$\partial_n p = 0 \quad \text{on} \quad \partial\Omega_{\text{top}}. \quad (1.4)$$

### 1.3. Notation

For any  $s \geq 0$  and given functions  $f : \Omega \rightarrow \mathbb{R}$ ,  $\varphi : \Gamma \rightarrow \mathbb{R}$  we set

$$\|f\|_s := \|f\|_{H^s(\Omega)}; \quad |\varphi|_s := \|\varphi\|_{H^s(\Gamma)}.$$

In particular, when  $s$  is not an integer, the corresponding fractional Sobolev space is defined by interpolation in a standard way. If  $f : [0, T] \times \Omega \rightarrow \mathbb{R}$ ,  $\varphi : [0, T] \times \Gamma \rightarrow \mathbb{R}$  are given time-dependent functions, then for any  $1 \leq p \leq \infty$  we set

$$\|f\|_{L_t^p H_x^s} := \|f\|_{L^p([0, T]; H^s(\Omega))}; \quad |\varphi|_{L_t^p H_x^s} := \|\varphi\|_{L^p([0, T]; H^s(\Gamma))}.$$

If  $i = 1, 2$  then  $f_{,i} := \partial_{x^i} f$  is the partial derivative of  $f$  with respect to the  $x^i$  coordinate. Similarly,  $f_{,ij} := \partial_{x^i} \partial_{x^j} f$  and so on. When differentiating with respect to the time variable  $t$ , we set  $f_{,t} = \partial_t f$ . For horizontal derivatives, we write

$$\bar{\partial} f := f_{,1}, \quad \bar{\partial}^k f := \bar{\partial}_{x^1}^k f.$$

We use  $C$  to denote a universal constant that may vary from line to line. In numerous estimates the sign  $\lesssim$  is used; by definition,  $X \lesssim Y$  if and only if there exists a universal constant  $C$  such that  $X \leq CY$ . We use  $P$  to denote a generic real polynomial with positive coefficients that can similarly vary from line to line. We always sum over repeated indices.

#### 1.4. Fixing the Domain

In order to obtain a priori estimates, and to facilitate the construction of solutions, we transform the Stefan problem to an equivalent problem on a fixed domain. To this end, we shall view  $\Gamma(t)$  as a graph over  $\mathbb{T}^1$  given by the *height function*  $h(t, \cdot) : \Gamma \rightarrow \mathbb{R}$

$$\Gamma(t) := \Psi(t, \Gamma).$$

In other words, the moving surface  $\Gamma(t)$  is parameterized as the graph of a signed height function  $h(t, x)$ , so that  $\Gamma(t) = \{\mathbf{x} \in \mathbb{T}^1 \times \mathbb{R} \mid \mathbf{x} = (x', h(t, x'))\}$ . With this parameterization, the outward unit normal  $n(t, x')$  to  $\Gamma(t)$  at the point  $(x', h(t, x'))$  is given by

$$n(t, x') = \frac{(\bar{\partial} h, -1)}{\sqrt{1 + |\bar{\partial} h|^2}}. \quad (1.5)$$

Assuming that  $h(t, \cdot)$  is sufficiently regular and remains a graph, we can define a diffeomorphism  $\Psi(t, \cdot) : \Omega \rightarrow \Omega(t)$  as an harmonic extension of the boundary diffeomorphism  $(x', h)$  by solving the elliptic equation

$$\Delta \Psi(t, \cdot) = 0 \quad \text{in } \Omega, \quad (1.6a)$$

$$\Psi(t, x', 0) = (x', h(t, x')) \quad x' \in \Gamma, \quad (1.6b)$$

$$\Psi(t, \cdot) = \text{Id} \quad \text{on } \partial\Omega_{\text{top}}, \quad (1.6c)$$

where  $\text{Id}$  denotes the identity map. The mapping  $\Psi(t, \cdot)$  is indeed a diffeomorphism; note that the map  $\Phi := \Psi - \text{Id}$  solves the problem

$$\Delta \Phi(t, \cdot) = 0 \quad \text{in } \Omega, \quad (1.7a)$$

$$\Phi(t, x', 0) = (0, h(t, x')) \quad x' \in \Gamma, \quad (1.7b)$$

$$\Phi(t, \cdot) = \mathbf{0} \quad \text{on } \partial\Omega_{\text{top}}, \quad (1.7c)$$

so that by elliptic estimates, we may conclude that  $\|\Psi(t, \cdot) - \text{Id}\|_{H^{4.5}(\Omega)} \lesssim |h(t, \cdot)|_{H^4(\Gamma)} \lesssim \epsilon_0$ , using assumption (1.3), and the continuity of the map  $t \mapsto h(t, \cdot)$  in  $H^4(\Gamma)$ , which will be proved below. By the inverse function theorem we infer that  $\Psi(t, \cdot)$  is a diffeomorphism.

As a consequence of (1.6),

$$\|\Psi\|_{H^s(\Omega)} \leq C \|\Psi\|_{H^{s-0.5}(\Gamma)}, \quad (1.8)$$

and thus  $\Psi(t, \cdot)$  gains a half-derivative of regularity in  $\Omega$  with respect to the height function  $h(t, \cdot)$  on  $\Gamma$ .

### 1.5. Reference Unit Normal, Unit Tangent, Line Element, and the Jacobian

We let

$$N = (0, -1), \quad T = (1, 0)$$

denote the outward pointing unit normal and tangent vectors to  $\Gamma = \mathbb{T}^1 \times \{x_2 = 0\}$ , respectively. The time-dependent unit normal  $n(t, \cdot)$  and tangent  $\tau(t, \cdot)$  vectors to  $\Gamma(t)$  are given by

$$n = Jg^{-1}A^T N, \quad \tau = Jg^{-1}A^T T, \quad (1.9)$$

where

$$J(t, \mathbf{x}) := \det \nabla \Psi(t, \mathbf{x}), \quad \mathbf{x} \in \Omega, \quad (1.10)$$

denotes the Jacobian determinant of  $\nabla \Psi$ , and

$$g(t, x) := \sqrt{1 + (\bar{\partial}h(t, x))^2}, \quad x \in \mathbb{T}^1, \quad (1.11)$$

where  $g^2 dx$  is the line element associated with the metric induced on  $\Gamma$ . Together with (1.9) we obtain the relationship

$$A_{\bullet}^2 := J^{-1}(\bar{\partial}h, -1), \quad |A_{\bullet}^2| = J^{-1}g. \quad (1.12)$$

The vector  $A_{\bullet}^2(t, \cdot)$  will play an important role in the derivation of energy identities as it is parallel to  $n(t, \cdot)$ .

**1.5.1. The Change of Variables** On the reference domain  $\Omega$ , we set

$$q := p \circ \Psi, \quad A := [\nabla \Psi]^{-1}, \quad w := \Psi_t, \quad v := -\nabla p \circ \Psi. \quad (1.13)$$

Note that, by the chain-rule, the relation  $v = -\nabla p \circ \Psi$  can be written as

$$v^i + A_i^k q_{,k} = 0 \quad \text{in } \Omega. \quad (1.14)$$

We also express  $p_t \circ \Psi$  in terms of  $q, v, w$ . Again, by the chain rule,  $p_t = q_t \circ \Psi^{-1} + \nabla q \circ \Psi^{-1} \cdot \Psi_t^{-1}$ . Since  $\Psi_t^{-1} = -A \circ \Psi^{-1} \Psi_t \circ \Psi^{-1}$  and  $w = \Psi_t$ , using (1.14) we obtain that

$$p_t \circ \Psi = q_t - q_{,k} A_r^k w^r = q_t + v \cdot w.$$

The transformed Laplacian  $\Delta_{\Psi} q := \Delta p \circ \Psi$  is defined as

$$\Delta_{\Psi} q = A_i^j (A_i^k q_{,k})_{,j}, \quad (1.15)$$

and we define

$$\nabla_{\Psi} q := A_i^k q_{,k} = \nabla p \circ \Psi. \quad (1.16)$$

**Remark 1.** (Differentiation rules) When differentiating the matrix  $A = [\nabla\Psi]^{-1}$ , for a given  $i, k \in \{1, 2\}$ ,

$$\partial_t A_i^k = -A_r^k w^r_{,s} A_i^s; \quad \bar{\partial} A_i^k = -A_r^k \bar{\partial} \Psi^r_{,s} A_i^s.$$

In particular, a simple application of the above identities, together with the product rule, show that for any given  $a, b \in \mathbb{N}$ :

$$\begin{aligned} \bar{\partial}^a \partial_t^b A_i^k &= -A_r^k \bar{\partial}^a \partial_t^b \Psi^r_{,s} A_i^s + \{\bar{\partial}^a \partial_t^b, A_i^k\}; \\ \{\bar{\partial}^m \partial_t^n, A_i^k\} &:= \sum_{l+l' \geq 1} a_{l,l'} \bar{\partial}^l \partial_t^{l'} (A_r^k A_i^s) \bar{\partial}^{m-l} \partial_t^{n-l'} \Psi^r_{,s}, \end{aligned} \quad (1.17)$$

where the term  $\{\cdot, \cdot\}$  is the *commutator* error. Here the constants  $a_{l,l'}$  are some universal constants, depending only on  $m, n, l$  and  $l'$  (where  $0 \leq l \leq m, 0 \leq l' \leq n$ ).

**1.5.2. Classical Stefan Problem in the New Variables** Using the family of diffeomorphisms  $\Psi(t, \cdot)$ , the classical Stefan problem (i.e. problem (1.1) with  $\sigma = 0$ ) on the fixed reference domain  $\Omega$  is given by

$$q_t - \Delta_\Psi q = -v \cdot w \quad \text{in } \Omega \times (0, T], \quad (1.18a)$$

$$v^i + A_i^k q_{,k} = 0 \quad \text{in } \Omega \times (0, T], \quad (1.18b)$$

$$q = 0 \quad \text{on } \Gamma \times [0, T], \quad (1.18c)$$

$$\Delta \Psi = 0 \quad \text{on } \Omega \times [0, T], \quad (1.18d)$$

$$\Psi = \text{Id} + h N \quad \text{on } \Gamma \times [0, T], \quad (1.18e)$$

$$\Psi = \text{Id} \quad \text{on } \partial\Omega_{\text{top}} \times [0, T], \quad (1.18f)$$

$$\Psi_t \cdot n(t) = -v \cdot n(t) \quad \text{on } \Gamma \times (0, T], \quad (1.18g)$$

$$v \cdot N = 0 \quad \text{on } \partial\Omega_{\text{top}} \times [0, T], \quad (1.18h)$$

$$\Psi(0, \cdot) = \Psi_0 \quad q(0, \cdot) = q_0 = p_0 \circ \Psi_0, \quad (1.18i)$$

where  $\Delta_\Psi q$  is defined in (1.15) and  $N = (0, 1)$  is the outward-pointing unit normal to  $\partial\Omega_{\text{top}}$ . Problem (1.18) is a reformulation of problem (1.1). Condition (1.18g) is equivalent to the evolution equation for the height function  $h(t, \cdot)$  which is given by

$$h_t(t, x) = -g(t, x) \nabla_\Psi q(t, x) \cdot n(t, x), \quad x \in \mathbb{T}^1, \quad (1.19)$$

where the quantity  $g$  is defined in (1.11) and  $\nabla_\Psi q$  is defined in (1.16). The time-evolution of the map  $\Psi(t, \cdot)$  is, in turn, coupled to the evolution of  $q(t, \cdot)$  via (1.18a).

**1.5.3. The Higher-Order Energy Function  $\mathcal{E}(t)$**  We define the higher-order energy function as

$$\begin{aligned} \mathcal{E}(t) = \mathcal{E}(q, h)(t) &:= \sum_{a+2b \leq 4} \|\bar{\partial}^a \partial_t^b v\|_{L_t^2 L_x^2}^2 + \sum_{a+2b \leq 3} \|\bar{\partial}^a \partial_t^b v\|_{C_t^0 L_x^2}^2 \\ &+ \sum_{b=0}^2 \|\partial_t^b q\|_{C_t^0 H_x^{4-2b}}^2 + \sum_{b=0}^2 \|\partial_t^b q\|_{L_t^2 H_x^{5-2b}}^2 \end{aligned}$$

$$+ \sum_{b=0}^2 |\partial_t^b h|_{C_t^0 H_x^{4-2b}}^2 + \sum_{b=0}^1 |\partial_t^b h_t|_{L_t^2 H_x^{3-2b}}^2, \quad (1.20)$$

where the time integrals in the  $L^2$ -norms above are over the time-interval  $[0, t]$ . We will show that  $\mathcal{E}(t)$  remains bounded on  $[0, T]$ .

**1.5.4. The Taylor Sign Condition** In order to obtain a locally well-posed problem for arbitrarily large initial data, we must impose the Taylor sign condition on the initial data as follows:

$$- \partial_n p_0 > 0 \quad \text{on } \Gamma(0). \quad (1.21)$$

Expressed in terms of  $q(0, \cdot)$ , (1.21) is written as

$$q_{0,2}|_{t=0} > 0 \quad \text{on } \Gamma. \quad (1.22)$$

The condition (1.22) ensures that

$$\inf_{x' \in \Gamma} q_{,2}(t, x', 0) > 0, \quad t \in [0, T], \quad (1.23)$$

if  $T > 0$  is taken sufficiently small. As mentioned in Sect. 1.1, we shall refer to (1.22) as the *Taylor sign condition* in analogy to the terminology used in the well-posedness theory in fluid mechanics [44,46]. The Taylor sign condition will provide positivity of the natural energy functional.

**Remark 2.** Note that  $q_{,2} = gJ^{-1}v \cdot n$  on  $\Gamma$ , with  $v$  defined by (1.13). By (2.12), we conclude that  $h_t = Jq_{,2}$  at time  $t = 0$ . Since the Jacobian remains positive on a short interval of time the Taylor sign condition (1.21) shows that  $h_t(0, x) < 0$  for all  $x \in \mathbb{T}^1$ . Thus, the domain  $\Omega(t)$  expands on a short interval of time.

**1.5.5. Compatibility Conditions** To ensure that the solution is continuously differentiable with respect to  $t$ , at  $t = 0$ , we must impose compatibility conditions on the initial data. In particular, restricting (1.18a) to  $\Gamma$  and evaluating at time  $t = 0$ , for  $H^4$  initial data, we find that

$$q_0 = 0 \quad \text{on } \Gamma, \quad (1.24a)$$

$$\Delta_{\Psi_0} q_0 + J_0^{-2} g_0^2 (q_0)_{,2} = 0 \quad \text{on } \Gamma. \quad (1.24b)$$

In the derivation of (1.24b) we have crucially, used (1.9) and the identity

$$\begin{aligned} v(0, \cdot) \cdot w(0, \cdot)|_{\Gamma} &= -A_i^k q_{0,k} \partial_t \Psi_t^i(0, \cdot) = -q_{0,2} A_i^k N_{,k} \Psi_t^i(0, \cdot) \\ &= J_0^{-1} g_0^{-2} q_{0,2} \Psi_t(0, \cdot) \cdot n_0 = J_0^{-1} g_0^{-2} q_{0,2} v(0, \cdot) \cdot n_0 \\ &= J_0^{-2} g_0^2 (q_0)_{,2}. \end{aligned}$$

Here  $J_0, g_0$  are the initial values of  $J, g$  defined in (1.10) and (1.11), respectively. Conditions (1.24) are satisfied for a large class of functions. Consider, simply,

a function  $q_0$  independent of  $x_1$  in the slab  $T_\epsilon = \mathbb{T}^1 \times [0, \epsilon]$  for some  $\epsilon > 0$ , and of the form

$$q_0(x_1, x_2) = -\alpha^2 \frac{x_2^2}{2} + \alpha x_2 \quad (1.25)$$

in  $T_\epsilon$  for some  $\alpha > 0$ . Condition (1.24a) is obviously satisfied, while (1.24b) reduces to the requirement that

$$\begin{aligned} 0 &= -\Delta_{\Psi_0} q_0 + J^{-2} g^2(q_0),_2^2 = -|A_\bullet^2|^2(q_0),_{22} + J^{-2} g^2(q_0),_2^2 \\ &= J^{-2} g^2\left((q_0),_{22} + (q_0),_2^2\right), \end{aligned}$$

where we have used (1.12). It is easily checked that for such  $q_0$ ,

$$(q_0),_{22} + (q_0),_2^2 = 0 \quad \text{on } \Gamma,$$

and therefore the condition (1.24b) is satisfied. Note that the assumption  $\alpha > 0$  ensures the validity of the Taylor sign condition  $(q_0),_2 > 0$ .

Since we imposed the homogeneous Neumann condition (1.18h) on the top boundary  $\partial\Omega_{\text{top}}$ , we impose the compatibility condition

$$(q_0),_2|_{t=0} = 0 \quad \text{on } \partial\Omega_{\text{top}}. \quad (1.26)$$

By employing a partition-of-unity of  $\Omega$ , we can now easily construct a  $q_0 \in H^4(\Omega)$  such that the compatibility conditions (1.24) and (1.26) are simultaneously satisfied.

**Remark 3.** The quadratic function  $q_0$  defined in (1.25) satisfies the compatibility conditions. This is one of many possible constructions of initial data satisfying the corresponding regularity and compatibility conditions.

### 1.6. Local-in-time Well-Posedness for the Classical Stefan Problem

We define

$$\mathcal{S}(t) := \{(q, h) : \mathcal{E}(q, h)(t) < \infty\}. \quad (1.27)$$

Our first result is a well-posedness statement for the classical Stefan problem.

**Theorem 1.1.** (Well-posedness of the classical Stefan problem) *Given initial conditions  $(q_0, h_0) \in \mathcal{S}(0)$  with  $q_0$  satisfying the Taylor sign condition (1.22) and the compatibility conditions (1.24)–(1.26), the problem (1.18) is locally-in-time well-posed, i.e. there is a  $T > 0$  such that and a unique solution  $(q, h)$  on the time interval  $[0, T]$  with initial data  $(q_0, h_0)$ , such that*

$$\sup_{t \in [0, T]} \mathcal{E}(q, h) \leq 2\mathcal{E}(q_0, h_0).$$

**Remark 4.** The definition of our higher-order energy function  $\mathcal{E}$  restricted to time  $t = 0$  requires an explanation of time-derivates of  $q$  and  $h$  at  $t = 0$ . Specifically, the values  $q_t|_{t=0}$ ,  $q_{tt}|_{t=0}$ ,  $h_t|_{t=0}$  and  $h_{tt}|_{t=0}$  are defined via space-derivatives using equations (1.18a) and (1.18g).



### 1.7. The Vanishing Surface Tension Limit

Our second main result establishes the vanishing surface tension limit. Denoting by  $\mathcal{H}$  the mean curvature of the free-boundary, in the  $\Psi$ -parametrization, the boundary condition (1.18c) is replaced with

$$q = \mathcal{H} = \sigma \frac{\bar{\partial}^2 \Psi \cdot n}{|\bar{\partial}^2 \Psi|^2} = -\sigma \frac{\bar{\partial}^2 h}{(1 + |\bar{\partial} h|^2)^{3/2}}; \quad (1.28)$$

then, the problem (1.18), with the boundary condition (1.28) replacing (1.18c), is the *Stefan problem with surface tension* formulated in harmonic coordinates. The high-order energy function adapted for the presence of surface tension is given by

$$\begin{aligned} \mathcal{E}^\sigma = \mathcal{E}^\sigma(q, h) &= \mathcal{E}(q, h) + \sigma \sum_{b=0}^2 |\partial_t^b h|_{C_t^0 H_x^{5-2b}}^2 \\ &+ \sigma \sum_{b=0}^1 |\partial_t^b h_t|_{L_t^2 H_x^{4-2b}}^2 + \sigma^2 \sum_{b=0}^2 |\partial_t^b h|_{L_t^\infty H_x^{4-2b}}^2, \quad \sigma > 0. \end{aligned} \quad (1.29)$$

#### 1.7.1. Compatibility Conditions For the Stefan Problem with Surface Tension

To ensure the spatial continuity of the temperature function  $q$  and its first time derivative  $q_t$  at time  $t = 0$ , we must impose two sets of compatibility conditions. The first condition is

$$q_0 = \sigma \mathcal{H}_0 = -\sigma g_0^{-3} \bar{\partial}^2 h_0 \quad \text{on } \Gamma, \quad (1.30)$$

where  $\mathcal{H}_0$  denotes the mean curvature of the initial free surface  $\Gamma_0$ , and  $g_0 = \sqrt{1 + (\bar{\partial} h_0)^2}$ . To obtain the second compatibility condition, we note that  $q_t|_\Gamma = -\sigma \partial_t \mathcal{H}|_{t=0}$ . From the boundary condition (1.18g) we can evaluate  $h_t$  at time  $t = 0$  as

$$h_t|_{t=0} = -g_0 \nabla_{\Psi_0} q_0 \cdot n_0, \quad (1.31)$$

where the subscript 0 refers to the initial values of the quantities  $g$ ,  $\Psi$ ,  $q$ , and  $n$  defined above. Therefore,

$$\begin{aligned} \partial_t \mathcal{H}|_{t=0} &= -g_0^{-3} \bar{\partial}^2 h_t|_{t=0} + 3g_0^{-5} \bar{\partial}^2 h_0 \bar{\partial} h_t|_{t=0} \bar{\partial} h_0 \\ &= g_0^{-3} \bar{\partial}^2 (g_0 \nabla_{\Psi_0} q_0 \cdot n_0) + 3q_0 g_0^{-2} \bar{\partial} (g_0 \nabla_{\Psi_0} q_0 \cdot n_0) \bar{\partial} h_0, \end{aligned} \quad (1.32)$$

where we have used (1.30) and (1.31) in the last line. After restricting (1.18a) to  $\Gamma$  at time  $t = 0$  and using (1.32), we find that

$$\begin{aligned} &-\sigma \left( g_0^{-3} \bar{\partial}^2 (g_0 \nabla_{\Psi_0} q_0 \cdot n_0) + 3q_0 g_0^{-2} \bar{\partial} (g_0 \nabla_{\Psi_0} q_0 \cdot n_0) \bar{\partial} h_0 \right) \\ &-\Delta_{\Psi_0} q_0 = -g_0 (\nabla_{\Psi_0} q_0 \cdot n_0) (A_0)_{2,q_0,k}^k. \end{aligned}$$

In particular, the right-hand side can be separated into the  $\sigma$ -dependent and  $\sigma$ -independent contributions, so that

$$\begin{aligned}
 -g_0 (\nabla_{\Psi_0} q_0 \cdot n_0) A_2^k q_{0,k} &= J_0^{-2} g_0^2(q_0)_{,2}^2 + \sigma \bar{\partial} \mathcal{H}_0 \left( g_0 (\nabla_{\Psi_0} q_0 \cdot n_0) (A_0)_2^1 \right. \\
 &\quad \left. + g_0 \left( (A_0)_\bullet^1 \cdot n_0 \right) A_2^2(q_0)_{,2} \right).
 \end{aligned}$$

Combining the two previous identities, we find the second compatibility condition to be

$$\Delta_{\Psi_0} q_0 + J_0^{-2} g_0^2(q_0)_{,2}^2 = \sigma \mathcal{C}(q_0, h_0), \quad (1.33)$$

where

$$\begin{aligned}
 \mathcal{C}(q_0, h_0) &:= - \left[ g_0^{-3} \bar{\partial}^2 (g_0 \nabla_{\Psi_0} q_0 \cdot n_0) + 3 q_0 g_0^{-2} \bar{\partial} (g_0 \nabla_{\Psi_0} q_0 \cdot n_0) \bar{\partial} h_0 \right] \\
 &\quad - \bar{\partial} \mathcal{H}_0 \left[ g_0 (\nabla_{\Psi_0} q_0 \cdot n_0) (A_0)_2^1 + g_0 \left( (A_0)_\bullet^1 \cdot n_0 \right) (A_0)_2^2(q_0)_{,2} \right].
 \end{aligned} \quad (1.34)$$

**1.7.2. Initial Data Satisfying Compatibility Conditions** When  $\Psi = \text{Id}$  (and therefore  $h_0(x) = 0$ ,  $g_0 = J_0 = 1$ ) the compatibility conditions (1.30) and (1.33)–(1.36) simplify significantly and take the form

$$q_0 = 0 \text{ and } (q_0)_{,22} + (q_0)_{,2}^2 = \sigma (q_0)_{,211} \text{ on } \Gamma \quad (1.35)$$

$$(q_0)_{,222}, (q_0)_{,211} \in C^0(\Gamma). \quad (1.36)$$

It is easy to check that the function  $q_0$  constructed in Section 1.5.5 satisfies (1.35)–(1.36). For general  $h$  satisfying  $|h|_4 \ll 1$  we can construct the initial temperature  $q_0$  satisfying (1.30) and (1.33)–(1.36) by perturbative methods, using, for instance, the implicit function theorem.

**1.7.3. Well-Prepared Initial Data** To obtain the vanishing surface tension limit, we need to define a suitable class of initial data  $(q_0^\sigma, h_0^\sigma)$ ,  $\sigma \geq 0$ .

**Definition 1.** (Well-prepared data) A family of initial data  $(q_0^\sigma, h_0^\sigma)_{\sigma \geq 0}$  such that  $\mathcal{E}(q_0^\sigma, h_0^\sigma) < \infty$  is well-prepared if it satisfies (1) compatibility conditions (1.30), (1.33)–(1.36) associated to the Stefan problem with surface tension, (2) the Taylor sign condition (1.22), and (3)  $\mathcal{E}(q_0^\sigma - q_0, h_0^\sigma - h_0) \rightarrow 0$  as  $\sigma \rightarrow 0$ .

We now demonstrate that the class of well-prepared initial data is non-empty. Let us assume for simplicity that  $\Psi_0 = \text{Id}$  and therefore the initial hypersurface  $\Gamma_0$  is flat. For  $\sigma \geq 0$  we have  $h^\sigma(x') = 0$ ,  $x' \in \mathbb{T}^1$ . Let  $b : \mathbb{T}^1 \rightarrow \mathbb{R}$  be a given smooth function and  $\alpha > 0$  a given positive real number. Consider a function  $q_0^\sigma$  independent of  $x_1$  in the slab  $T_\epsilon = \mathbb{T}^1 \times [0, \epsilon]$  for some  $0 < \epsilon < 1$  and of the form

$$q_0^\sigma(x_1, x_2) = -\alpha^2 \frac{x_2^2}{2} + \alpha x_2 + \sigma b(x_1) x_2^3, \quad (x_1, x_2) \in T_\epsilon.$$

It is straightforward to check that conditions (1.35)–(1.36) are both satisfied with this choice of  $q_0^\sigma$ . Moreover, The Taylor sign condition holds since  $q_{0,2}^\sigma = \alpha > 0$  for any  $\sigma \geq 0$  and the convergence requirement (3) in Definition 1 is clearly satisfied. Outside the slab  $T_\epsilon$  we can extend the function  $q_0^\sigma$  smoothly so that the Neumann boundary condition  $\partial_N q_0^\sigma$  is satisfied on  $\partial\Omega_{\text{top}}$ .

**1.7.4. The Vanishing Surface Tension Limit** For a given  $T > 0$  let

$$C_t^1 C_x^0 \cap C_t^0 C_x^2 := \left\{ (q, h) : q \in C^1([0, T]; C^0(\Omega)) \cap C^0([0, T]; C^2(\Omega)), \right. \\ \left. h \in C^1([0, T]; C^0(\Gamma)) \cap C^0([0, T]; C^2(\Gamma)) \right\} \quad (1.37)$$

with the associated norm:

$$\|(q, h)\|_{C_t^1 C_x^0 \cap C_t^0 C_x^2} = \max_{t \in [0, T], x \in \Omega} (|q(t, x)| + |\partial_t q(t, x)| \\ + |\nabla q(t, x)| + |\nabla^2 q(t, x)|) \\ + \max_{t \in [0, T], x' \in \Gamma} (|h(t, x')| + |\partial_t h(t, x')| \\ + |\bar{\partial} h(t, x')| + |\bar{\partial}^2 h(t, x')|). \quad (1.38)$$

**Theorem 1.2.** (The limit of zero surface tension) *Let  $(q_0^\sigma, h_0^\sigma)_{\sigma \geq 0}$  be a sequence of well-prepared initial conditions in the sense of Definition 1 such that*

$$\mathcal{E}(q_0^\sigma - q_0, h_0^\sigma - h_0) \rightarrow 0 \text{ as } \sigma \rightarrow 0.$$

*Let  $(q^\sigma(t, \cdot), h^\sigma(t, \cdot))_{\sigma \geq 0}$  denote the corresponding sequence of solutions to the Stefan problem with surface tension, such that  $(q^\sigma(0, \cdot), h^\sigma(0, \cdot)) = (q_0^\sigma, h_0^\sigma)$ . Then, there exists a  $\sigma$ -independent time  $T > 0$  and a constant  $C$  depending only on  $(q_0, h_0)$  such that*

$$\mathcal{E}^\sigma(q^\sigma, h^\sigma)(T) \leq C \quad \sigma \geq 0$$

for all  $\sigma \geq 0$ .

*Furthermore, the sequence  $(q^\sigma, h^\sigma)$  converges in the  $C_t^1 C_x^0 \cap C_t^0 C_x^2$ -norm to the unique solution  $(q, h)$  of the classical Stefan problem (1.18) with  $\sigma = 0$  and the initial data  $(q(0), h(0)) = (q_0, h_0)$ .*

## 1.8. Prior Results and a Motivation For the Current Treatment

There is a large literature on the classical one-phase Stefan problem. For a comprehensive overview, we refer the reader to MEIRMANOV [40] and VISINTIN [49]. The first *weak* solutions were defined by KAMENOMOSTSKAYA [35], LADYZHENSKAYA, SOLONNIKOV and URALCEVA [38]. These weak solutions were analyzed by FRIEDMAN, KINDERLEHRER [24–26], CAFFARELLI, EVANS [6, 7], wherein the regularity of weak solutions was established. Since the problem satisfies a maximum principle, it is ideally suited to the so-called *viscosity solutions* approach. The existence and regularity of viscosity solutions was established by ATHANASOPOULOS, CAFFARELLI, AND SALSA in [4, 5]. The existence of viscosity solutions in the one-phase case was proven by KIM [36] and in the two-phase case by KIM AND POŽAR [37]. A local-in time regularity theorem was proven in [13] which in particular shows that initially Lipschitz free-boundaries become  $C^1$  over a possibly smaller spatial region. For an exhaustive overview and introduction to the regularity theory of such solutions we refer the reader to CAFFARELLI AND SALSA [8], see also more recent results [12, 13].

The local existence of classical solutions for the classical Stefan problem was shown by MEIRMANOV (see [40] and references therein) and HANZAWA [34]. In the first approach, the author regularizes the problem by adding artificial viscosity to (1.1b) and fixes the moving domain by switching to so-called von Mises variables. The obtained solutions however, lose derivatives with respect to the assumed regularity on the initial data. Similarly, in [34] the author uses Nash–Moser iteration to obtain a local-in-time solution, however again with a significant derivative loss with respect to the initial data. A local existence result for the one-phase  $n$ -dimensional Stefan problem is proved in [28], where the required regularity class for the temperature function is  $W_p^{2,1}$  with  $p > n + 2$ . For the two-phase Stefan problem a local existence result is presented in [41] in the framework of  $L^p$  maximal regularity, where the corresponding functional spaces of Sobolev-type require  $p > n + 3$ , where  $n$  is the dimension of the ambient space.

In related work, local and global existence for the one-phase and two-phase Muskat problems has been established in [10, 14–16]. For the local and global well-posedness of the one-phase Hele-Shaw problem and optimal decay rates of the solutions, see [9] and the references therein.

As to the Stefan problem with surface tension (also known as the Stefan problem with Gibbs–Thomson correction), a global weak existence theory (without uniqueness) is given in [1, 39, 45]. In [27] the authors consider the Stefan problem with small surface tension i.e.  $\sigma \ll 1$  whereby (1.1c) is replaced by  $v = \sigma \kappa$ . The local existence of classical solutions is studied in [43]. In [23], the authors prove a local existence and uniqueness result for classical solutions under a smallness assumption on the initial datum close to flat hypersurfaces. The global existence close to flat hyper-surfaces is proved in [30] and close to stationary spheres for the two-phase problem in [29] and later in [42].

With the Gibbs–Thomson correction, problem (1.1) can account for phenomena such as phase nucleation and undercooling (superheating); it is also used in modeling crystal growth [49]. This is a small-scale model as opposed to the macro-scale classical Stefan problem. In this sense, there is a fundamental importance in rigorously understanding the link between the two models. As explained in [48, 49], one can associate a free energy to the Stefan problem with surface tension defined by

$$F_\sigma(\tilde{p}, \tilde{\Gamma}) = \int_\Omega \tilde{p} \, dx + \sigma |\tilde{\Gamma}|,$$

where  $\tilde{p}$ ,  $\tilde{\Gamma}$  are time-independent. Then, in the the sense of  $\Gamma$ -convergence of De Giorgi [22], the free energy  $F_\sigma(\tilde{p}, \tilde{\Gamma})$  converges to the free energy for the classical Stefan problem, see [49]. This is, however, a completely time-independent consideration and does not address the vanishing  $\sigma$ -limit of time-dependent solutions to the full non-linear problem (1.1). In the context of the water wave problem, the vanishing surface tension limit in two and three dimensions has been studied in [2, 3]; for the full Euler equations, see [17].

Turning our attention to the Stefan problem, we can observe that there are two parallel developments in the existence theory for weak solutions briefly mentioned above. The first one applies to the classical Stefan problem and is motivated by the

validity of maximum principle; suitable notions of weak and viscosity solutions have been established [4–6, 26, 38]. The second development refers to the problem with surface tension, wherein the weak solution existence results are in BV-type spaces, and rely upon the gradient-flow structure of the problem. From the point of view of the vanishing surface tension, it is natural to ask whether the two concepts are compatible in any rigorous mathematical manner. The answer is inconclusive due to a lack compactness. While the control of solutions constructed in [1, 39] is strong enough to pass to some limit as  $\sigma \rightarrow 0$ , it is too weak to guarantee a sharp interface in the limit. In other words, it is not clear how to preclude the formation of so-called mushy regions [49].

We develop a *new* energy method for the Stefan problem with and without surface tension and prove the vanishing surface tension limit. The *well-posedness* is established in  $H^k$  Sobolev spaces using a combination of energy estimates for tangential derivatives and elliptic-type estimates for added parabolic-type regularity. Our framework is motivated by the analysis of the free-surface incompressible Euler equations of COUTAND AND SHKOLLER [19, 20].

Precise statements of our results are given in Theorems 1.1 and 1.2. The estimates that we use are nonlinear in nature and they fundamentally exploit the intricate energy structure of the problem. In particular, no derivative loss occurs with respect to the regularity of the initial data. This framework is particularly convenient, as it allows us to rigorously establish the vanishing surface tension limit locally-in-time, as formulated in Theorem 1.2. In this way, we link two fundamental models of phase transitions that are valid on different spatial scales, thus answering the open question explained above. In forthcoming work, we shall extend our results to the two-phase Stefan problem, providing the analog of Theorem 1.1 [33], while the question of global-in-time stability of steady states using this functional-analytic framework has been addressed in [31, 32].

### 1.9. Methodology and Outline of the Paper

There are three main ingredients in our approach to the Stefan problem. First, we replace the study of the heat equation for temperature with the equation for a velocity field  $u(t, x)$  which satisfies the equation  $u + \nabla p = 0$ . Second, we introduce the so-called Arbitrary Lagrange-Eulerian (ALE) variables, in which we introduce a family of diffeomorphisms  $\Psi(t, \cdot) : \Omega \rightarrow \Omega(t)$  which fix the moving domain. With respect to this change-of-variables, we define, respectively, the new velocity and temperature fields  $v = u \circ \Psi$  and  $q = p \circ \Psi$ ; in these variables, the velocity equation becomes  $v + \nabla p \circ \Psi = 0$ . This equation contains the geometry of the evolving free-boundary, and by the use of energy estimates for tangential derivatives, we are able to naturally estimate the second-fundamental form as

$$\int_{\Gamma} q_{,2} |\bar{\partial}^k \Psi \cdot A_{\bullet}^2|^2 dx' \approx \int_{\Gamma} q_{,2} |\bar{\partial}^k h|^2 dx', \quad (1.39)$$

for  $k$  some positive integer. In the original Eulerian framework (1.1), the energy dissipation law is given by

$$\frac{1}{2} \frac{d}{dt} \int_{\Omega(t)} p(t, x)^2 dx + \int_{\Omega(t)} |\nabla p(t, x)|^2 dx = 0.$$

This basic energy law is too weak to control the regularity of the evolving free-boundary. Observe that our higher-order control of the free-boundary given by (1.39) naturally produces the stability condition; in particular, the Taylor sign condition (1.22) arises as a coefficient to the second-fundamental form, and its sign determines either the control or growth of the curvature and its derivatives via a Gronwall-type inequality.

A further subtlety consists in the discovery of another coercive energy term which is defined on the whole domain  $\Omega$  (phase), displayed in the fourth line of (2.24). It contains terms of the general form

$$\|\bar{\partial}^a \partial_t^b q + \bar{\partial}^a \partial_t^b \Psi \cdot v\|_{L_t^\infty L_x^2}^2 \quad \text{and} \quad \|\bar{\partial}^a \partial_t^b q_t + \bar{\partial}^a \partial_t^b \Psi_t \cdot v\|_{L_t^2 L_x^2}^2$$

for  $a, b$  as in (2.24). They are *intrinsically* linked to the problem and contain information about the regularity of the divergence of the velocity  $v$ . Taking  $a = 0$  and  $b = 1$ , the first term above becomes the norm of the ALE-divergence of  $v$ , as it is easily seen from (1.18a):

$$\|q_t + v \cdot w\|_{L_t^\infty L_x^2} = \|\operatorname{div}_\Psi v\|_{L_t^\infty L_x^2}.$$

The gauge condition (1.6) allows us to get optimal Sobolev regularity for  $\Psi$  and hence for the temperature function  $q$ . This allows us to prove that the energy  $\mathcal{E}$  defined in (2.16) is in fact bounded by the coercive quadratic form (the “natural energy”)  $\mathcal{F}$  (2.24) dictated by the Stefan problem.

Condition (1.21) is the exact equivalent of the Taylor sign condition, necessary for well-posedness of free-surface incompressible Euler equations without surface tension [19] or the water wave problem [50]. If the initial temperature  $q_0$  is non-negative, it is implied by Hopf’s lemma, at least over a short period of time. In a short time regime, we prove a uniform lower bound on  $\lambda$  [cf. (1.23)], thus enabling us to close the estimates.

In many free-boundary problems, constructing the solution is, in general, a challenging problem, despite the (possible) availability of good a-priori estimates. Our main technical idea, to make the construction as straightforward as possible, is to regularize the problem via *horizontal convolution by layers* as introduced in [19] in the study of well-posedness of the incompressible Euler equation on a moving domain. In addition to that, we also regularize the Stefan condition  $p = 0$  on  $\Gamma(t)$  by modifying it into a Robin-type condition. If  $\kappa > 0$  is a suitable regularization parameter, to each  $\kappa$  we shall associate an energy functional  $\mathcal{E}_\kappa$  which will be shown to satisfy the following energy inequality:

$$\mathcal{E}_\kappa(t) \leq C\mathcal{E}_\kappa(0) + C(t + \sqrt{t})P(\sqrt{\mathcal{E}_\kappa}),$$

where  $P$  is something with the leading order cubic contribution. Such a polynomial inequality, through a continuity argument, leads to uniform-in- $\kappa$  time of existence  $[0, T]$  and the bound

$$\mathcal{E}_\kappa(t) \leq 2C\mathcal{E}_\kappa(0).$$

Passing to the limit as  $\kappa \rightarrow 0$ , we recover the solution of the Stefan problem (1.18). Our regularization is intrinsic to the problem and it does not rely on formulating a sequence of iterated linear problems.

The second part of this work focuses on the problem of the vanishing surface tension limit. Once the well-posedness framework of Theorem 1.1 is set-up, the idea is rather straightforward. Namely, at the level of energy, the presence of surface tension simply augments the high-order energy functional by a  $\sigma$ -dependent contribution coming from the boundary  $\Gamma$ , so as to obtain (1.29). The goal is to prove a uniform-in- $\sigma$  upper bound on  $\mathcal{E}^\sigma$  on a  $\sigma$ -independent time interval  $[0, T]$ . This is made possible by one fundamental property of  $\mathcal{E}^\sigma$ : it distinguishes between two boundary energy contributions of general forms

$$|\sqrt{q, 2} \bar{\partial}^a \partial_t^b h|_0^2 \quad \text{and} \quad \sigma |\bar{\partial}^{a+1} \partial_t^b h|_0^2$$

for suitable  $a, b \in \mathbb{N}_0$ . Since the error terms are at least of cubic order, we can afford to estimate *all* lower order terms in terms of the  $\sigma$ -independent energy term, while the two terms with highest number of derivatives get bounded via the  $\sigma$ -dependent energy contribution. With uniform estimates in hand, we can pass to the limit as  $\sigma \rightarrow 0$ .

The plan of the paper is as follows. In Sect. 2.1 we introduce the  $\kappa$ -regularized problem and the associated high-order energy  $\mathcal{E}_\kappa$ . We then state the energy identities (Lemma 2.2), prove that  $\mathcal{E}_\kappa$  is controlled by the natural energy  $\mathcal{F}_\kappa$  (Proposition 2.5) and finally prove Lemma 2.2. In Sect. 2.7, we provide the energy estimates for the error terms. Passage to the vanishing surface tension limit is explained in Sect. 3. In Sect. 4 we explain how to extend our results to the three dimensional setting.

## 2. Local Well-Posedness For the Classical Stefan Problem

### 2.1. A Nonlinear Regularization of the Stefan Problem: the $\kappa$ -Problem

We regularize the problem by using the *horizontal convolution by layers*, introduced in [19] in the study of well-posedness of the incompressible Euler equation on a moving domain.

**Definition 2.** (Horizontal convolution-by-layers) Let  $\rho_\kappa$  be a  $C^\infty(\mathbb{R})$ -bump function supported in a ball of radius  $\kappa$  defined through:  $\rho_\kappa(x) := \frac{1}{\kappa} \rho(\frac{x}{\kappa})$ , where

$$\rho(x) = \begin{cases} c_* e^{-1/(1-|x|^2)}, & |x| < 1, \\ 0 & |x| \geq 1 \end{cases} \quad (2.1)$$

and constant  $c_*$  is such that  $\int_{\mathbb{R}} \rho(x') dx' = 1$ . For any given  $g : \Omega \rightarrow \mathbb{R}$  we define the horizontal convolution by layers of  $g$  via

$$\Lambda_\kappa g(x^1, x^2) := \int_{\Gamma} g(x^1, x^2) \rho_\kappa(x^1 - x') dx'.$$

We also define the standard 2-D sequence of mollifiers:  $\eta_\kappa(x) = \kappa^{-2}\eta(x/\kappa)$  where  $\eta(x) = c_*e^{-1/(1-|x|^2)}$  for  $|x| < 1$  and  $\eta(x) = 0$  for  $|x| \geq 1$ , and  $c_*$  is chosen so that  $\int_{\mathbb{R}^2} \eta(x)dx = 0$ . To formulate the regularized problem, we introduce the following quantities:

$$\Psi_\kappa(t, x') = (x', h_\kappa(t, x')) \quad \text{with} \quad h_\kappa(t, x') := \Lambda_\kappa \Lambda_\kappa h(t, x'), \quad x' \in \Gamma,$$

and we define  $\Psi_\kappa$  on  $\Omega$  as a harmonic extension of its boundary value on  $\Gamma$  as in (1.6). Analogously to (1.8), the following trace estimate is true:

$$\|\partial_t^a \Psi_\kappa\|_{H^s(\Omega)} \lesssim \|\partial_t^a \Psi_\kappa\|_{H^{s-0.5}(\Gamma)}, \quad s > 0.5, \quad a \in \mathbb{N}. \quad (2.2)$$

We also denote  $J_\kappa := \det \nabla \Psi_\kappa$ . Furthermore,

$$\mathbb{A} := [\nabla \Psi_\kappa]^{-1}; \quad \mathbb{W} := \partial_t \Psi_\kappa.$$

In analogy to (1.12), we introduce

$$\mathbb{A}_\bullet^2 := \frac{(\bar{\partial} h_\kappa, -1)}{J_\kappa}; \quad \mathbb{A} := J_\kappa^{-1} \mathbb{A}. \quad (2.3)$$

For  $\kappa > 0$ , we now define a nonlinear regularization of the Stefan problem, which we call the  $\kappa$ -problem (1.18), in which the coefficients are smoothed by use of the horizontal convolution operator  $\Lambda_\kappa$ . On a time interval  $[0, T_\kappa]$ , the  $\kappa$ -problem is given as

$$q_t - \Delta_{\Psi_\kappa} q = -v \cdot \mathbb{W} + \alpha \quad \text{in } [0, T_\kappa] \times \Omega, \quad (2.4a)$$

$$v^j + \mathbb{A}_i^k q_{,k} = 0 \quad \text{in } [0, T_\kappa] \times \Omega, \quad (2.4b)$$

$$q = -\kappa^2 v \cdot \mathbb{A}_\bullet^2 + \kappa^2 \beta(t, x') \quad \text{on } [0, T_\kappa] \times \Gamma, \quad (2.4c)$$

$$\Psi_t \cdot n_\kappa = -v \cdot n_\kappa \quad \text{on } [0, T_\kappa] \times \Gamma, \quad (2.4d)$$

$$v \cdot N = 0 \quad \text{on } [0, T_\kappa] \times \partial\Omega_{\text{top}}, \quad (2.4e)$$

$$\Psi(0, \cdot) = \Psi_0 \quad q(0, \cdot) = \mathcal{Q}_0^k, \quad (2.4f)$$

where

$$\Delta_{\Psi_\kappa}(t, \cdot) = 0 \quad \text{in } \Omega, \quad (2.5a)$$

$$\Psi_\kappa(t, x', 0) = (x', h_\kappa(t, x')) \quad x' \in \Gamma, \quad (2.5b)$$

$$\Psi_\kappa(t, \cdot) = \text{Id} \quad \text{on } \partial\Omega_{\text{top}}, \quad (2.5c)$$

$\Delta_{\Psi_\kappa} q := \mathbb{A}_i^j (\mathbb{A}_i^k q_{,k})_{,j}$ , and the time-independent forcing function  $\alpha(x)$  is given by

$$\alpha = J_0^{-2} g_0^2 [q_{0,2}]^2 - J_\kappa(0, \cdot)^{-2} g_\kappa(0, \cdot)^2 [q_{0,2}]^2. \quad (2.6)$$

Here

$$g_\kappa(t, x) := \sqrt{1 + (\bar{\partial} h_\kappa(t, x))^2},$$



and  $\beta(t, x')$  is defined as

$$\beta(t, x') := \sum_{k=0}^2 \frac{t^k}{k!} \partial_t^k (v \cdot \xi_{\bullet}^2)|_{t=0}. \quad (2.7)$$

Note that we use the subscript and superscript  $\kappa$  on dependent variables in which there is explicit use of the horizontal convolution operator  $\Lambda_\kappa$ ; of course, all of the  $q$ ,  $h$ , and  $\Psi$  implicitly depend on  $\kappa$  as well, but for notational convenience, we do not indicate this implicit dependence on  $\kappa$ .

The presence of the horizontal mollification operator  $\Lambda_\kappa$  in the approximate  $\kappa$ -problem changes the compatibility conditions on the the initial data. The addition of the forcing functions  $\alpha(x)$  and  $\beta(t, x')$  ensure that the compatibility conditions (1.24) are modified to be

$$Q_0^\kappa = 0 \quad \text{on } \Gamma, \quad (2.8a)$$

$$\Delta_{\Psi_0^\kappa} Q_0^\kappa = -J_0^{-2} g_0^2 [q_{0,2}]^2 \quad \text{on } \Gamma, \quad (2.8b)$$

where  $\Psi_0^\kappa = \Psi_\kappa(0, \cdot)$ . The approximated initial temperature function  $Q_0^\kappa$  is then defined as the solution of the fourth-order elliptic equation

$$\Delta_{\Psi_0^\kappa} \Delta_{\Psi_0^\kappa} Q_0^\kappa = \eta_\kappa * E(\Delta_{\Psi_0} \Delta_{\Psi_0} q_0) \quad \text{in } \Omega, \quad (2.9a)$$

$$Q_0^\kappa = 0 \quad \text{on } \Gamma, \quad (2.9b)$$

$$\Delta_{\Psi_0^\kappa} Q_0^\kappa = -J_0^{-2} g_0^2 [q_{0,2}]^2 \quad \text{on } \Gamma, \quad (2.9c)$$

where  $E$  continuously maps  $H^k(\Omega)$  to  $H^k(\mathbb{R}^2)$  for all  $k \geq 0$ . The fourth-order elliptic equation (2.9) can be written as a system of second-order equations given by

$$\Delta_{\Psi_0^\kappa} Q_0^\kappa = R_0^\kappa \quad \text{in } \Omega, \quad (2.10a)$$

$$\Delta_{\Psi_0^\kappa} R_0^\kappa = \eta_\kappa * E(\Delta_{\Psi_0} \Delta_{\Psi_0} q_0) \quad \text{in } \Omega, \quad (2.10b)$$

$$Q_0^\kappa = 0 \quad \text{on } \Gamma, \quad (2.10c)$$

$$R_0^\kappa = -J_0^{-2} g_0^2 [q_{0,2}]^2 \quad \text{on } \Gamma. \quad (2.10d)$$

According to the basic elliptic regularity theorem with Sobolev class coefficients, Theorem 3.6 in [11], we obtain estimates for  $R_0^\kappa$  and then  $Q_0^\kappa$  which show that

$$\|Q_0^\kappa\|_4^2 \leq C \mathcal{E}(q_0, h_0),$$

the constant  $C$  being independent of  $\kappa$ . Thus we see that  $Q_0^\kappa \rightarrow q_0$  in  $H^s(\Omega)$ ,  $s \in [0, 4)$ , and so we have an approximated initial temperature function  $Q_0^\kappa \in H^4(\Omega)$ , which satisfies the compatibility conditions (2.8). Again, from the elliptic system (2.10) and the Sobolev embedding theorem,  $Q_0^\kappa \rightarrow q_0$  in  $C^1(\overline{\Omega})$ , and hence the Taylor sign condition (1.22) remains valid for  $Q_0^\kappa$ , so that

$$(Q_0^\kappa)_{,2} \Big|_{t=0} > 0 \quad \text{for sufficiently small } \kappa > 0. \quad (2.11)$$

In equation (2.4d),  $n_\kappa$  denotes the outer unit normal with respect to the regularized surface  $\Gamma_\kappa$ , i.e. in the coordinate representation

$$n_\kappa = \frac{(\bar{\partial}h_\kappa, -1)}{\sqrt{1 + |\bar{\partial}h_\kappa|^2}} = g_\kappa^{-1}(\bar{\partial}h_\kappa, -1).$$

Note that the corresponding unit tangent to  $\Gamma_\kappa$  is given via

$$\tau_\kappa = \frac{\bar{\partial}\Psi_\kappa}{|\bar{\partial}\Psi_\kappa|} = \frac{(1, \bar{\partial}h_\kappa)}{\sqrt{1 + |\bar{\partial}h_\kappa|^2}} = g_\kappa^{-1}(1, \bar{\partial}h_\kappa).$$

In analogy to (2.12), equation (2.4d) can be reformulated as an evolution equation for  $h$ , given by

$$h_t(t, x) = g_\kappa(t, x)v \cdot n_\kappa(t, x), \quad x \in \mathbb{T}^1. \quad (2.12)$$

**Remark 5.** (The regularization (2.4c)) The approximate  $\kappa$ -problem uses horizontal convolution by layers together with carefully chosen artificial viscosity terms. This approximation scheme provides a simple existence theory for the  $\kappa$ -problem while maintaining the nonlinear energy structure.

**Remark 6.** We introduce the regularization (2.4c) to circumvent a technical difficulty of closing the energy estimates at the level of highest-in-time differentiated problem. The problem arises from the commutation of the horizontal convolution operator appearing in the terms of the following schematic form:

$$\int_{\Gamma} \Lambda_\kappa \Lambda_\kappa \Psi_{tt} \cdot \Psi_{ttt} \mathcal{T} dx',$$

where  $\mathcal{T}$  is a lower order term. Of course, when performing a-priori estimates (i.e. assuming that the solutions to the *original* problem are smooth enough to justify all the integrations by parts), such an issue does not arise.

### 2.1.1. Solutions to the $\kappa$ -Problem

**Theorem 2.1.** *Let  $\kappa > 0$  be fixed. Let  $(Q_0^\kappa, h_0) \in H^4(\Omega) \times H^4(\Gamma)$  be given initial data satisfying the compatibility conditions (2.8). Then there is a time  $T_\kappa$  depending on  $\kappa$ , such that there exists a unique solution  $(q, h) = (q(\kappa), h(\kappa))$  to (2.4) on the time interval  $[0, T_\kappa]$ . The solution satisfies*

$$\begin{aligned} & \sum_{a=0}^2 \left( \|\partial_t^a q\|_{C_t^0 H_x^{4-2a}} + \|\partial_t^a q\|_{L_t^2 H_x^{5-2a}} + \|q_{ttt}\|_{L_t^2(H_x^1)'} \right. \\ & \left. + \|\partial_t^{a+1} h\|_{L_t^2 H_x^{4-2a}} \right) + \sum_{a=0}^1 \|\partial_t^{a+1} h\|_{C_t^0 H_x^{3-2a}} < \infty, \end{aligned} \quad (2.13)$$

where  $H^1(\Omega)'$  denotes the dual space of  $H^1(\Omega)$ .

**Proof.** We briefly sketch the proof. For  $T_\kappa$  fixed (and taken sufficiently small) and for  $K > 0$ , we define the closed set

$$\begin{aligned} Z_K := & \left\{ h : [0, T] \times \Gamma \rightarrow \mathbb{R}, |\partial_t^a h \in C([0, T_\kappa], \right. \\ & H^{4-2a}(\Gamma)) \cap L^2([0, T_\kappa], H^{5-2a}(\Gamma)), a = 0, 1, 2, \\ & \left. \sum_{a=0}^2 \left( |\partial_t^a h|_{C_t^0 H^{4-2a}}^2 + |\partial_t^a h|_{L_t^2 H^{5-2a}}^2 \right) \leq K, \right. \\ & \left. h_0 \text{ and } Q_0^\kappa \text{ satisfy compatibility conditions (2.8)} \right\}. \end{aligned} \quad (2.14)$$

Given  $h \in Z_K$ , we define  $h_\kappa = \Lambda_\kappa^2 h$ , and then we define its harmonic extension  $\Psi_\kappa$  by solving (2.5). We then define the corresponding  $\mathcal{A}$ ,  $\mathcal{A}$ , and  $J_\kappa$ , and consider the weak formulation of the parabolic problem (2.4a)–(2.4c): for all test functions  $\phi \in H^1(\Omega)$  and a.e.  $t \in [0, T]$ ,

$$\begin{aligned} \langle q_t J_\kappa, \phi \rangle + \int_\Omega q_{,k} \mathcal{A}_i^k \mathcal{A}_i^j \phi_{,j} J_\kappa dx + \frac{1}{\kappa^2} \int_\Gamma q \phi dx_1 \\ = \int_\Omega q_{,k} \mathcal{A}_i^k \Phi_i^j \phi dx + \int_\Omega \alpha \phi J_\kappa dx + \int_\Gamma \beta \phi dx_1, \end{aligned} \quad (2.15)$$

together with the initial condition

$$q(0, x) = Q_0^\kappa(x).$$

Since  $\mathcal{A}$ ,  $\mathcal{A}$ , and  $J_\kappa$  are in  $C^\infty$ , and since  $\mathcal{A}_i^k \mathcal{A}_i^j \geq \lambda$  for  $\lambda > 0$ , and the compatibility conditions are satisfied, standard parabolic theory provides the existence of a unique solution on a short time-interval  $[0, T_\kappa]$  with the desired regularity properties. In particular, it is a standard argument to establish the existence of a unique solution in  $q \in L^2(0, T_\kappa; H^5(\Omega))$  which satisfies the estimate (2.13).

Using a Galerkin scheme on (2.15), we obtain unique solutions in  $L^2(0, T_\kappa; H^1(\Omega))$  for  $q, q_t$ , and  $q_{tt}$  and also find that  $q_{ttt} \in L^2(0, T_\kappa; H^1(\Omega)')$ , where  $H^1(\Omega)'$  denotes the dual space of  $H^1(\Omega)$ . Standard parabolic regularity theory, as in [47], shows that  $q \in L^2(0, T_\kappa; H^5(\Omega))$  and that  $q_t \in L^2(0, T_\kappa; H^3(\Omega))$ .

With the solution  $q$ , we define the associated velocity field  $v$  using (2.4b). We then update the height function  $h$  as

$$\Phi(h)(t) := h_0 + \int_0^t g_\kappa(\tau) v \cdot n_\kappa(\tau) d\tau, \quad t \in [0, T].$$

Choosing  $T_\kappa$  sufficiently small, it can be shown that  $\Phi$  maps  $Z_K$  into itself, and that  $\Phi$  is a contraction map. The fixed-point of  $\Phi$  is then a solution to the  $\kappa$ -problem (2.4).  $\square$

**Remark 7.** A priori, the time of existence  $T_\kappa$  may converge to 0 as  $\kappa \rightarrow 0$ . By obtaining  $\kappa$ -independent bounds on solutions to (2.4), we will prove that, in fact, the time of existence is independent of  $\kappa$  and given by  $T > 0$ .

## 2.2. The Higher-Order Energy Function Compatible with the $\kappa \rightarrow 0$ Asymptotics

The asymptotically consistent higher-order energy function associated to our sequence of regularized  $\kappa$ -problems is given by

$$\begin{aligned}
 \mathcal{E}_\kappa = \mathcal{E}_\kappa(q, h) &:= \sum_{a+2b \leq 4} \|\bar{\partial}^a \partial_t^b v\|_{L_t^2 L_x^2}^2 + \sum_{a+2b \leq 3} \|\bar{\partial}^a \partial_t^b v\|_{C_t^0 L_x^2}^2 \\
 &+ \kappa^2 \sum_{a+2b \leq 4} |\bar{\partial}^a \partial_t^b h_t|_{L_t^2 L_x^2}^2 + \kappa^2 \sum_{a+2b \leq 3} |\bar{\partial}^a \partial_t^b h_t|_{C_t^0 L_x^2}^2 \\
 &+ \sum_{b=0}^2 \|\partial_t^b q\|_{C_t^0 H_x^{4-2b}}^2 + \sum_{b=0}^2 \|\partial_t^b q\|_{L_t^2 H_x^{5-2b}}^2 \\
 &+ \sum_{b=0}^2 |\Lambda_\kappa h|_{C_t^0 H_x^{4-2b}}^2 + \sum_{b=0}^1 |\partial_t \Lambda_\kappa h|_{L_t^2 H_x^{3-2b}}^2. \tag{2.16}
 \end{aligned}$$

As a consequence of Theorem 2.1 the map  $t \mapsto \mathcal{E}_\kappa(t)$  is continuous on  $[0, T_\kappa]$ .

## 2.3. Bounds on Lower-Order Norms

Let

$$\begin{aligned}
 \mathcal{A}_\kappa(t) &= \sum_{a+2b \leq 2} \|\bar{\partial}^a \partial_t^b v\|_{L_t^2 L_x^2}^2 + \sum_{a+2b \leq 1} \|\bar{\partial}^a \partial_t^b v\|_{L_t^\infty L_x^2}^2 \\
 &+ \kappa^2 \sum_{a+2b \leq 2} |\bar{\partial}^a \partial_t^b h_t|_{L_t^2 L_x^2}^2 + \kappa^2 \sum_{a+2b \leq 1} |\bar{\partial}^a \partial_t^b h_t|_{L_t^\infty L_x^2}^2 \\
 &+ \sum_{b=0}^1 \|\partial_t^b q\|_{L_t^\infty H_x^{2-2b}}^2 + \sum_{b=0}^1 \|\partial_t^b q\|_{L_t^2 H_x^{3-2b}}^2 \\
 &+ \sum_{b=0}^1 |\partial_t^b \Lambda_\kappa h|_{L_t^\infty H_x^{2-2b}}^2 + |\Lambda_\kappa h|_{L_t^2 H_x^{3-2b}}^2.
 \end{aligned}$$

We then assume that

$$\mathcal{A}_\kappa(t) \leq \mathcal{E}_\kappa(0) + 1, \quad t \in [0, T_\kappa]. \tag{2.17}$$

By the fundamental theorem of calculus it is easy to see that

$$\mathcal{A}_\kappa(t) \leq \mathcal{A}_\kappa(0) + t \sup_{0 \leq s \leq t} \mathcal{E}_\kappa(t) \leq \mathcal{E}_\kappa(0) + t \sup_{0 \leq s \leq t} \mathcal{E}_\kappa(t).$$

In Sect. 2.8 we will prove an a priori bound for  $\mathcal{E}_\kappa$  independent of  $\kappa$  and show that the time of existence  $T$  is independent of  $\kappa$ . The bound (2.17) will then be justified a posteriori using the fundamental theorem of calculus, smallness of  $T_\kappa$ , and the definition of  $\mathcal{E}_\kappa$ . By choosing  $T_\kappa$  possibly smaller we assume that for certain  $\delta > 0$ ,

$$\min_{x' \in \Gamma} q_{,2}(t, x') > \delta \quad \text{and} \quad |\bar{\partial} h_\kappa(t, \cdot)|_\infty^2 \leq 1/2, \quad t \in [0, T_\kappa], \tag{2.18}$$

where  $(q, h)$  is the solution of the  $\kappa$ -problem (2.4). The first inequality is true by continuity-in-time of the energy  $\mathcal{E}_\kappa$  and the Taylor sign condition (2.11). The second inequality follows from the continuity-in-time and smallness of  $|\bar{\partial} h_0|_3$  (1.3).

## 2.4. The Energy Identities

In this section we collect the high-order energy identities in two lemmas stated below. We use the notation  $\mathcal{T}$  for those error terms which in a straightforward way are seen to satisfy the energy bound of the form:

$$\int_0^t |\mathcal{T}(s)| ds \lesssim tP(\mathcal{E}_\kappa);$$

this bound will then always follow from the standard  $L^\infty - L^2 - L^2$  type estimates. Here and in the rest of the paper  $P(\cdot)$  stands for a generic polynomial satisfying  $P(0) = 0$ .

**Lemma 2.2.** *Assume that  $(q, h)$  is a solution to the regularized Stefan problem (2.4) given by Theorem 2.1. Then the following identities hold:*

(i)

$$\begin{aligned} & \int_\Omega |\bar{\partial}^4 v|^2 + \frac{1}{2} \frac{d}{dt} \int_\Gamma (-q, 2) |\bar{\partial}^4 \Lambda_\kappa \Psi \cdot \mathfrak{A}_\bullet^2|^2 + \frac{1}{2} \frac{d}{dt} \int_\Omega (\bar{\partial}^4 q + \bar{\partial}^4 \Psi_\kappa \cdot v)^2 \\ & + \kappa^2 \int_\Gamma J_\kappa^{-1} |\bar{\partial}^4 h_t|^2 = \int_\Omega \mathcal{R}_1 + \int_\Gamma \mathcal{R}_2 + \mathcal{T}; \end{aligned} \quad (2.19)$$

$$\begin{aligned} & \int_\Omega |\bar{\partial}^2 \partial_t v|^2 + \frac{1}{2} \frac{d}{dt} \int_\Gamma (-q, 2) |\bar{\partial}^2 \partial_t \Lambda_\kappa \Psi \cdot \mathfrak{A}_\bullet^2|^2 + \frac{1}{2} \frac{d}{dt} \int_\Omega (\bar{\partial}^2 \partial_t q + \bar{\partial}^2 \partial_t \Psi_\kappa \cdot v)^2 \\ & + \kappa^2 \int_\Gamma J_\kappa^{-1} |\bar{\partial}^2 \partial_t h_t|^2 = \int_\Omega \mathcal{R}_3 + \int_\Gamma \mathcal{R}_4 + \mathcal{T}; \end{aligned} \quad (2.20)$$

$$\begin{aligned} & \int_\Omega |\partial_t^2 v|^2 + \frac{1}{2} \frac{d}{dt} \int_\Gamma (-q, 2) |\partial_t^2 \Lambda_\kappa \Psi \cdot \mathfrak{A}_\bullet^2|^2 + \frac{1}{2} \frac{d}{dt} \int_\Omega (\partial_t^2 q + \partial_t^2 \Psi_\kappa \cdot v)^2 \\ & + \kappa^2 \int_\Gamma J_\kappa^{-1} |\partial_{tt} h_t|^2 = \int_\Omega \mathcal{R}_5 + \int_\Gamma \mathcal{R}_6 + \mathcal{T}, \end{aligned} \quad (2.21)$$

where  $\mathcal{R}_i$ ,  $i = 1, \dots, 6$ , are error terms given below respectively by (2.41), (2.42), (2.43), (2.44), (2.45) and (2.46).

(ii)

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \int_\Omega |\bar{\partial}^3 v|^2 + \int_\Gamma (-q, 2) |\bar{\partial}^3 \Lambda_\kappa \Psi_t \cdot \mathfrak{A}_\bullet^2|^2 + \int_\Omega (\bar{\partial}^3 q_t + \bar{\partial}^3 \mathfrak{w} \cdot v)^2 \\ & + \frac{\kappa^2}{2} \frac{d}{dt} \int_\Gamma J_\kappa^{-1} |\bar{\partial}^3 h_t|^2 = \int_\Omega \mathcal{S}_1 + \int_\Gamma \mathcal{S}_2 + \mathcal{T}; \end{aligned} \quad (2.22)$$

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \int_\Omega |\bar{\partial} v_t|^2 + \int_\Gamma (-q, 2) |\bar{\partial} \Lambda_\kappa \Psi_{tt} \cdot \mathfrak{A}_\bullet^2|^2 + \int_\Omega (\bar{\partial} q_{tt} + \bar{\partial} \mathfrak{w}_t \cdot v)^2 \\ & + \frac{\kappa^2}{2} \frac{d}{dt} \int_\Gamma J_\kappa^{-1} |\bar{\partial} h_{tt}|^2 = \int_\Omega \mathcal{S}_3 + \int_\Gamma \mathcal{S}_4 + \mathcal{T}, \end{aligned} \quad (2.23)$$

where  $\mathcal{S}_i$ ,  $i = 1, \dots, 4$ , are error terms given below respectively by (2.51), (2.52), (2.53), (2.54).

We postpone the proof of Lemma 2.2 to Sect. 2.6.

2.5. *Equivalence of the Higher-Order Norm  $\mathcal{E}_\kappa$  and the Natural Energy Function  $\mathcal{F}_\kappa$*

By summing the left-hand sides of the identities (2.19)–(2.23) from Lemma 2.2, the *natural* coercive quadratic form  $\mathcal{F}_\kappa$  that arises as the energy takes the form

$$\begin{aligned}
 \mathcal{F}_\kappa = & \sum_{a+2b \leq 4} \|\bar{\partial}^a \partial_t^b v\|_{L_t^2 L_x^2}^2 + \frac{1}{2} \sum_{a+2b \leq 3} \|\bar{\partial}^a \partial_t^b v\|_{L_t^\infty L_x^2}^2 \\
 & + \kappa^2 \sum_{a+2b \leq 4} |J_\kappa^{-1/2} \bar{\partial}^a \partial_t^b h_t|_{L_t^2 L_x^2}^2 + \frac{\kappa^2}{2} \sum_{a+2b \leq 3} |J_\kappa^{-1} \bar{\partial}^a \partial_t^b h_t|_{L_t^\infty L_x^2}^2 \\
 & + \frac{1}{2} \sum_{a+2b \leq 4; } |\sqrt{q_{,2}} J_\kappa^{-1/2} \bar{\partial}^a \partial_t^b \Lambda_\kappa h|_{L_t^\infty L_x^2}^2 \\
 & + \sum_{a+2b \leq 3} |\sqrt{q_{,2}} J_\kappa^{-1/2} \bar{\partial}^a \partial_t^b \Lambda_\kappa h_t|_{L_t^2 L_x^2}^2 \\
 & + \frac{1}{2} \sum_{a+2b \leq 4; } \|\bar{\partial}^a \partial_t^b q + \bar{\partial}^a \partial_t^b \Psi_\kappa \cdot v\|_{L_t^\infty L_x^2}^2 \\
 & + \sum_{a+2b \leq 3; } \|\bar{\partial}^a \partial_t^b q_t + \bar{\partial}^a \partial_t^b \Psi_{\kappa t} \cdot v\|_{L_t^2 L_x^2}^2. \tag{2.24}
 \end{aligned}$$

The mathematical reason for imposing the Taylor sign condition (1.23) now becomes apparent. In order for the second line in the definition of  $\mathcal{F}_\kappa$  (2.24) above to make sense we must have

$$\min_{x' \in \Gamma} (q_{,2})(t, x', 0) > 0,$$

as was assumed in (1.23) for the (unregularized) classical Stefan problem. In order to perform the estimates in the next section, it is crucial to show that the energy  $\mathcal{E}_\kappa$  is bounded by  $\mathcal{F}_\kappa$ . To prove this statement we first establish the following temperature estimate.

**Lemma 2.3.** *Let  $(q, h)$  be a solution of the regularized problem (2.4) given by Theorem 2.1. Assume that the a priori assumption (2.17) holds on  $[0, T_\kappa]$ . Then:*

(i)

$$\sum_{a=0}^2 \|\partial_t^a q\|_{L_t^\infty H_x^{4-2a}}^2 \lesssim \mathcal{F}_\kappa \quad \text{on } [0, T_\kappa]. \tag{2.25}$$

(ii)

$$\|q\|_{L_t^2 H_x^{4.5}}^2 + \sum_{a=1}^2 \|\partial_t^a q\|_{L_t^2 H_x^{5-2a}}^2 \lesssim \mathcal{F}_\kappa \quad \text{on } [0, T_\kappa]. \tag{2.26}$$

**Proof.** We use elliptic regularity theory and the a priori assumption (2.17) to show that  $\|q\|_{L_t^\infty H_x^2} \lesssim \mathcal{F}_\kappa$  since  $\Delta \Psi_\kappa q = q_t + v \cdot \mathring{w} + \alpha$ ,  $\|q_t\|_{L_t^\infty L_x^2} \lesssim \mathcal{F}_\kappa$ ,  $\|v\|_{L_t^\infty L_x^2} \|w\|_{L_t^\infty L_x^2} \leq \|v\|_{L_t^\infty L_x^2} \|h_t\|_{L_t^\infty H_x^1} \lesssim \mathcal{F}_\kappa$ , and  $\|\alpha\|_{L_t^\infty L_x^2} \lesssim \mathcal{F}_\kappa(0) \lesssim \mathcal{F}_\kappa$ . Differentiating (2.4a) with respect to  $x^j$  ( $j = 1, 2$ ), we obtain that  $\Delta \Psi_\kappa q_{,j} = (\mathring{A}_i^m \mathring{A}_i^n)_{,j} q_{,mn} + (q_t + v \cdot \mathring{w})_{,j} + \alpha_{,j}$ . Furthermore, since  $q_{,j} = \Psi_{\kappa,j} \cdot v$  we have that

$$\begin{aligned} \|q_{,j}\| &\lesssim \|\Psi_{\kappa,j}\|_{L_t^\infty L_x^\infty} \|v\|_{L_t^\infty L_x^2} + \|\Psi_{\kappa,j}\|_{L_t^\infty L_x^\infty} \|v_t\|_{L_t^\infty L_x^2} \\ &\lesssim \|h_t\|_{L_t^\infty H_x^2} \|v\|_{L_t^\infty L_x^2} + \|\nabla \Psi_\kappa\|_{L_t^\infty H^{1.5}} \|v_t\|_{L_t^\infty L_x^2} \lesssim \mathcal{F}_\kappa, \end{aligned}$$

where we have used the trace bound (2.2) and the a priori assumption (2.17). Note that

$$\begin{aligned} \|(v \cdot \mathring{w})_{,j}\|_{L_t^\infty H_x^1} &\lesssim \|v\|_{L_t^\infty H_x^1} \|\mathring{w}\|_{L_t^\infty L_x^\infty} + \|v\|_{L_t^\infty L_x^2} \|\mathring{w}_{,j}\|_{L_t^\infty L_x^\infty} \\ &\lesssim \|q\|_{L_t^\infty H_x^2} \|h_{\kappa,t}\|_{L_t^\infty H_x^1} + \|q\|_{L_t^\infty H_x^2} \|h_{\kappa,t}\|_{L_t^\infty H_x^2} \lesssim \mathcal{F}_\kappa, \end{aligned}$$

where we have used the bound (2.17) in the last estimate. It is easy to see that  $\|(\mathring{A}_i^m \mathring{A}_i^n)_{,j} q_{,mn}\|_{L_t^\infty L_x^2} \lesssim \mathcal{F}_\kappa (1 + P(\mathcal{A}_\kappa)) \lesssim \mathcal{F}_\kappa$ , where  $P$  stands for a generic polynomial. Finally,  $\|\nabla \alpha\|_{L_t^\infty L_x^2} \lesssim \mathcal{F}_\kappa$ . Thus, by the elliptic theory again, we conclude

$$\|q\|_{L_t^\infty H_x^3}^2 \lesssim \mathcal{F}_\kappa.$$

Differentiating (2.4a) with respect to  $t$ , we obtain  $\Delta \Psi_\kappa q_t = -(\mathring{A}_i^m \mathring{A}_i^n)_{,t} q_{,mn} + v_{tt} + (v \cdot \mathring{w})_t$  (since  $\alpha$  is independent of  $t$ ). Again, using  $\|v_{tt}\|_{L_t^\infty L_x^2}^2 \lesssim \mathcal{F}_\kappa$ , the previous estimates and the bound  $\|h_{\kappa,t}\|_{L_t^\infty H_x^2}^2 \lesssim \mathcal{F}_\kappa$ , elliptic regularity implies  $\|q_t\|_{L_t^\infty H_x^2} \lesssim \mathcal{F}_\kappa$ . Furthermore  $\|q_{tt}\|_{L_t^\infty L_x^2}^2 \lesssim \|q_{tt}\|_{L_t^\infty L_x^2}^2 + \|\mathring{w}_t \cdot v\|_{L_t^\infty L_x^2}^2 + \|\mathring{w}_t \cdot v\|_{L_t^\infty L_x^2}^2 \lesssim \mathcal{F}_\kappa$ . The last equality follows from the third line on the definition (2.16) of  $\mathcal{F}_\kappa$  and a simple bound on the  $L_t^\infty L_x^\infty$ -norm of  $v$ , which follows from Sobolev embedding. Finally, choose any  $j, k \in \{1, 2\}$ . Applying  $\partial_{x^j} \partial_{x^k}$  to (2.4a), we arrive at the elliptic equation

$$\begin{aligned} \Delta \Psi_\kappa q_{,jk} &= -(\mathring{A}_i^m \mathring{A}_i^n)_{,j} q_{,mnk} - (\mathring{A}_i^m \mathring{A}_i^n)_{,k} q_{,mnj} \\ &\quad - (\mathring{A}_i^m \mathring{A}_i^n)_{,jk} q_{,mn} + (q_t + v \cdot \mathring{w})_{,jk} + \alpha_{,jk}. \end{aligned}$$

By the estimates already derived above, (2.16), (2.6), and (2.17), the right-hand side is bounded by  $\mathcal{F}_\kappa (1 + P(\mathcal{A}_\kappa)) \lesssim \mathcal{F}_\kappa$  in  $L_t^\infty L_x^2$ -norm. Thus, by elliptic regularity, we finally conclude  $\|q\|_{L_t^\infty H^4} \lesssim \mathcal{F}_\kappa$ , concluding the proof of (2.25).

To prove (2.26) we start with the easiest case  $b = 2$ . For  $j = 1, 2$ , we have

$$q_{,jtt} = (\Psi_{\kappa,j} \cdot v)_{tt} = \Psi_{\kappa,jtt} \cdot v + 2\Psi_{\kappa,jt} \cdot v_t + \Psi_{\kappa,j} \cdot v_{tt}.$$

From the above we easily infer that

$$\begin{aligned} \int_0^t \|\nabla q_{tt}\|_0^2 d\tau &\lesssim \int_0^t \left( \|\nabla \Psi_{\kappa,tt}\|_0^2 \|v\|_\infty^2 + \|\Psi_{\kappa,jt}\|_\infty^2 \|v_t\|_0^2 + \|\Psi_{\kappa,j}\|_\infty^2 \|v_{tt}\|_0^2 \right) d\tau \\ &\lesssim P(\mathcal{F}_\kappa). \end{aligned} \tag{2.27}$$

We have thereby used the trace estimate (2.2) to obtain

$$\|\nabla \Psi_{\kappa, tt}\|_0^2 \lesssim |h_{\kappa, tt}|_{0,5}^2 \lesssim \mathcal{F}_\kappa,$$

where we have used the definition (2.24) in the last bound above. On the other hand, using the a priori bound (2.17) and the Sobolev embedding we conclude that  $\|v\|_\infty \lesssim \|v\|_1 \lesssim \|q\|_2 \leq \mathcal{E}_\kappa(0) + 1 \lesssim 1$ . The remaining two terms on the right-hand side of (2.27) are estimated in a similar fashion. If  $b = 1$ , we apply the same ideas using (2.25), (2.17), and the Sobolev embeddings. To prove  $\|q\|_{L_t^2 H_x^{4.5}}^2 \lesssim \mathcal{F}_\kappa$  we need to use an interpolation estimate. The strategy consists of estimating  $\|q\|_{L_t^2 H_x^5}^2$  and  $\|q\|_{L_t^2 H_x^4}^2$  separately and then interpolating between the two estimates. The reader may consult [31] for the details.  $\square$

**Remark 8.** The regularity of  $q \in L_t^2 H_x^{4.5}$  can in fact be improved.

**Lemma 2.4.** (Optimal regularity for  $\Psi_\kappa$  and  $q$ ) *Suppose that the pair  $(q, h)$  is a solution of the  $\kappa$ -problem (2.4) given by Theorem 2.1, and that the basic assumption (2.17) holds on  $[0, T_\kappa]$ . Then  $\int_0^{T_\kappa} (\|\Psi_\kappa\|_5^2 + \|q\|_5^2) dt \leq C\mathcal{E}_\kappa(t)$ .*

**Proof.** *Step 1.* We will first prove that  $\int_0^{T_\kappa} \|\Psi_\kappa\|_5^2 dt \leq C\mathcal{E}(t)$ . Since  $q = 0$  on  $\Gamma$  it follows that

$$v(x, t) \cdot \tau_\kappa(x, t) = 0 \text{ on } \Gamma, \quad (2.28)$$

where we recall that  $\tau_\kappa$  is the unit tangent vector to  $\Psi_\kappa(t, \Gamma)$ . Applying the horizontal derivative  $\bar{\partial}$  to (2.28), and using the fact that  $\bar{\partial}\tau_\kappa = g_\kappa^{-1}\bar{\partial}^2\Psi_\kappa \cdot n_\kappa n_\kappa$  and that  $\bar{\partial}^2\Psi_\kappa \cdot n_\kappa = -g_\kappa^{-1}\bar{\partial}^2 h_\kappa$ , we find that

$$\bar{\partial}^2 h_\kappa = \frac{g_\kappa^2 \bar{\partial} v \cdot \tau_\kappa}{v \cdot n_\kappa}. \quad (2.29)$$

The dominator in (2.32) is strictly positive for  $T_\kappa$  small enough by the Taylor sign condition (2.11). For any  $W : \Omega \rightarrow \mathbb{R}^2$  we define

$$\text{curl}_\Psi W = \varepsilon_{ji} A_j^s W^i, \quad (2.30)$$

where  $\varepsilon_{21} = -\varepsilon_{12} = 1$ ,  $\varepsilon_{11} = \varepsilon_{22} = 0$ . By the tangential trace inequality (see [11]),

$$\left| \bar{\partial}^4 v \cdot \tau_\kappa \right|_{H^{-\frac{1}{2}}(\Gamma)} \lesssim \|\text{curl}_\Psi \bar{\partial}^3 v\|_{L^2(\Omega)} + \|\bar{\partial}^4 v\|_{L^2(\Omega)}. \quad (2.31)$$

We observe that

$$\begin{aligned} \text{curl}_\Psi \bar{\partial}^3 v &= \bar{\partial}^3 (\text{curl}_\Psi v) - \varepsilon_{ji} \sum_{m=1}^3 c_m \bar{\partial}^m A_j^s \bar{\partial}^{3-m} v^j, \\ &= -\varepsilon_{ji} \sum_{m=1}^3 c_m \bar{\partial}^m A_j^s \bar{\partial}^{3-m} v^j, \end{aligned}$$



where we have used the identity  $\operatorname{curl}_\Psi v = (\operatorname{curl} \nabla p) \circ \Psi = 0$ . Using the Cauchy–Schwarz inequality and the definition of  $\mathcal{E}_\kappa$ , we obtain

$$\|\operatorname{curl}_\Psi \bar{\partial}^3 v\|_0 \lesssim \sqrt{\mathcal{E}_\kappa}.$$

From (2.31) and the definition (2.16) of  $\mathcal{E}_\kappa$ , we obtain

$$\int_0^t |\bar{\partial}^4 v \cdot \tau_\kappa|_{H^{-\frac{1}{2}}(\Gamma)}^2 d\sigma \lesssim \mathcal{E}_\kappa(t), \quad 0 \leq t \leq T_\kappa. \quad (2.32)$$

Using (2.32) and (2.31) it follows easily that  $\int_0^t |h_\kappa|_{4.5}^2 d\sigma \leq \mathcal{E}_\kappa(t)$ ,  $0 \leq t \leq T_\kappa$ . Recalling that  $\Psi_\kappa$  is the harmonic extension of  $(x', \bar{h}_\kappa(x'))$ ,  $x' \in \Gamma$  the optimal trace inequality (1.8) implies that  $\int_0^t \|\Psi_\kappa\|_5^2 d\sigma \leq C\mathcal{E}_\kappa(t)$  for any  $t \in [0, T_\kappa]$ .

*Step 2.* The fact that  $\int_0^{T_\kappa} \|q\|_5^2 dt \leq C\mathcal{E}(t)$  follows from Step 1 and the elliptic regularity result in Theorem 3.6 in [11].  $\square$

As a consequence of Lemmas 2.3 and 2.4 we obtain the following key bound between the norm  $\mathcal{E}_\kappa$  and the energy  $\mathcal{F}_\kappa$ .

**Proposition 2.5.** (Norm-energy equivalence) *Let  $(q, h)$  be a solution of the  $\kappa$ -problem (2.4) given by Theorem 2.1. Assume that the a priori assumption (2.17) holds on  $[0, T_\kappa]$ . Then  $\mathcal{E}_\kappa \lesssim \mathcal{F}_\kappa$  on  $[0, T_\kappa]$ .*

**Proof.** Due to the Taylor sign condition (2.18), the boundary integrals in  $\mathcal{E}_\kappa$  and  $\mathcal{F}_\kappa$  satisfy

$$\begin{aligned} & \sum_{b=0}^2 |\Lambda_\kappa h|_{L_t^\infty H_x^{4-2b}}^2 + \sum_{b=0}^1 |\Lambda_\kappa h_t|_{L_t^2 H_x^{3-2b}}^2 \\ & \lesssim \sum_{a+2b \leq 4; } |\sqrt{-q} \cdot 2\bar{\partial}^a \partial_t^b \Lambda_\kappa h|_{L_t^\infty L_x^2}^2 + \sum_{a+2b \leq 3} |\sqrt{-q} \cdot 2\bar{\partial}^a \partial_t^b \Lambda_\kappa h_t|_{L_t^2 L_x^2}^2. \end{aligned}$$

The remaining estimates now follow directly from Lemmas 2.3 and 2.4.

## 2.6. Proof of Lemma 2.2

*Proof of part (i) of Lemma 2.2.* Applying the tangential differential operator  $\bar{\partial}^4$  to the equation (2.4b), multiplying it by  $\bar{\partial}^4 v^i$  and integrating over  $\Omega$ , we obtain

$$(\bar{\partial}^4 v^i + \bar{\partial}^4 \mathbb{A}_i^k q_{,k} + \mathbb{A}_i^k \bar{\partial}^4 q_{,k}, \bar{\partial}^4 v^i)_{L^2} = \sum_{l=1}^3 c_l (\bar{\partial}^l \mathbb{A}_i^k \bar{\partial}^{4-l} q_{,k}, \bar{\partial}^4 v^i)_{L^2}, \quad (2.33)$$

where  $c_l = \binom{4}{l}$ . Recalling (1.17), we write

$$\bar{\partial}^4 \mathbb{A}_i^k = -\mathbb{A}_i^s \bar{\partial}^4 \Psi_{\kappa,s}^r \mathbb{A}_r^k + \{\bar{\partial}^4, \mathbb{A}_i^k\}, \quad (2.34)$$

where  $\{\bar{\partial}^4, \mathbb{A}_i^k\}$  stands for the lower order commutator defined in (1.17). With this identity, we obtain

$$\begin{aligned}
 (\bar{\partial}^4 \mathbb{A}_i^k q_{,k}, \bar{\partial}^4 v^i)_{L^2(\Omega)} &= -(\mathbb{A}_i^s \bar{\partial}^4 \Psi_{\kappa,s}^r \mathbb{A}_r^k q_{,k}, \bar{\partial}^4 v^i)_{L^2(\Omega)} + (\{\bar{\partial}^4, \mathbb{A}_i^k\} q_{,k}, \bar{\partial}^4 v^i)_{L^2(\Omega)} \\
 &= - \int_{\Gamma} q_{,k} \mathbb{A}_i^s \bar{\partial}^4 \Psi_{\kappa}^r \mathbb{A}_r^k \bar{\partial}^4 v^i N^s + \int_{\Omega} \mathbb{A}_i^s \bar{\partial}^4 \Psi_{\kappa}^r \mathbb{A}_r^k q_{,k} \bar{\partial}^4 v_{,s}^i + \mathcal{T} \\
 &= - \int_{\Gamma} q_{,k} \mathbb{A}_i^s \bar{\partial}^4 \Psi_{\kappa}^r \mathbb{A}_r^k \bar{\partial}^4 v^i N^s - \int_{\Omega} \mathbb{A}_i^s \bar{\partial}^4 \Psi_{\kappa}^r v^r \bar{\partial}^4 v_{,s}^i + \mathcal{T},
 \end{aligned} \tag{2.35}$$

where we have used  $(\mathbb{A}_i^s)_{,s} = 0$  and the identity  $v^r = -\mathbb{A}_r^k q_{,k}$  to write the last line more concisely. Furthermore, integrating by parts with respect to  $x^k$ ,

$$(\mathbb{A}_i^k \bar{\partial}^4 q_{,k}, \bar{\partial}^4 v^i)_{L^2} = \int_{\Omega} \mathbb{A}_i^k \bar{\partial}^4 q_{,k} \bar{\partial}^4 v^i = \int_{\Gamma} \mathbb{A}_i^k \bar{\partial}^4 q \bar{\partial}^4 v^i \cdot N^k - \int_{\Omega} \mathbb{A}_i^k \bar{\partial}^4 q \bar{\partial}^4 v_{,k}^i. \tag{2.36}$$

Note that the the boundary contribution coming from the fixed boundary  $\partial\Omega_{\text{top}}$  vanishes due to the boundary condition (2.4e), which further reduces to  $v^2 = 0$  on  $\partial\Omega_{\text{top}}$ . Summing (2.35) and (2.36), we obtain

$$\begin{aligned}
 (\bar{\partial}^4 \mathbb{A}_i^k q_{,k} + \mathbb{A}_i^k \bar{\partial}^4 q_{,k}, \bar{\partial}^4 v^i)_{L^2(\Omega)} &= - \int_{\Gamma} q_{,k} \mathbb{A}_i^s \bar{\partial}^4 \Psi_{\kappa}^r \mathbb{A}_r^k \bar{\partial}^4 v^i N^s \\
 &\quad + \int_{\Gamma} \mathbb{A}_i^k \bar{\partial}^4 q \bar{\partial}^4 v^i \cdot N^k - \int_{\Omega} \mathbb{A}_i^k \bar{\partial}^4 v_{,k}^i (\bar{\partial}^4 q + \bar{\partial}^4 \Psi_{\kappa} \cdot v) + \mathcal{T}.
 \end{aligned} \tag{2.37}$$

The first three terms on the right-hand side of (2.37) will be the source of positive definite quadratic contributions to the energy. To extract the quadratic coercive contribution from the first integral on the right-hand side of (2.37), we simplify it to

$$\begin{aligned}
 - \int_{\Gamma} q_{,k} \mathbb{A}_i^s \bar{\partial}^4 \Psi_{\kappa}^r \mathbb{A}_r^k \bar{\partial}^4 v^i N^s &= \int_{\Gamma} q_{,2} \mathbb{A}_i^2 \bar{\partial}^4 \Psi_{\kappa}^r \mathbb{A}_r^2 \bar{\partial}^4 v^i + \int_{\Gamma} q_{,1} \mathbb{A}_i^2 \bar{\partial}^4 \Psi_{\kappa}^r \mathbb{A}_r^1 \bar{\partial}^4 v^i \\
 &= \int_{\Gamma} q_{,2} \bar{\partial}^4 \Psi_{\kappa} \cdot \mathbb{A}_{\bullet}^2 \bar{\partial}^4 v \cdot \mathbb{A}_{\bullet}^2 + \int_{\Gamma} q_{,1} \bar{\partial}^4 \Psi_{\kappa}^r \mathbb{A}_r^1 \bar{\partial}^4 v \cdot \mathbb{A}_{\bullet}^2.
 \end{aligned}$$

We rewrite the expression  $\bar{\partial}^4 v \cdot \mathbb{A}_{\bullet}^2$  and thereby use the boundary condition (2.4d):

$$\begin{aligned}
 \bar{\partial}^4 v \cdot \mathbb{A}_{\bullet}^2 &= \bar{\partial}^4 w \cdot \mathbb{A}_{\bullet}^2 + \bar{\partial}^4 (v + w) \cdot \mathbb{A}_{\bullet}^2 \\
 &= \bar{\partial}^4 w \cdot \mathbb{A}_{\bullet}^2 + \bar{\partial}^4 \left( \underbrace{(v + w) \cdot \mathbb{A}_{\bullet}^2}_{=0} \right) - \sum_{l=1}^4 a_l \bar{\partial}^{4-l} (v + w) \cdot \bar{\partial}^l \mathbb{A}_{\bullet}^2 \\
 &= \bar{\partial}^4 w \cdot \mathbb{A}_{\bullet}^2 - \sum_{l=1}^4 a_l \bar{\partial}^{4-l} (v + w) \cdot \bar{\partial}^l \mathbb{A}_{\bullet}^2.
 \end{aligned}$$

Due to the above identity and recalling  $\Psi_\kappa = \Lambda_\kappa \Lambda_\kappa \Psi$ , we obtain

$$\begin{aligned} \int_\Gamma q_{,2} \bar{\partial}^4 \Psi_\kappa \cdot n_\kappa \bar{\partial}^4 v \cdot \mathbb{A}_\kappa^2 &= \int_\Gamma q_{,2} \bar{\partial}^4 \Lambda_\kappa \Lambda_\kappa \Psi \cdot \mathbb{A}_\bullet^2 \bar{\partial}^4 w \cdot \mathbb{A}_\bullet^2 \\ &\quad - \sum_{l=1}^4 a_l \int_\Gamma q_{,2} \bar{\partial}^4 \Psi_\kappa \cdot \mathbb{A}_\bullet^2 \bar{\partial}^{4-l} (v+w) \cdot \bar{\partial}^l \mathbb{A}_\bullet^2. \end{aligned} \quad (2.38)$$

The first term on the right-hand side of (2.38) is rewritten in the following way:

$$\begin{aligned} \int_\Gamma q_{,2} \bar{\partial}^4 \Lambda_\kappa \Lambda_\kappa \Psi \cdot \mathbb{A}_\bullet^2 \bar{\partial}^4 w \cdot \mathbb{A}_\bullet^2 &= \int_\Gamma q_{,2} \Lambda_\kappa \bar{\partial}^4 \Psi \cdot \mathbb{A}_\bullet^2 \Lambda_\kappa \bar{\partial}^4 \Psi_t \cdot \mathbb{A}_\bullet^2 \\ &\quad + \int_\Gamma \bar{\partial}^4 \Lambda_\kappa \Psi \left[ \Lambda_\kappa [q_{,2} (\bar{\partial}^4 \Psi_t \cdot \mathbb{A}_\bullet^2) \mathbb{A}_\bullet^2] - q_{,2} (\bar{\partial}^4 \Lambda_\kappa \Psi_t \cdot \mathbb{A}_\bullet^2) \mathbb{A}_\bullet^2 \right] \\ &= \frac{1}{2} \partial_t \int_\Gamma q_{,2} |\bar{\partial}^4 \Lambda_\kappa \Psi \cdot \mathbb{A}_\bullet^2|^2 - \frac{1}{2} \int_\Gamma q_{,2t} |\bar{\partial}^4 \Lambda_\kappa \Psi \cdot \mathbb{A}_\bullet^2|^2 \\ &\quad - \int_\Gamma q_{,2} \bar{\partial}^4 \Lambda_\kappa \Psi \cdot \mathbb{A}_\bullet^2 \bar{\partial}^4 \Lambda_\kappa \Psi \cdot \mathbb{A}_\bullet^2 \\ &\quad + \int_\Gamma \Lambda_\kappa \bar{\partial}^4 \Psi \left[ \Lambda_\kappa [q_{,2} (\bar{\partial}^4 \Psi_t \cdot \mathbb{A}_\bullet^2) \mathbb{A}_\bullet^2] - q_{,2} (\Lambda_\kappa \bar{\partial}^4 \Psi_t \cdot \mathbb{A}_\bullet^2) \mathbb{A}_\bullet^2 \right] \\ &= \frac{1}{2} \partial_t \int_\Gamma q_{,2} |\bar{\partial}^4 \Lambda_\kappa \Psi \cdot \mathbb{A}_\bullet^2|^2 + \int_\Gamma \Lambda_\kappa \bar{\partial}^4 \Psi \left[ \Lambda_\kappa [q_{,2} (\bar{\partial}^4 \Psi_t \cdot \mathbb{A}_\bullet^2) \mathbb{A}_\bullet^2] \right. \\ &\quad \left. - q_{,2} (\Lambda_\kappa \bar{\partial}^4 \Psi_t \cdot \mathbb{A}_\bullet^2) \mathbb{A}_\bullet^2 \right] + \mathcal{T}. \end{aligned} \quad (2.39)$$

The second term on the right-hand side of (2.37) turns into

$$\begin{aligned} \int_\Gamma \mathbb{A}_i^k \bar{\partial}^4 q \bar{\partial}^4 v^i \cdot N^k &= - \int_\Gamma \bar{\partial}^4 q \bar{\partial}^4 v \cdot \mathbb{A}_\bullet^2 \\ &= \kappa^2 \int_\Gamma \bar{\partial}^4 h_t (\bar{\partial}^4 (v \cdot \mathbb{A}_\bullet^2) - \sum_{l=0}^3 a_l \bar{\partial}^l v \bar{\partial}^{4-l} \mathbb{A}_\bullet^2) \\ &\quad + \kappa^2 \int_\Gamma \beta(t, x') \bar{\partial}^4 v \cdot \mathbb{A}_\bullet^2 \\ &= \kappa^2 \int_\Gamma J_\kappa^{-1} |\bar{\partial}^4 h_t|^2 - \kappa^2 \sum_{l=0}^3 a_l \int_\Gamma \bar{\partial}^4 h_t \bar{\partial}^l v \bar{\partial}^{4-l} \mathbb{A}_\bullet^2 \\ &\quad + \kappa^2 \int_\Gamma \bar{\partial}^4 \beta(t, x') \bar{\partial}^4 v \cdot \mathbb{A}_\bullet^2. \end{aligned}$$

where we have used the boundary condition (2.4c) in the second equality above (recall  $v \cdot \mathbb{A}_\bullet^2 = w \cdot \mathbb{A}_\bullet^2 = h_t$ ). As to the third term on the right-hand side of (2.37), note that

$$\begin{aligned}
 \mathfrak{A}_i^k \bar{\partial}^4 v^i{}_{,k} &= \bar{\partial}^4 (\mathfrak{A}_i^k v^i{}_{,k}) - \sum_{l=1}^4 c_l \bar{\partial}^l \mathfrak{A}_i^k \bar{\partial}^{4-l} v^i{}_{,k} \\
 &= -\bar{\partial}^4 (q_t + v \cdot \mathfrak{w}) - \sum_{l=1}^4 c_l \bar{\partial}^l \mathfrak{A}_i^k \bar{\partial}^{4-l} v^i{}_{,k},
 \end{aligned}$$

where  $\mathfrak{A}_i^k v^i{}_{,k} = -\operatorname{div}_{\Psi_\kappa} v = -(q_t + v \cdot \mathfrak{w}) + \alpha$  by the parabolic equation (2.4a). Thus

$$\begin{aligned}
 & - \int_{\Omega} \mathfrak{A}_i^k \bar{\partial}^4 v^i{}_{,k} (\bar{\partial}^4 q + \bar{\partial}^4 \Psi_\kappa \cdot v) \\
 &= \int_{\Omega} \bar{\partial}^4 (q_t + \Psi_{\kappa t} \cdot v - \alpha) (\bar{\partial}^4 q + \bar{\partial}^4 \Psi_\kappa \cdot v) \\
 & \quad + \sum_{l=1}^4 c_l \int_{\Omega} \bar{\partial}^l \mathfrak{A}_i^k \bar{\partial}^{3-l} v^i{}_{,k} (\bar{\partial}^4 q + \bar{\partial}^4 \Psi_\kappa \cdot v) \\
 &= \frac{1}{2} \partial_t \int_{\Omega} (\bar{\partial}^4 q + \bar{\partial}^4 \Psi_\kappa \cdot v)^2 \\
 & \quad + \int_{\Omega} \left( \sum_{l=1}^4 d_l \bar{\partial}^{4-l} \Psi_{\kappa t} \cdot \bar{\partial}^l v - \bar{\partial}^4 \Psi_\kappa \cdot v_t \right) (\bar{\partial}^4 q + \bar{\partial}^4 \Psi_\kappa \cdot v) \\
 & \quad + \sum_{l=1}^4 c_l \int_{\Omega} \bar{\partial}^l \mathfrak{A}_i^k \bar{\partial}^{3-l} v^i{}_{,k} (\bar{\partial}^4 q + \bar{\partial}^4 \Psi_\kappa \cdot v) + \int_{\Omega} \bar{\partial}^4 \alpha (\bar{\partial}^4 q + \bar{\partial}^4 \Psi_\kappa \cdot v).
 \end{aligned} \tag{2.40}$$

Combining (2.37), (2.38), (2.39) and (2.40) we obtain the identity (2.19) with the error terms  $\mathcal{R}_1$  and  $\mathcal{R}_2$  given by:

$$\begin{aligned}
 \mathcal{R}_1 &:= \sum_{l=1}^3 c_l \bar{\partial}^l \mathfrak{A}_i^k \bar{\partial}^{4-l} q_{,k} \bar{\partial}^4 v^i - \left( \sum_{l=1}^4 c_l \bar{\partial}^l \mathfrak{A}_i^k \bar{\partial}^{4-l} v^i{}_{,k} \right. \\
 & \quad + \sum_{l=1}^4 d_l \bar{\partial}^{4-l} \mathfrak{w} \cdot \bar{\partial}^l v - \bar{\partial}^4 \Psi_\kappa \cdot v_t \left. \right) (\bar{\partial}^4 q + \bar{\partial}^4 \Psi_\kappa \cdot v) \\
 & \quad + \bar{\partial}^4 \alpha (\bar{\partial}^4 q + \bar{\partial}^4 \Psi_\kappa \cdot v);
 \end{aligned} \tag{2.41}$$

$$\begin{aligned}
 \mathcal{R}_2 &:= -\Lambda_\kappa \bar{\partial}^4 \Psi \left[ \Lambda_\kappa [(-q, {}_2) (\bar{\partial}^4 \Psi_t \cdot \mathfrak{A}_\bullet^2) \mathfrak{A}_\bullet^2] - (-q, {}_2) (\bar{\partial}^4 \Lambda_\kappa \Psi_t \cdot \mathfrak{A}_\bullet^2) \mathfrak{A}_\bullet^2 \right] \\
 & \quad + \sum_{l=1}^4 a_l (-q, {}_2) \bar{\partial}^4 \Psi_\kappa \cdot \mathfrak{A}_\bullet^2 \bar{\partial}^{4-l} (v + w) \cdot \bar{\partial}^l \mathfrak{A}_\bullet^2 \\
 & \quad + q_{,1} \bar{\partial}^4 \Psi_\kappa^r \mathfrak{A}_r^1 \bar{\partial}^4 v \cdot \mathfrak{A}_\bullet^2 + \kappa^2 \sum_{l=0}^3 a_l \bar{\partial}^4 h_t \bar{\partial}^l v \bar{\partial}^{4-l} \mathfrak{A}_\bullet^2 \\
 & \quad - \kappa^2 \bar{\partial}^4 \beta \bar{\partial}^4 v \cdot \mathfrak{A}_\bullet^2.
 \end{aligned} \tag{2.42}$$

Applying the tangential differential operator  $\bar{\partial}^2 \partial_t$  to the equation (2.4b), multiplying it by  $\bar{\partial}^2 \partial_t v^i$  and integrating over  $\Omega$ , we obtain, in a completely analogous fashion, identity (2.20), claimed in Lemma 2.2 with error terms  $\mathcal{R}_3$  and  $\mathcal{R}_4$  given by:

$$\begin{aligned} \mathcal{R}_3 := & \sum_{1 \leq m+n \leq 2} c_{mn} \bar{\partial}^m \partial_t^n \mathbb{A}_i^k \bar{\partial}^{2-m} \partial_t^{1-n} q_{,k} \bar{\partial}^2 \partial_t v^i \\ & - \left( \sum_{1 \leq m+n \leq 2} c_{mn} \bar{\partial}^m \partial_t^n \mathbb{A}_i^k \bar{\partial}^{2-m} \partial_t^{1-n} v^i_{,k} \right. \\ & + \sum_{0 \leq m+n \leq 2} d_{mn} \bar{\partial}^m \partial_t^n \Psi_{\kappa t} \cdot \bar{\partial}^{2-m} \partial_t^{1-n} v - \bar{\partial}^2 \Psi_{\kappa t} \cdot v_t \\ & \left. \times (\bar{\partial}^2 q_t + \bar{\partial}^2 \Psi_{\kappa t} \cdot v) \right); \end{aligned} \quad (2.43)$$

$$\begin{aligned} \mathcal{R}_4 := & -\Lambda_\kappa \bar{\partial}^2 \Psi_t \left[ \Lambda_\kappa [q, 2 (\bar{\partial}^2 \partial_t \Psi_t \cdot \mathbb{A}_\bullet^2) \mathbb{A}_\bullet^2] - q, 2 (\bar{\partial}^2 \partial_t \Lambda_\kappa \Psi_t \cdot \mathbb{A}_\bullet^2) \mathbb{A}_\bullet^2 \right] \\ & + \sum_{l+l' \geq 1} a_{l,l'} \int_\Gamma q, 2 \bar{\partial}^l \partial_t^{l'} \Psi_\kappa \cdot \mathbb{A}_\bullet^2 \bar{\partial}^{2-l} \partial_t^{1-l'} (v+w) \cdot \partial_t^l \mathbb{A}_\bullet^2 \\ & - q, 1 \bar{\partial}^2 \partial_t \Psi_\kappa \mathbb{A}_r^1 \bar{\partial}^2 \partial_t v \cdot \mathbb{A}_\bullet^2 + \kappa^2 \sum_{0 \leq l+l' < 3} \bar{\partial}^2 \partial_t h_t \bar{\partial}^l \partial_t^{l'} v \cdot \bar{\partial}^{2-l} \partial_t^{1-l'} \mathbb{A}_\bullet^2 \\ & - \kappa^2 \bar{\partial}^2 \partial_t \beta \bar{\partial}^2 \partial_t v \cdot \mathbb{A}_\bullet^2. \end{aligned} \quad (2.44)$$

Finally, applying  $\partial_{tt}$  to the equation (2.4b), multiplying it by  $\partial_{tt} v^i$  and integrating over  $\Omega$ , the last identity (2.21) of Lemma 2.2 follows with error terms  $\mathcal{R}_5$  and  $\mathcal{R}_6$  given by

$$\begin{aligned} \mathcal{R}_5 := & -(\mathbb{A}_i^k{}_{,tt} v^i{}_{,k} + 2\mathbb{A}_i^k{}_{,t} v^i{}_{,kt} + 2v_t \cdot \mathbb{A}_t + v_{tt} \cdot \mathbb{A}_w \\ & - \Psi_{\kappa tt} \cdot v_t)(q_{tt} + \Psi_{\kappa tt} \cdot v); \end{aligned} \quad (2.45)$$

$$\begin{aligned} \mathcal{R}_6 := & -\Lambda_\kappa \Psi_{tt} \left[ \Lambda_\kappa [q, 2 (\Psi_{tt} \cdot \mathbb{A}_\bullet^2) \mathbb{A}_\bullet^2] - q, 2 (\partial_{tt} \Lambda_\kappa \Psi_t \cdot \mathbb{A}_\bullet^2) \mathbb{A}_\bullet^2 \right] \\ & + \sum_{l=1}^2 a_l^2 \int_\Gamma q, 2 \partial_t^2 \Psi_\kappa \cdot \mathbb{A}_\bullet^2 \partial_t^{2-l} (v+w) \cdot \partial_t^l \mathbb{A}_\bullet^2 - q, 1 \partial_{tt} \Psi_\kappa \mathbb{A}_r^1 \partial_{tt} v \cdot \mathbb{A}_\bullet^2 \\ & + \kappa^2 \sum_{l'=0}^1 \partial_{tt} h_t \partial_t^{l'} v \partial_t^{2-l'} \mathbb{A}_\bullet^2 \\ & - \kappa^2 \partial_{tt} \beta \partial_{tt} v \cdot \mathbb{A}_\bullet^2. \end{aligned} \quad (2.46)$$

*Proof of part (ii) of Lemma 2.2.* Applying the tangential operator  $\bar{\partial}^3 \partial_t$  to the equation (2.4b), multiplying by  $\bar{\partial}^3 v^i$  and integrating over  $\Omega$  we obtain

$$(\bar{\partial}^3 \partial_t v^i, \bar{\partial}^3 v^i)_{L^2} + (\bar{\partial}^3 \partial_t (\mathbb{A}_i^k q, k), \bar{\partial}^3 v^i)_{L^2} = 0,$$

implying

$$\begin{aligned} & \frac{1}{2} \partial_t \int_\Omega |\bar{\partial}^3 v^i|^2 + (\bar{\partial}^3 \partial_t \mathbb{A}_i^k q, k + \mathbb{A}_i^k \bar{\partial}^3 \partial_t q, k, \bar{\partial}^3 v^i)_{L^2} \\ & = \sum_{\substack{l+\bar{l}=3, k+\bar{k}=1 \\ 0 < l+k < 4}} c_{l,k,\bar{l},\bar{k}} (\bar{\partial}^l \partial_t^k \mathbb{A}_i^k \bar{\partial}^{\bar{l}} \partial_t^{\bar{k}} q, k, \bar{\partial}^3 v^i)_{L^2}. \end{aligned} \quad (2.47)$$

Recalling (1.17), we write

$$\bar{\partial}^3 \partial_t \mathbb{A}_i^k = -\mathbb{A}_r^k \bar{\partial}^3 \mathbb{w}_{,s}^r \mathbb{A}_i^s + \{\bar{\partial}^3 \partial_t, \mathbb{A}_i^k\}. \quad (2.48)$$

Using this decomposition we have

$$\begin{aligned} (\bar{\partial}^3 \partial_t \mathbb{A}_i^k q_{,k} + \mathbb{A}_i^k \bar{\partial}^3 \partial_t q_{,k}, \bar{\partial}^3 v^i)_{L^2} &= - \int_{\Omega} \mathbb{A}_r^k \bar{\partial}^3 \mathbb{w}_{,s}^r \mathbb{A}_i^s q_{,k} \bar{\partial}^3 v^i \\ &\quad + \int_{\Omega} \mathbb{A}_i^k \bar{\partial}^3 \partial_t q_{,k} \bar{\partial}^3 v^i + \mathcal{T}, \end{aligned} \quad (2.49)$$

where the commutator term has been absorbed in the error  $\mathcal{T}$ . Integrating by parts with respect to  $s$  and  $k$  in the first two integrals on the right-hand side above respectively, we obtain analogously to the proof of Lemma 2.2:

$$\begin{aligned} &- \int_{\Omega} \mathbb{A}_r^k \bar{\partial}^3 \mathbb{w}_{,s}^r \mathbb{A}_i^s q_{,k} \bar{\partial}^3 v^i + \int_{\Omega} \mathbb{A}_i^k \bar{\partial}^3 \partial_t q_{,k} \bar{\partial}^3 v^i \\ &= \int_{\Gamma} q_{,2} \bar{\partial}^3 \mathbb{w} \cdot \mathbb{A}_{\bullet}^2 \bar{\partial}^3 w \cdot \mathbb{A}_{\bullet}^2 - \int_{\Gamma} \bar{\partial}^3 \partial_t q \bar{\partial}^3 v \cdot \mathbb{A}_{\bullet}^2 + \int_{\Gamma} q_{,1} \mathbb{A}_r^1 \bar{\partial}^3 \mathbb{w}^r \bar{\partial}^3 v \cdot \mathbb{A}_{\bullet}^2 \\ &\quad + \int_{\Omega} (\bar{\partial}^3 q_t + \bar{\partial}^3 \mathbb{w} \cdot v)^2 + \int_{\Omega} \left( \sum_{l=1}^3 d_l \bar{\partial}^{3-l} \mathbb{w} \cdot \bar{\partial}^l v \right. \\ &\quad \left. + \sum_{l=1}^3 e_l \bar{\partial}^l A_i^s \bar{\partial}^{3-l} v^i_{,s} \right) (\bar{\partial}^3 q_t + \bar{\partial}^3 \mathbb{w} \cdot v) + \mathcal{T}. \end{aligned} \quad (2.50)$$

Note further that the first term on the right-hand side above can be, similarly to (2.39), further written as

$$\begin{aligned} \int_{\Gamma} q_{,2} \bar{\partial}^3 \mathbb{w} \cdot \mathbb{A}_{\bullet}^2 \bar{\partial}^3 w \cdot \mathbb{A}_{\bullet}^2 &= \int_{\Gamma} q_{,2} |\bar{\partial}^3 \Lambda_{\kappa} w \cdot \mathbb{A}_{\bullet}^2|^2 \\ &\quad + \int_{\Gamma} \Lambda_{\kappa} \bar{\partial}^3 w \cdot [\Lambda_{\kappa} [q_{,2} (\bar{\partial}^3 w \cdot \mathbb{A}_{\bullet}^2) \mathbb{A}_{\bullet}^2] \\ &\quad - q_{,2} (\bar{\partial}^3 \Lambda_{\kappa} w \cdot \mathbb{A}_{\bullet}^2) \mathbb{A}_{\bullet}^2]. \end{aligned}$$

The second term on the right-hand side of (2.50) reads, using the boundary condition (2.4c)

$$\begin{aligned} \int_{\Gamma} \bar{\partial}^3 \partial_t q \bar{\partial}^3 v \cdot \mathbb{A}_{\bullet}^2 &= \kappa^2 \int_{\Gamma} \bar{\partial}^3 \partial_t h_t (\bar{\partial}^3 (v \cdot \mathbb{A}_{\bullet}^2) - \sum_{l=0}^2 c_l \bar{\partial}^l v \cdot \bar{\partial}^{3-l} \mathbb{A}_{\bullet}^2) \\ &\quad + \kappa^2 \int_{\Gamma} \bar{\partial}^3 \partial_t \beta \bar{\partial}^3 v \cdot \mathbb{A}_{\bullet}^2 \\ &= \frac{\kappa^2}{2} \frac{d}{dt} \int_{\Gamma} J_{\kappa}^{-1} |\bar{\partial}^3 h_t|^2 - \kappa^2 \sum_{l=0}^2 c_l \int_{\Gamma} \bar{\partial}^3 \partial_t h_t \bar{\partial}^l v \cdot \bar{\partial}^{3-l} \mathbb{A}_{\bullet}^2 \\ &\quad + \kappa^2 \int_{\Gamma} \bar{\partial}^3 \partial_t \beta \bar{\partial}^3 v \cdot \mathbb{A}_{\bullet}^2 + \mathcal{T}, \end{aligned}$$

where the error term  $\mathcal{T}$  denotes the lower order terms containing the time derivative of  $J_\kappa$ . We also used the regularized boundary condition (2.4c) in the first equality above. Combining (2.47)–(2.50) and the last identity we obtain the identity (2.22) with error terms  $\mathcal{S}_1$  and  $\mathcal{S}_2$  given by

$$\begin{aligned} \mathcal{S}_1 := & \sum_{\substack{l+\bar{l}=3, k+\bar{k}=1 \\ 0 < l+k < 4}} c_{l,k,\bar{l},\bar{k}} \bar{\partial}^l \partial_t^k \mathbb{A}_i^k \bar{\partial}^{\bar{l}} \partial_t^{\bar{k}} q_{,k} \\ & - \left( \sum_{l=1}^3 d_l \bar{\partial}^{3-l} \mathbb{w} \cdot \bar{\partial}^l v + \sum_{l=1}^3 e_l \bar{\partial}^l \mathbb{A}_i^s \bar{\partial}^{3-l} v^i_{,s} \right) \left( \bar{\partial}^3 q_t + \bar{\partial}^3 \mathbb{w} \cdot v \right); \end{aligned} \quad (2.51)$$

$$\begin{aligned} \mathcal{S}_2 := & -\Lambda_\kappa \bar{\partial}^3 w \cdot [\Lambda_\kappa [q_{,2} (\bar{\partial}^3 w \cdot \mathbb{A}_\bullet^2) \mathbb{A}_\bullet^2] - q_{,2} (\bar{\partial}^3 \Lambda_\kappa w \cdot \mathbb{A}_\bullet^2) \mathbb{A}_\bullet^2] \\ & + \sum_{l=1}^3 c_l (-q_{,2}) \bar{\partial}^3 \mathbb{w} \cdot \mathbb{A}_\bullet^2 (\bar{\partial}^{3-l} (v+w) \cdot \bar{\partial}^l \mathbb{A}_\bullet^2) - q_{,1} \mathbb{A}_r^1 \bar{\partial}^3 \mathbb{w}^r \bar{\partial}^3 v \cdot \mathbb{A}_\bullet^2 \\ & + \kappa^2 \sum_{l=0}^2 c_l \bar{\partial}^3 \partial_t h_t \bar{\partial}^l v \cdot \bar{\partial}^{3-l} \mathbb{A}_\bullet^2 - \kappa^2 \bar{\partial}^3 \partial_t \beta \bar{\partial}^3 v \cdot \mathbb{A}_\bullet^2. \end{aligned} \quad (2.52)$$

Applying the tangential operator  $\bar{\partial} \partial_t^2$  to the equation (2.4b), multiplying by  $\bar{\partial} \partial_t v^i$  and integrating over  $\Omega$  we obtain the identity (2.23) in an analogous way, with error terms  $\mathcal{S}_3$  and  $\mathcal{S}_4$  given by

$$\begin{aligned} \mathcal{S}_3 := & (v_{,s} \cdot \bar{\partial} \Psi_{\kappa tt} \mathbb{A}_i^s - \{\bar{\partial} \partial_t^2, \mathbb{A}_i^k\} q_{,k}) \bar{\partial} v_t^i \\ & + \sum_{1 \leq m+n \leq 2} c_{mn} \bar{\partial}^a \partial_t^b \mathbb{A}_i^k \bar{\partial}^{1-a} \partial_t^{2-b} q_{,k} \bar{\partial} v_{tt}^i \\ & + \left( \sum_{1 \leq m+n \leq 2} d_{mn} \bar{\partial}^a \partial_t^b \mathbb{A}_i^s \bar{\partial}^{1-a} \partial_t^{1-b} v^i_{,s} \right. \\ & \left. - (\Psi_{tt} \cdot \bar{\partial} v + \bar{\partial} \Psi_t \cdot v_t + \Psi_t \bar{\partial} v_t) \right) (\bar{\partial} q_{tt} + \bar{\partial} \Psi_{\kappa,tt} \cdot v); \end{aligned} \quad (2.53)$$

$$\begin{aligned} \mathcal{S}_4 := & q_{,2} \bar{\partial} \mathbb{w}_t \cdot \mathbb{A}_\bullet^2 [(\bar{\partial} (v+w) \cdot \mathbb{A}_\bullet^2) + (\mathbb{w}_t + v_t) \cdot \bar{\partial} \mathbb{A}_\bullet^2 + (\mathbb{w} + v) \cdot \bar{\partial} \mathbb{A}_\bullet^2] \\ & - \bar{\partial} \partial_t \Lambda_\kappa w \cdot \mathbb{A}_\bullet^2 \left[ \Lambda_\kappa ((-q_{,2}) \mathbb{A}_\bullet^2 \bar{\partial} \partial_t w \cdot \mathbb{A}_\bullet^2) - q_{,2} \bar{\partial} \partial_t \Lambda_\kappa w \mathbb{A}_\bullet^2 \right] \\ & - q_{,1} \mathbb{A}_r^1 \bar{\partial} \partial_t \mathbb{w}^r \bar{\partial} \partial_t v \cdot \mathbb{A}_\bullet^2 + \kappa^2 \bar{\partial} \partial_t h_t (\bar{\partial} \partial_t (v \cdot \mathbb{A}_\bullet^2) - \bar{\partial} \partial_t v \cdot \mathbb{A}_\bullet^2) \\ & - \kappa^2 \bar{\partial} \partial_{tt} \beta \bar{\partial} \partial_t v \cdot \mathbb{A}_\bullet^2. \end{aligned} \quad (2.54)$$

### 2.7. Nonlinear Energy Estimates

The following proposition states the desired energy bound for the classical Stefan problem (with  $\sigma = 0$ ), will subsequently lead to a uniform-in- $\kappa$  time of existence for our family solutions to the regularized  $\kappa$ -problems (2.4).

**Proposition 2.6.** (Main energy inequality) *There exists a constant  $C$  independent of  $\kappa$  and a generic polynomial function  $P$  such that for any  $t \in [0, T^\kappa]$  we have the following bound:*

$$\mathcal{E}_\kappa(t) \leq C \mathcal{E}_\kappa(0) + C(t + \sqrt{t}) P(\mathcal{E}_\kappa). \quad (2.55)$$

The proof of the proposition proceeds by systematically estimating error terms in the energy identities from Section 2.4. We shall implicitly use the a priori bound (2.17) freely throughout the proof without explicitly making reference to it.

*Step 1. Estimates for  $\int_0^t \int_\Omega \mathcal{R}_1$  defined by (2.41)* We start by estimating the integral  $\sum_{l=1}^3 \int_0^t \int_\Omega c_l \bar{\partial}^l \mathcal{A}_i^k \bar{\partial}^{4-l} q_{,k} \bar{\partial}^4 v^i$  [the first term appearing in (2.41).] If  $l = 1$ , we have

$$\begin{aligned} \left| \int_0^t \int_\Omega \bar{\partial} \mathcal{A}_i^k \bar{\partial}^3 q_{,k} \bar{\partial}^4 v^i \right| &\leq \|\bar{\partial} \mathcal{A}_i^k\|_{L_t^\infty L_x^\infty} \int_0^t \|\bar{\partial}^3 q_{,k}\|_{L^2} \|\bar{\partial}^4 v^i\|_{L^2} \\ &\lesssim \left\| \int_0^t \bar{\partial} \partial_t (\mathcal{A}_i^k) \right\|_{H^{1.5}} \|\bar{\partial}^3 q_{,k}\|_{L_t^2 L_x^2} \|\bar{\partial}^4 v^i\|_{L_t^2 L_x^2} \\ &\lesssim \sqrt{t} \|\mathfrak{w}\|_{L_t^2 H_x^{3.5}} \|\bar{\partial}^3 q_{,k}\|_{L_t^2 L_x^2} \|\bar{\partial}^4 v^i\|_{L_t^2 L_x^2} \lesssim \sqrt{t} P(\mathcal{E}_\kappa). \end{aligned}$$

For  $l = 2, 3$  we have

$$\begin{aligned} \left| \int_0^t \int_\Omega \bar{\partial}^l \mathcal{A}_i^k \bar{\partial}^{4-l} q_{,k} \bar{\partial}^4 v^i \right| &\lesssim \|\bar{\partial}^l \mathcal{A}_i^k\|_{L_t^\infty H_x^{0.5}} \|\bar{\partial}^{4-l} q_{,k}\|_{H^{0.5}} \int_0^t \|\bar{\partial}^4 v^i\|_{L^2} \\ &\lesssim \sqrt{t} \|\nabla(\Psi_\kappa - \text{Id})\|_{L_t^\infty H_x^{2.5}} \|q\|_{L_t^\infty H_x^{3.5}} \|\bar{\partial}^4 v^i\|_{L_t^2 L_x^2} \\ &\lesssim \sqrt{t} P(\mathcal{E}_\kappa). \end{aligned}$$

We proceed to estimate the integral  $\sum_{l=1}^4 c_l \int_0^t \int_\Omega \bar{\partial}^l \mathcal{A}_i^k \bar{\partial}^{4-l} v^i_{,k} (\bar{\partial}^4 q + \bar{\partial}^4 \Psi_\kappa \cdot v)$  [the second term appearing in (2.41)]. Only cases  $l = 1$  and  $l = 4$  deserve special attention, while the cases  $l = 2$  and  $l = 3$  are estimated by a routine application of the Cauchy–Schwarz inequality and the Sobolev embedding. When  $l = 1$ , we can use Lemma B.2 to conclude that

$$\begin{aligned} &\left| \int_0^t \int_\Omega \bar{\partial} \mathcal{A}_i^k \bar{\partial}^3 v^i_{,k} (\bar{\partial}^4 q + \bar{\partial}^4 \Psi_\kappa \cdot v) \right| \\ &\leq \int_0^t \|\bar{\partial} \mathcal{A}_i^k\|_{0.5} \|v^i_{,k}\|_{2.5} \|\bar{\partial}^4 q + \bar{\partial}^4 \Psi_\kappa \cdot v\|_0 \\ &+ \|\bar{\partial} \mathcal{A}_i^k\|_{L_t^\infty L_x^\infty} \int_0^t \|v^i_{,k}\|_{2.5} (\|\bar{\partial}^4 q\|_{0.5} + \|(\bar{\partial}^4 \Psi_\kappa \cdot v)\|_{0.5}) \\ &\lesssim \left\| \int_0^t \nabla^2 \partial_t (\mathcal{A}_i^k) \right\|_{L_t^\infty L_x^2} \int_0^t \|q\|_{4.5} (\|\bar{\partial}^4 q\|_{0.5} + \|\bar{\partial}^4 \Psi_\kappa \cdot v\|_{0.5}) \\ &+ \left\| \int_0^t \bar{\partial} \partial_t (\mathcal{A}_i^k) \right\|_{H^{1.5}} \int_0^t \|q\|_{4.5}^2 \\ &+ \|\nabla^2 \Psi_\kappa\|_{L_t^\infty H_x^{1.5}} \|v\|_{L_t^\infty H_x^{0.5}} \|\nabla^2 \Psi_\kappa\|_{L_t^\infty H_x^{2.5}} \int_0^t \|q\|_{4.5} \\ &\lesssim \sqrt{t} \|\mathfrak{w}\|_{L_t^2 H_x^3} \|q\|_{L_t^2 H_x^{4.5}}^2 + \sqrt{t} \|\mathfrak{w}\|_{L_t^2 H_x^3} \|\nabla^2 \Psi_\kappa\|_{L_t^\infty H_x^{2.5}} \int_0^t \|q\|_{4.5} \\ &+ C \sqrt{t} \|\mathfrak{w}\|_{L_t^2 H_x^{3.5}} \int_0^t \|q\|_{4.5}^2 \\ &+ C \sqrt{t} \|\nabla^2 \Psi_\kappa\|_{L_t^\infty H_x^{1.5}} \|\nabla^2 \Psi_\kappa\|_{L_t^\infty H_x^{2.5}} \|q\|_{L_t^2 H_x^{4.5}} \lesssim \sqrt{t} P(\mathcal{E}_\kappa). \end{aligned}$$



As for the case  $l = 4$ , we use Lemma B.2 again and obtain

$$\begin{aligned}
 & \left| \int_0^t \int_{\Omega} \bar{\partial}^4 \mathcal{A}_i^k v^{i,k} (\bar{\partial}^4 q + \bar{\partial}^4 \Psi_{\kappa} \cdot v) \right| \\
 & \leq \int_0^t \int_{\Omega} \|\mathcal{A}_i^k\|_{3.5} \|v^{i,k} (\bar{\partial}^4 q + \bar{\partial}^4 \Psi_{\kappa} \cdot v)\|_{0.5} \\
 & \leq \int_0^t \|\nabla^2 \Psi_{\kappa}\|_{2.5} \|v^{i,k}\|_{W^{0.5,\infty}} \|\bar{\partial}^4 q + \bar{\partial}^4 \Psi_{\kappa} \cdot v\|_{0.5} \\
 & \leq \|\nabla^2 \Psi_{\kappa}\|_{L_t^{\infty} H_x^{2.5}} \|q\|_{L_t^{\infty} H_x^{3.5}} \int_0^t (\|q\|_{4.5} + \|\nabla^2 \Psi_{\kappa}\|_{2.5}) \\
 & \leq \sqrt{t} \|\nabla^2 \Psi_{\kappa}\|_{L_t^{\infty} H_x^{2.5}} \|q\|_{L_t^{\infty} H_x^{3.5}} \|q\|_{L_t^2 H_x^{4.5}} + t \|q\|_{L_t^{\infty} H_x^{3.5}} \|\nabla^2 \Psi_{\kappa}\|_{L_t^{\infty} H_x^{2.5}}^2 \\
 & \lesssim (\sqrt{t} + t) P(\mathcal{E}_{\kappa}).
 \end{aligned}$$

The next error term to estimate is  $\sum_{l=1}^4 d_l \int_0^t \int_{\Omega} \bar{\partial}^{4-l} \mathcal{W} \cdot \bar{\partial}^l v (\bar{\partial}^4 q + \bar{\partial}^4 \Psi_{\kappa} \cdot v)$  [the third term appearing in (2.41)]. If  $l = 4$ , we estimate

$$\begin{aligned}
 & \left| \int_0^t \int_{\Omega} \mathcal{W} \cdot \bar{\partial}^4 v (\bar{\partial}^4 q + \bar{\partial}^4 \Psi_{\kappa} \cdot v) \right| \leq \int_0^t \|\mathcal{W}\|_{\infty} \|\bar{\partial}^4 v\|_0 \|\bar{\partial}^4 q + \bar{\partial}^4 \Psi_{\kappa} \cdot v\|_0 \\
 & \leq \sqrt{t} \|\mathcal{W}\|_{L_t^{\infty} H_x^{1.5}} \|\bar{\partial}^4 q + \bar{\partial}^4 \Psi_{\kappa} \cdot v\|_{L_t^{\infty} L_x^2} \|\bar{\partial}^4 v\|_{L_t^2 L_x^2} \lesssim \sqrt{t} P(\mathcal{E}_{\kappa});
 \end{aligned}$$

and analogously for  $l = 3$

$$\begin{aligned}
 & \left| \int_0^t \int_{\Omega} \bar{\partial} \mathcal{W} \cdot \bar{\partial}^3 v (\bar{\partial}^4 q + \bar{\partial}^4 \Psi_{\kappa} \cdot v) \right| \leq \sqrt{t} \|\mathcal{W}\|_{L_t^{\infty} H_x^{2.5}} \|\bar{\partial}^4 q \\
 & \quad + \bar{\partial}^4 \Psi_{\kappa} \cdot v\|_{L_t^{\infty} L_x^2} \|\bar{\partial}^3 v\|_{L_t^2 L_x^2} \lesssim \sqrt{t} P(\mathcal{E}_{\kappa}).
 \end{aligned}$$

For  $l = 1, 2$ , we have

$$\begin{aligned}
 & \left| \int_0^t \int_{\Omega} \bar{\partial}^l \mathcal{W} \cdot \bar{\partial}^{4-l} v (\bar{\partial}^4 q + \bar{\partial}^4 \Psi_{\kappa} \cdot v) \right| \\
 & \leq \int_0^t \|\bar{\partial}^l \mathcal{W}\|_0 \|\bar{\partial}^{4-l} v\|_{\infty} \|\bar{\partial}^4 q + \bar{\partial}^4 \Psi_{\kappa} \cdot v\|_0 \\
 & \leq \sqrt{t} \|\mathcal{W}\|_{L_t^{\infty} H_x^2} \|\bar{\partial}^4 q + \bar{\partial}^4 \Psi_{\kappa} \cdot v\|_{L_t^{\infty} L_x^2} \|q\|_{L_t^2 H_x^{4.5}} \lesssim \sqrt{t} P(\mathcal{E}_{\kappa}).
 \end{aligned}$$

The next-to-last term on the right-hand side of (2.41) is estimated as follows:

$$\begin{aligned}
 & \left| \bar{\partial}^4 \Psi_{\kappa} \cdot v_t (\bar{\partial}^4 q + \bar{\partial}^4 \Psi_{\kappa} \cdot v) \right| \leq \sqrt{t} \|\bar{\partial}^4 \Psi_{\kappa}\|_{L_t^{\infty} L_x^2} \|\bar{\partial}^4 q \\
 & \quad + \bar{\partial}^4 \Psi_{\kappa} \cdot v\|_{L_t^{\infty} L_x^2} \|v_t\|_{L_t^2 H_x^{1.5}} \lesssim \sqrt{t} P(\mathcal{E}_{\kappa}).
 \end{aligned}$$

Finally, to bound  $\int_0^t \int_{\Omega} \bar{\partial}^4 \alpha (\bar{\partial}^4 q + \bar{\partial}^4 \Psi_{\kappa} \cdot v)$  [the last term appearing in (2.41)] we integrate by parts and use the Cauchy–Schwarz inequality to obtain

$$\begin{aligned}
 & \left| \int_0^t \int_{\Omega} \bar{\partial}^4 \alpha (\bar{\partial}^4 q + \bar{\partial}^4 \Psi_{\kappa} \cdot v) \right| \leq \|\bar{\partial}^3 \alpha\|_{L_t^{\infty} L_x^2} \sqrt{t} \|\bar{\partial}^5 q + \bar{\partial} (\bar{\partial}^4 \Psi_{\kappa} \cdot v)\|_{L_t^2 L_x^2} \\
 & \lesssim \mathcal{E}_{\kappa}(0) + t \mathcal{E}_{\kappa}(t).
 \end{aligned}$$

Note that we used Lemma 2.4 and a priori bound (2.17).

Estimates for the error term  $\int_0^t \int_{\Gamma} \mathcal{R}_2$  defined by (2.42). For any  $i, j \in \{1, 2\}$  set  $F = q_{,2} \mathfrak{A}_i^2 \mathfrak{A}_j^2$ ,  $G = \bar{\partial}^3 \Psi_i^j$  and apply Lemma B.1 to conclude

$$\begin{aligned} & \left| \int_0^t \left| \Lambda_{\kappa} [q_{,2} (\bar{\partial}^4 \Psi_t \cdot \mathfrak{A}_{\bullet}^2) \mathfrak{A}_{\bullet}^2] - q_{,2} (\bar{\partial}^4 \Lambda_{\kappa} \Psi_t \cdot \mathfrak{A}_{\bullet}^2) \mathfrak{A}_{\bullet}^2 \right|^2 \right. \\ & \lesssim \int_0^t |q_{,2} \mathfrak{A}_{\bullet}^2 : \mathfrak{A}_{\bullet}^2|_{W^{1,\infty}(\Gamma)}^2 |\Lambda_{\kappa} w|_3^2 \\ & \lesssim \sup_{0 \leq s \leq t} |q_{,2} \mathfrak{A}_{\bullet}^2 : \mathfrak{A}_{\bullet}^2|_2^2 \int_0^t |\Lambda_{\kappa} w|_3^2 \lesssim P(\mathcal{E}_{\kappa}), \end{aligned}$$

where we estimate  $|\Lambda_{\kappa} w|_{L_t^2 H_x^3}$  using (2.12):

$$|h_t|_{L_t^2 H_x^3}^2 \lesssim \int_0^t |\bar{\partial}^3 (\sqrt{1 + |\bar{\partial} h_{\kappa}|^2} (v \cdot \mathfrak{A}_{\bullet}^2))|^2 \lesssim P(\mathcal{E}_{\kappa}) \int_0^t \|q\|_{4.5}^2 \lesssim P(\mathcal{E}_{\kappa}). \quad (2.56)$$

Note that we bounded  $|v|_3$  by relating it to its norm over  $\Omega$  via the trace estimate

$$|v|_{L_t^2 H_x^3}^2 \lesssim \|v\|_{L_t^2 H_x^{3.5}}^2 \lesssim P(\mathcal{E}_{\kappa}) \int_0^t \|q\|_{4.5}^2 \lesssim P(\mathcal{E}_{\kappa}). \quad (2.57)$$

Thus,

$$\begin{aligned} & \left| \int_0^t \int_{\Gamma} \Lambda_{\kappa} \bar{\partial}^4 \Psi \left[ \Lambda_{\kappa} [q_{,2} (\bar{\partial}^4 \Psi_t \cdot \mathfrak{A}_{\bullet}^2) \mathfrak{A}_{\bullet}^2] - q_{,2} (\bar{\partial}^4 \Lambda_{\kappa} \Psi_t \cdot \mathfrak{A}_{\bullet}^2) \mathfrak{A}_{\bullet}^2 \right] \right| \\ & \lesssim P(\mathcal{E}_{\kappa})^{1/2} \left( \int_0^t \int_{\Gamma} |\Lambda_{\kappa} \bar{\partial}^4 \Psi|^2 \right)^{1/2} \lesssim t P(\mathcal{E}_{\kappa}). \end{aligned}$$

Finally, we treat the last term on the right-hand side of (2.42). For  $1 \leq l \leq 2$ , we have

$$\begin{aligned} & \int_0^t \int_{\Gamma} q_{,2} \bar{\partial}^4 \Psi_{\kappa} \cdot \mathfrak{A}_{\bullet}^2 \bar{\partial}^{4-l} (v + \mathfrak{w}) \cdot \bar{\partial}^l \mathfrak{A}_{\bullet}^2 \\ & \leq |q_{,2} \bar{\partial}^l \mathfrak{A}_{\bullet}^2|_{L_t^{\infty} L_x^{\infty}} \int_0^t |\bar{\partial}^4 \Psi_{\kappa} \cdot \mathfrak{A}_{\bullet}^2|_0 |\bar{\partial}^{4-l} (v + \mathfrak{w})|_0 \\ & \lesssim |q_{,2}|_{L_t^{\infty} H_x^1} |\bar{\partial}^l \mathfrak{A}_{\bullet}^2|_{L_t^{\infty} H_x^1} |\bar{\partial}^4 \Psi_{\kappa} \cdot \mathfrak{A}_{\bullet}^2|_{L_t^{\infty} L_x^2} \sqrt{t} (|v|_{L_t^2 H_x^3} + |\mathfrak{w}|_{L_t^2 H_x^3}) \\ & \lesssim \sqrt{t} P(\mathcal{E}_{\kappa}), \end{aligned}$$

where estimates (2.56) and (2.57) were used in the last inequality. If  $l = 3$ , we apply a similar estimate, bounding the term  $\bar{\partial}^l \mathfrak{A}_{\bullet}^2 = \bar{\partial}^3 \mathfrak{A}_{\bullet}^2$  in  $L^2$ -norm and  $\bar{\partial}^{4-l} (v + \mathfrak{w}) = \bar{\partial} (v + \mathfrak{w})$  via  $L^{\infty}$  norm and Sobolev embedding leading to:

$$\left| \int_0^t \int_{\Gamma} q_{,2} \bar{\partial}^4 \Psi_{\kappa} \cdot \mathfrak{A}_{\bullet}^2 \bar{\partial} (v + \mathfrak{w}) \cdot \bar{\partial}^3 \mathfrak{A}_{\bullet}^2 \right| \lesssim \sqrt{t} P(\mathcal{E}_{\kappa}).$$

Case  $l = 4$  is the trickiest error term as four derivatives fall on  $\mathfrak{A}_{\bullet}^2$ , thus creating a term that at highest order contains five derivatives of  $\Psi$ , which is more than the

number of derivatives allowed by our energy  $\mathcal{E}_\kappa$ . However, we have the following identity:

$$\begin{aligned} \int_{\Gamma} q_{,2} \bar{\partial}^4 \Psi_\kappa \cdot \mathcal{A}_\bullet^2 (v + \mathfrak{w}) \cdot \bar{\partial}^4 \mathcal{A}_\bullet^2 &= \frac{1}{2} \int_{\Gamma} \bar{\partial} [q_{,2} \frac{\mathfrak{w} \cdot \tau_\kappa}{|\bar{\partial} \tau_\kappa|}] (\bar{\partial}^4 \Psi_\kappa \cdot \mathcal{A}_\bullet^2)^2 \\ &+ \int_{\Gamma} q_{,2} \frac{\mathfrak{w} \cdot \tau_\kappa}{|\bar{\partial} \tau_\kappa|} \bar{\partial}^4 \Psi_\kappa \cdot \mathcal{A}_\bullet^2 \bar{\partial}^4 \Psi_\kappa \cdot \bar{\partial} \mathcal{A}_\bullet^2 \\ &+ \int_{\Gamma} q_{,2} \bar{\partial}^4 \Psi_\kappa \cdot \mathcal{A}_\bullet^2 E, \end{aligned} \quad (2.58)$$

where  $E$  is the lower order error term given by

$$E = \mathfrak{w} \cdot \tau_\kappa \sum_{l=1}^3 e_l \bar{\partial}^{l+1} \Psi_\kappa \bar{\partial}^{4-l} (|\bar{\partial} \Psi_\kappa|^{-1}) \cdot \mathcal{A}_\bullet^2 + \sum_{l=1}^3 e_l \mathfrak{w} \cdot \tau_\kappa \bar{\partial}^l \tau_\kappa \bar{\partial}^{4-l} \mathcal{A}_\bullet^2. \quad (2.59)$$

To prove (2.58) we first note that

$$v + \mathfrak{w} = (v + \mathfrak{w}) \cdot n_\kappa n_\kappa + (v + \mathfrak{w}) \cdot \tau_\kappa \tau_\kappa = (v + \mathfrak{w}) \cdot \tau_\kappa \tau_\kappa,$$

where we have used the boundary condition (2.4d). Therefore, we have the equality

$$\begin{aligned} (v + \mathfrak{w}) \cdot \bar{\partial}^4 \mathcal{A}_\bullet^2 &= (v + \mathfrak{w}) \cdot \tau_\kappa \tau_\kappa \cdot \bar{\partial}^4 \mathcal{A}_\bullet^2 = \mathfrak{w} \cdot \tau_\kappa \tau_\kappa \cdot \bar{\partial}^4 \mathcal{A}_\bullet^2 \\ &= -\mathfrak{w} \cdot \tau_\kappa \bar{\partial}^4 \tau_\kappa \cdot \mathcal{A}_\bullet^2 + \sum_{l=1}^3 e_l \mathfrak{w} \cdot \tau_\kappa \bar{\partial}^l \tau_\kappa \bar{\partial}^{4-l} \mathcal{A}_\bullet^2, \end{aligned}$$

where we first used the identity  $v \cdot \tau_\kappa = 0$  and in the last line we used the product rule expansion of the identity  $0 = \bar{\partial}^4 (\tau_\kappa \cdot \mathcal{A}_\bullet^2)$  with  $e_l$  the corresponding binomial coefficients. Since  $\tau_\kappa = \frac{\bar{\partial} \Psi_\kappa}{|\bar{\partial} \Psi_\kappa|}$ , we have

$$\begin{aligned} \bar{\partial}^4 \tau_\kappa \cdot \mathcal{A}_\bullet^2 &= \frac{\bar{\partial}^5 \Psi_\kappa}{|\bar{\partial} \Psi_\kappa|} \cdot \mathcal{A}_\bullet^2 + \sum_{l=1}^3 e_l \bar{\partial}^{l+1} \Psi_\kappa \bar{\partial}^{4-l} (|\bar{\partial} \Psi_\kappa|^{-1}) \cdot \mathcal{A}_\bullet^2 \\ &+ \bar{\partial} \Psi_\kappa \bar{\partial}^4 (|\bar{\partial} \Psi_\kappa|^{-1}) \cdot \mathcal{A}_\bullet^2 \\ &= \frac{\bar{\partial}^5 \Psi_\kappa}{|\bar{\partial} \Psi_\kappa|} \cdot \mathcal{A}_\bullet^2 + \sum_{l=1}^3 e_l \bar{\partial}^{l+1} \Psi_\kappa \bar{\partial}^{4-l} (|\bar{\partial} \Psi_\kappa|^{-1}) \cdot \mathcal{A}_\bullet^2, \end{aligned}$$

where we simply used the product rule to expand  $\bar{\partial}^4 (\frac{\bar{\partial} \Psi}{|\bar{\partial} \Psi|})$  and the orthogonality of  $\bar{\partial} \Psi_\kappa$  and  $\mathcal{A}_\bullet^2$  in the last line. Combining the previous two identities, we may write

$$(v + \mathfrak{w}) \cdot \bar{\partial}^4 \mathcal{A}_\bullet^2 = -\frac{\bar{\partial}^5 \Psi_\kappa}{|\bar{\partial} \Psi_\kappa|} \cdot \mathcal{A}_\bullet^2 \mathfrak{w} \cdot \tau_\kappa + E,$$

where the error term  $E$  is given by (2.59). We thus obtain

$$\begin{aligned} \int_{\Gamma} q_{,2} \bar{\partial}^4 \Psi_{\kappa} \cdot \mathbb{A}_{\bullet}^2(v + \mathbb{w}) \cdot \bar{\partial}^4 \mathbb{A}_{\bullet}^2 &= - \int_{\Gamma} q_{,2} \frac{\mathbb{w} \cdot \tau_{\kappa}}{|\bar{\partial} \tau_{\kappa}|} \bar{\partial}^4 \Psi_{\kappa} \cdot \mathbb{A}_{\bullet}^2 \bar{\partial}^5 \Psi_{\kappa} \cdot \mathbb{A}_{\bullet}^2 \\ &\quad + \int_{\Gamma} q_{,2} \bar{\partial}^4 \Psi_{\kappa} \cdot \mathbb{A}_{\bullet}^2 E. \end{aligned}$$

Note that first integral on the right-hand side has a symmetry allowing us to extract a full tangential derivative at the level of highest order terms:

$$\begin{aligned} &- \int_{\Gamma} q_{,2} \frac{\mathbb{w} \cdot \tau_{\kappa}}{|\bar{\partial} \tau_{\kappa}|} \bar{\partial}^4 \Psi_{\kappa} \cdot \mathbb{A}_{\bullet}^2 \bar{\partial}^5 \Psi_{\kappa} \cdot \mathbb{A}_{\bullet}^2 \\ &= - \frac{1}{2} \int_{\Gamma} q_{,2} \frac{\mathbb{w} \cdot \tau_{\kappa}}{|\bar{\partial} \tau_{\kappa}|} \bar{\partial} [(\bar{\partial}^4 \Psi_{\kappa} \cdot \mathbb{A}_{\bullet}^2)^2] + \int_{\Gamma} q_{,2} \frac{\mathbb{w} \cdot \tau_{\kappa}}{|\bar{\partial} \tau_{\kappa}|} \bar{\partial}^4 \Psi_{\kappa} \cdot \mathbb{A}_{\bullet}^2 \bar{\partial}^4 \Psi_{\kappa} \cdot \bar{\partial} \mathbb{A}_{\bullet}^2 \\ &= \frac{1}{2} \int_{\Gamma} \bar{\partial} [q_{,2} \frac{\mathbb{w} \cdot \tau_{\kappa}}{|\bar{\partial} \tau_{\kappa}|}] (\bar{\partial}^4 \Psi_{\kappa} \cdot \mathbb{A}_{\bullet}^2)^2 + \int_{\Gamma} q_{,2} \frac{\mathbb{w} \cdot \tau_{\kappa}}{|\bar{\partial} \tau_{\kappa}|} \bar{\partial}^4 \Psi_{\kappa} \cdot \mathbb{A}_{\bullet}^2 \bar{\partial}^4 \Psi_{\kappa} \cdot \bar{\partial} \mathbb{A}_{\bullet}^2, \end{aligned}$$

where we have used integration by parts in the second equation. Finally, summing the previous two identities we arrive at (2.58).

Note that  $\Psi_{\kappa}$  enters the right-hand side of the above identity at most with 4 derivatives. By standard  $L^{\infty} - L^2 - L^2$  type estimates and identity (2.58), we finally arrive at

$$\left| \int_0^t \int_{\Gamma} q_{,2} \bar{\partial}^4 \Psi_{\kappa} \cdot \mathbb{A}_{\bullet}^2(v + \mathbb{w}) \cdot \bar{\partial}^4 \mathbb{A}_{\bullet}^2 \right| \lesssim \sqrt{t} P(\mathcal{E}_{\kappa}). \quad (2.60)$$

Before we estimate the third term on the right-hand side of (2.42), we first rewrite:

$$\bar{\partial}^4 v \cdot \mathbb{A}_{\bullet}^2 = -\bar{\partial}^4 (\mathbb{w} \cdot \mathbb{A}_{\bullet}^2) - \sum_{l=0}^3 a_l \bar{\partial}^l v \bar{\partial}^{4-l} \mathbb{A}_{\bullet}^2 = -J_{\kappa}^{-1} \bar{\partial}^4 h_{\kappa,t} - \sum_{l=0}^3 a_l \bar{\partial}^l v \bar{\partial}^{4-l} \mathbb{A}_{\bullet}^2,$$

where  $a_l$ ,  $l = 0, \dots, 3$  are the corresponding binomial coefficients. As a consequence, we have

$$\begin{aligned} &\left| \int_0^t \int_{\Gamma} q_{,1} \bar{\partial}^4 \Psi_{\kappa}^r \mathbb{A}_r^1 \bar{\partial}^4 v \cdot \mathbb{A}_{\bullet}^2 \right| \leq \kappa^2 \left| \int_0^t \int_{\Gamma} (v \cdot \mathbb{A}_{\bullet}^2 + \beta)_{,1} \bar{\partial}^4 \Psi_{\kappa}^r \mathbb{A}_r^1 J_{\kappa}^{-1} \bar{\partial}^4 h_t \right| \\ &\quad + \kappa^2 \sum_{l=0}^3 a_l \left| \int_0^t \int_{\Gamma} (v \cdot \mathbb{A}_{\bullet}^2 + \beta)_{,1} \bar{\partial}^4 \Psi_{\kappa}^r \mathbb{A}_r^1 \bar{\partial}^l v \bar{\partial}^{4-l} \mathbb{A}_{\bullet}^2 \right|. \end{aligned} \quad (2.61)$$

The first term on the right-hand side above is easily bounded as follows:

$$\begin{aligned} &\kappa^2 \left| \int_0^t \int_{\Gamma} (v \cdot \mathbb{A}_{\bullet}^2 + \beta)_{,1} \bar{\partial}^4 \Psi_{\kappa}^r \mathbb{A}_r^1 J_{\kappa}^{-1} \bar{\partial}^4 h_t \right| \\ &\leq \sqrt{t} \kappa |v \cdot \mathbb{A}_{\bullet}^2 + \beta)_{,1} \mathbb{A}_r^1 J_{\kappa}^{-1}|_{L_r^{\infty} L_x^{\infty}} |\bar{\partial}^4 \Psi_{\kappa}|_{L_r^{\infty} L_x^2 \kappa} |\bar{\partial}^4 h_t|_{L_t^2 L_x^2} \\ &\lesssim \sqrt{t} \kappa P(\mathcal{E}_{\kappa}). \end{aligned}$$

The second term on the right-hand side of (2.61) is a sum, and the hardest summand to bound is created when  $l = 0$ . In this case, roughly speaking we bound  $|\bar{\partial}^4 \mathbb{A}_\bullet^2|_0$  by  $\kappa^{-1} |\Lambda_\kappa \Psi|_4$  trading one tangential derivative on  $\bar{\partial}^4 \Lambda_\kappa \Lambda_\kappa \nabla \Psi$  for a bound on  $\Lambda_\kappa \nabla \Psi$  in  $H^3$ , at the expense of a factor of  $\kappa^{-1}$ . Using this observation we obtain

$$\begin{aligned} & \kappa^2 \left| \int_0^t \int_\Gamma (v \cdot \mathbb{E}_\bullet^2 + \beta),_1 \bar{\partial}^4 \Psi_\kappa \mathbb{A}_r^1 v \bar{\partial}^4 \mathbb{A}_\bullet^2 \right| \\ & \lesssim \kappa^2 \sqrt{t} |(v \cdot \mathbb{A}_\bullet^2 + \beta),_1 \mathbb{A}_r^1|_{L_t^\infty L_x^\infty} |\kappa^{-1} \bar{\partial}^4 \Lambda_\kappa|_{L_t^\infty L_x^2} \\ & \lesssim \sqrt{t} \kappa P(\mathcal{E}_\kappa). \end{aligned}$$

The next-to-last term on the right-hand side of (2.42) is again a sum and the hardest term to estimate is created again when  $l = 0$ . We use the same idea as in the previous estimate to obtain

$$\left| \kappa^2 \int_0^t \int_\Gamma \bar{\partial}^4 h_t v \cdot \bar{\partial}^4 \mathbb{A}_\bullet^2 \right| \lesssim \sqrt{t} \kappa |\bar{\partial}^4 h_t|_{L_t^2 L_x^2} |v|_{L_t^\infty L_x^\infty} \kappa \kappa^{-1} |\Lambda_\kappa \Psi|_{L_t^\infty L_x^2} \lesssim \sqrt{t} P(\mathcal{E}_\kappa).$$

Note that we exploited the presence of the  $\kappa$ -dependent energy term in our energy  $\mathcal{E}_\kappa$ , using the bound  $\kappa |\bar{\partial}^4 h_t|_{L_t^2 L_x^2} \leq \sqrt{\mathcal{E}_\kappa}$ . In an analogous manner, we conclude

$$\left| \int_\Omega \mathcal{R}_3 + \mathcal{R}_5 dx \right| + \left| \int_\Gamma \mathcal{R}_4 + \mathcal{R}_6 dx' \right| \lesssim (t + \sqrt{t}) P(\mathcal{E}_\kappa),$$

where we note that the commutator term, i.e. the first term on the right-hand side of (2.46) deserves special attention. Due to the absence of spatial derivatives in the term  $\Psi_{ttt}$  in

$$-\Lambda_\kappa \Psi_{tt} \left[ \Lambda_\kappa [q, {}_2 (\Psi_{ttt} \cdot \mathbb{A}_\bullet^2) \mathbb{A}_\bullet^2] - q, {}_2 (\partial_{tt} \Lambda_\kappa \Psi_t \cdot \mathbb{A}_\bullet^2) \mathbb{A}_\bullet^2 \right],$$

we cannot apply the commutator bound from Lemma B.1 in the form stated. Here, we crucially exploit the  $\kappa$ -dependent term in the energy  $\mathcal{E}_\kappa$ . Note that

$$\begin{aligned} & \left| \int_0^t \int_\Gamma \Lambda_\kappa \Psi_{tt} \left[ \Lambda_\kappa [q, {}_2 (\Psi_{ttt} \cdot \mathbb{A}_\bullet^2) \mathbb{A}_\bullet^2] - q, {}_2 (\partial_{tt} \Lambda_\kappa \Psi_t \cdot \mathbb{A}_\bullet^2) \mathbb{A}_\bullet^2 \right] \right| \\ & \leq \sqrt{t} |\Lambda_\kappa \Psi_{tt}|_{L_t^\infty L_x^2} |q, {}_2 \mathbb{A}_\bullet^2 : \mathbb{A}_\bullet^2|_{L_t^\infty W_x^{1,\infty}} \kappa |\Psi_{ttt}|_{L_t^2 L_x^2} \\ & \leq \sqrt{t} P(\mathcal{E}_\kappa), \end{aligned} \tag{2.62}$$

where we gain one power of  $\kappa$  in the second line above from the commutator estimate and then absorb it into the energy contribution  $\kappa |\Psi_{ttt}|_{L_t^2 L_x^2}$ . The last term on the right-hand side of (2.42) contains the  $\beta$ -contribution from the regularized Dirichlet condition (2.4c). It is easily estimated using the Cauchy-Schwarz inequality by a term of the form  $C t m_0 + C t \kappa^2 |\bar{\partial}^4 h_t|_{L^2 L^2}^2$  which in turn is smaller than a constant multiple of  $t m_0 + t \mathcal{E}_\kappa$ . Here  $m_0$  is a constant, which depends only on the initial data.

Estimates for  $\int_{\Omega} \mathcal{S}_1$  and  $\int_{\Gamma} \mathcal{S}_2$ . In the first term on the right-hand side of (2.51), the hardest terms to estimate correspond to the cases  $(\bar{l}, \bar{k}) = (2, 1)$  and  $(l, k) = (2, 1)$ . If  $(\bar{l}, \bar{k}) = (2, 1)$ , then

$$\begin{aligned}
 & \left| \int_0^t \int_{\Omega} \bar{\partial} \mathbb{A}_i^k \bar{\partial}^2 \partial_t q_{,k} \bar{\partial}^3 v^i \right| \\
 & \leq \int_0^t \int_{\Omega} |\bar{\partial}(\bar{\partial} \mathbb{A}_i^k \bar{\partial}^3 v^i) \bar{\partial} \partial_t q_{,k}| \\
 & \leq \int_0^t \|\bar{\partial}^2 \mathbb{A}_i^k\|_{\infty} \|\bar{\partial}^3 v^i\|_0 \|\bar{\partial} \partial_t q_{,k}\|_0 \\
 & \quad + \int_0^t \|\bar{\partial} \partial_t \mathbb{A}_i^k\|_{\infty} \|\bar{\partial}^4 v^i\|_0 \|\bar{\partial} \partial_t q_{,k}\|_0 \\
 & \lesssim \sqrt{t} \|\bar{\partial}^2 \mathbb{A}_i^k\|_{L_t^{\infty} H_x^{1.5}} \|\bar{\partial}^3 v\|_{L_t^{\infty} L_x^2} \|q_t\|_{L_t^2 H_x^2} \\
 & \quad + \left\| \int_0^t \bar{\partial} \mathbb{A}_i^k(s) ds \right\|_{H^{1.5}} \|\bar{\partial}^4 v^i\|_{L_t^2 L_x^2} \|q_t\|_{L_t^2 H_x^2} \\
 & \lesssim \sqrt{t} \|\nabla^2 \Psi_{\kappa}\|_{L_t^{\infty} H_x^{2.5}} \|\bar{\partial}^3 v\|_{L_t^{\infty} L_x^2} \|q_t\|_{L_t^2 H_x^2} \\
 & \quad + t \|\nabla w\|_{L_t^2 H_x^1} \|\bar{\partial}^4 v^i\|_{L_t^2 L_x^2} \|q_t\|_{L_t^2 H_x^2} \\
 & \lesssim (t + \sqrt{t}) P(\mathcal{E}_{\kappa}).
 \end{aligned}$$

Assume now that  $(l, k) = (2, 1)$ , then

$$\left| \int_0^t \int_{\Omega} \bar{\partial}^2 \partial_t \mathbb{A}_i^k \bar{\partial} q_{,k} \bar{\partial}^3 v^i \right| \lesssim \sqrt{t} \|q\|_{L_t^{\infty} H_x^2} \|\bar{\partial}^3 v\|_{L_t^{\infty} L_x^2} \|\mathbb{w}\|_{L_t^2 H_x^3} \lesssim \sqrt{t} P(\mathcal{E}_{\kappa}).$$

The second error term is rather straightforward: for any  $l = 2, 3$ ,

$$\begin{aligned}
 & \left| \int_0^t \int_{\Omega} \bar{\partial}^{3-l} \mathbb{w} \cdot \bar{\partial}^l v (\bar{\partial}^3 q_t + \bar{\partial}^3 \mathbb{w} \cdot v) \right| \\
 & \leq \|\bar{\partial}^{3-l} \mathbb{w}\|_{L_t^{\infty} H_x^{1.5}} \int_0^t \|\bar{\partial}^l v\|_0 \|\bar{\partial}^3 q_t + \bar{\partial}^3 \mathbb{w} \cdot v\|_0 \\
 & \lesssim \sqrt{t} \|\mathbb{w}\|_{L_t^{\infty} H_x^{2.5}} \|\bar{\partial}^l v\|_{L_t^{\infty} L_x^2} \|\bar{\partial}^3 q_t + \bar{\partial}^3 \mathbb{w} \cdot v\|_{L_t^2 L_x^2} \lesssim \sqrt{t} P(\mathcal{E}_{\kappa}).
 \end{aligned}$$

If  $l = 1$ , then

$$\begin{aligned}
 \left| \int_0^t \int_{\Omega} \bar{\partial}^2 \mathbb{w} \cdot \bar{\partial} v (\bar{\partial}^3 q_t + \bar{\partial}^3 \mathbb{w} \cdot v) \right| & \lesssim \|Dv\|_{L_t^{\infty} H_x^{1.5}} \|\mathbb{w}\|_{L_t^{\infty} H_x^2} \|\bar{\partial}^3 q_t \\
 & \quad + \bar{\partial}^3 \mathbb{w} \cdot v\|_{L_t^2 L_x^2} \lesssim \sqrt{t} P(\mathcal{E}_{\kappa}).
 \end{aligned}$$

Similar analysis yields:

$$\sum_{l=1}^3 \left| \int_0^t \int_{\Omega} \bar{\partial}^l \mathbb{A}_i^s \bar{\partial}^{3-l} v^i \cdot s (\bar{\partial}^3 q_t + \bar{\partial}^3 \mathbb{w} \cdot v) \right| \lesssim \sqrt{t} P(\mathcal{E}_{\kappa}).$$

As for the error term (2.52), we start by applying Lemma B.1 to deal with the commutator term. For any  $i, j \in \{1, 2\}$  set  $F = q_{,2} \mathbb{A}_i^2 \mathbb{A}_j^2$ ,  $G = \bar{\partial}^2 w$  and apply Lemma B.1 to obtain

$$\begin{aligned} & \int_0^t \left| \Lambda_\kappa [q_{,2} (\bar{\partial}^3 w \cdot \mathbb{A}_\bullet^2 \mathbb{A}_\bullet^2)] - q_{,2} (\bar{\partial}^3 \Lambda_\kappa w \cdot \mathbb{A}_\bullet^2 \mathbb{A}_\bullet^2) \right|^2 \\ & \lesssim \int_0^t |q_{,2} \mathbb{A}_\bullet^2 : \mathbb{A}_\bullet^2|_{W^{1,\infty}}^2 |w|_2^2 \\ & \lesssim t \sup_{0 \leq s \leq t} |q_{,2} \mathbb{A}_\bullet^2 : \mathbb{A}_\bullet^2|_2^2 \sup_{0 \leq s \leq t} |w|_2^2 \lesssim t P(\mathcal{E}_\kappa), \end{aligned} \quad (2.63)$$

where, in order to bound  $|w|_{L_t^\infty H_x^2}$ , we use the equation (2.4d) analogously to the bound (2.56). Upon using the Cauchy-Schwarz inequality we get

$$\left| \int_0^t \int_\Gamma \Lambda_\kappa \bar{\partial}^3 w \cdot [\Lambda_\kappa [q_{,2} (\bar{\partial}^3 w \cdot \mathbb{A}_\bullet^2 \mathbb{A}_\bullet^2)] - q_{,2} (\bar{\partial}^3 \Lambda_\kappa w \cdot \mathbb{A}_\bullet^2 \mathbb{A}_\bullet^2)] \right| \leq \sqrt{t} P(\mathcal{E}_\kappa).$$

As for the second term on the right-hand side of (2.52), we obtain

$$\begin{aligned} & \left| \int_0^t \int_\Gamma q_{,2} (\bar{\partial}^{3-l}(v+w) \cdot \bar{\partial}^l \mathbb{A}_\bullet^2) \bar{\partial}^3 w \cdot \mathbb{A}_\bullet^2 \right| \\ & \leq \int_0^t |q_{,2}|_\infty (|v|_2 + |w|_2) |\bar{\partial}^3 \mathbb{A}_\bullet^2|_0 |\bar{\partial}^3 \Lambda_\kappa w|_0 \\ & \lesssim \|q\|_{L_t^\infty H_x^{2.5}} (\|v\|_{L_t^\infty H_x^{2.5}} + |w|_2) |\bar{\partial} \Psi - \text{Id}|_3 \int_0^t |\bar{\partial}^3 \Lambda_\kappa w|_0 \lesssim \sqrt{t} P(\mathcal{E}), \end{aligned}$$

where the term  $|w|_{L_t^\infty H_x^2}^2$  is bounded by  $P(\mathcal{E}_\kappa)$  for the same reason as in (2.63). The last term on the right-hand side of (2.52) is a sum, and the hardest term to bound is created when  $l = 0$ . We must integrate by parts with respect to the time variable to obtain

$$\begin{aligned} \kappa^2 \int_0^t \int_\Gamma \bar{\partial}^3 \partial_t h_t v \cdot \bar{\partial}^3 \mathbb{A}_\bullet^2 &= \kappa^2 \int_\Gamma \bar{\partial}^3 h_t v \cdot \bar{\partial}^3 \mathbb{A}_\bullet^2 \Big|_0^t - \kappa^2 \int_0^t \int_\Gamma \bar{\partial}^3 h_t v_t \cdot \bar{\partial}^3 \mathbb{A}_\bullet^2 \\ &\quad - \kappa^2 \int_0^t \int_\Gamma \bar{\partial}^3 h_t v \cdot \bar{\partial}^3 \partial_t \mathbb{A}_\bullet^2. \end{aligned} \quad (2.64)$$

Now observe that

$$\begin{aligned} \kappa^2 \int_\Gamma \bar{\partial}^3 h_t v \cdot \bar{\partial}^3 \mathbb{A}_\bullet^2 \Big|_0^t &\lesssim \kappa^2 m_0 + \kappa |\bar{\partial}^3 h_t|_{L_t^\infty L_x^2} |v|_{L_t^\infty L_x^\infty} \kappa \left| \int_0^t \partial_t \bar{\partial}^3 \mathbb{A}_\bullet^2 \Big|_{L_t^\infty L_x^2} \right| \\ &\lesssim \sqrt{\mathcal{E}_\kappa} \sqrt{\mathcal{E}_\kappa} \sqrt{t} \kappa |\bar{\partial}^4 h_t|_{L_t^2 L_x^2} \lesssim \sqrt{t} P(\mathcal{E}_\kappa), \end{aligned}$$

where  $m_0$  depends only on the initial conditions. As for the remaining three terms on the right-hand side of (2.64), they are straightforward to bound using the standard energy estimates. We arrive at

$$\kappa^2 \left| \int_0^t \int_\Gamma \bar{\partial}^3 \partial_t h_t v \cdot \bar{\partial}^3 \mathbb{A}_\bullet^2 \right| \leq \kappa^2 m_0 + \sqrt{t} (1 + \kappa) P(\mathcal{E}_\kappa).$$

In an analogous manner we conclude

$$\left| \int_0^t \int_\Omega \mathcal{S}_3 dx \right| + \left| \int_0^t \int_\Gamma \mathcal{S}_4 dx' \right| \lesssim (t + \sqrt{t}) P(\mathcal{E}_\kappa).$$

## 2.8. Proof of Theorem 1.1

The polynomial inequality (2.55) replaces the typically used Gronwall inequality. Since the constants appearing in (2.55) are independent of  $\kappa$  a standard continuity argument (see for instance Section 9 of [18]) yields the existence of a  $\kappa$ -independent time  $T$  such that

$$\mathcal{E}_\kappa(t) \leq C\mathcal{E}_\kappa(0) \leq C\mathcal{E}(0) + 1$$

for  $\kappa$  small enough.

Since  $\mathcal{E}(t) \leq \mathcal{E}_\kappa(t)$ ,  $t \in [0, T]$  (recall the definitions (1.20) of  $\mathcal{E}$  and (2.16) of  $\mathcal{E}_\kappa$ ), we obtain the uniform bound

$$\mathcal{E}(q^\kappa, h^\kappa) \leq C\mathcal{E}(0) + 1,$$

where  $(q^\kappa, h^\kappa)_\kappa$  is a family of solutions to the  $\kappa$ -regularized problem (2.4),  $0 \leq \kappa \leq 1$ . Note that the assumptions (2.18) remain valid (on a possibly smaller) time interval  $[0, T]$ , as both  $|\bar{\partial}h|_{L_t^\infty L_x^\infty}$  and  $\delta$  are easily controlled by the energy  $\mathcal{E}$ . By the fundamental theorem of calculus, it is clear that on a possibly smaller time interval  $[0, T]$  we have

$$\sup_{0 \leq t \leq T} \mathcal{A}(t) \leq \mathcal{E}_\kappa(0) + T \sup_{0 \leq t \leq T} \mathcal{E}_\kappa(t) \leq \mathcal{E}_\kappa(0) + \frac{1}{2},$$

thus justifying a posteriori the a priori assumption (2.17). Thus, passing to the weak limit as  $\kappa \rightarrow 0$  we obtain a solution on the time interval  $[0, T]$  which belongs to the space  $\mathcal{S}(T)$  defined in (1.27). Since  $\mathcal{S}(T)$  embeds compactly into  $C_t^1 C_x^0 \cap C_t^0 C_x^2$  the solution is also classical.

**Uniqueness.** We only present a brief sketch of the uniqueness argument. A simple application of the energy method also implies uniqueness of the solution. Assume that  $(\tilde{q}, \tilde{h})$  also solves (2.4) with the corresponding  $\tilde{\Psi}$ ,  $\tilde{v}$ ,  $\tilde{w}$ . Then the pair  $(r, \rho) := (q - \tilde{q}, h - \tilde{h})$  satisfies the following system of equations:

$$r_t - A_i^j (A_i^k r_{,k}),_j = (\Delta_\Psi - \Delta_{\tilde{\Psi}})q - (v - \tilde{v}) \cdot w + \tilde{v}(w - \tilde{w}) \text{ in } \Omega; \quad (2.65a)$$

$$(v - \tilde{v})^i + A_i^k r_{,k} + \tilde{q}_{,k} (A_i^k - \tilde{A}_i^k) = 0 \text{ in } \Omega; \quad (2.65b)$$

$$r = 0 \text{ on } \Gamma; \quad (2.65c)$$

$$\rho_t = -r_{,2} \text{ on } \Gamma; \quad (2.65d)$$

$$\partial_n r = 0 \text{ on } \partial\Omega_{\text{top}}. \quad (2.65e)$$

Furthermore, initially  $(r(0, x), \rho(0, x')) = (0, 0)$ . Applying  $\bar{\partial}$  to the identity (2.65b), multiplying by  $(\bar{\partial}(v - \tilde{v}))^i$  and integrating over  $\Omega$ , we derive the first identity in analogy to the proof of Lemma 2.2. Similarly, applying  $\partial_t$  to (2.65b), multiplying by  $(v - \tilde{v})^i$  and integrating, we obtain the second energy identity. The natural quadratic form that emerges is equivalent to

$$\begin{aligned} E := & \|\bar{\partial}(v - \tilde{v})\|_{L_t^2 L_x^2}^2 + \|v - \tilde{v}\|_{L^\infty L^2}^2 + \|r\|_{L^\infty H_x^1}^2 \\ & + \|r_t\|_{L_t^2 L_x^2}^2 + |\rho|_{L_t^\infty L_x^2}^2 + |\rho_t|_{L_t^2 L_x^2}^2. \end{aligned}$$



Furthermore, we have an a-priori control of the high-order derivatives of the two solutions, i.e. for some  $M > 0$ :  $\mathcal{E}(q, h) + \mathcal{E}(\tilde{q}, \tilde{h}) < M$ . From here, we can easily prove the polynomial bound

$$E(t) \leq tP(E(t)),$$

which in particular, uses the fact that the initial values for  $\rho$  and  $r$  are 0. We infer that  $E = 0$  and hence the uniqueness follows.

**Continuity in Time.** Since  $q \in L_t^2 H_x^5$  and  $q_t \in L_t^2 H_x^3$ , it follows that  $q \in C_t^0 H_x^4$ ; similarly, since  $q_{tt} \in L_t^2 H_x^3$ , then  $q_t \in C_t^0 H_x^2$ . Passing to the limit as  $\kappa \rightarrow 0$  in (2.32),

$$\bar{\partial}^2 h = \frac{g^2 \bar{\partial} v \cdot \tau}{v \cdot n}, \quad (2.66)$$

where  $v \cdot n > 0$  by the Taylor sign condition. By passing to the limit as  $\kappa \rightarrow 0$  in Lemma 2.4, we have that  $\Psi \in L_t^2 H_x^5$ , and we also have that  $\Psi \in L_t^2 H_x^{3.5}$ , from which it follows that  $\Psi \in C_t^0 H_x^4$ . Since  $q \in C_t^0 H_x^4$  and, and since  $v = -\nabla \Psi q \in L_t^2 H_x^4$ , it follows that  $v \in L_t^2 H_x^4 \cap C_t^0 H_x^3$ ; hence,  $\bar{\partial} v \cdot \tau \in L_t^2 H^{2.5}(\Gamma)$ . Then, since  $g$  and  $n$  are in  $L_t^\infty H^3(\Gamma)$ , and  $v \cdot n \in L_t^\infty H^{2.5}(\Gamma)$ , we see from (2.66) that

$$h \in L_t^2 H^{4.5}(\Gamma).$$

Since  $h_t = gv \cdot n$  on  $\Gamma$ , we then have that

$$h_t \in L_t^2 H^{3.5}(\Gamma),$$

from which it follows that

$$h \in C_t^0 H^4(\Gamma).$$

Since  $h_t = gv \cdot n$  on  $\Gamma$ , and since  $g$  and  $n$  are in  $C_t^0 H^3(\Gamma)$  and  $v \in C_t^0 H^{2.5}(\Gamma)$ , then  $h_t \in C_t^0 H^{2.5}(\Gamma)$ . Using that  $h_{tt} = \partial_t [gv \cdot n]$  on  $\Gamma$  and the fact that  $v_t \in C_t^0 H^{0.5}(\Gamma)$ , we also have that  $h_{tt} \in C_t^0 H^{0.5}(\Gamma)$ .

It remains to show that  $q_{tt} \in C_t^0 L_x^2$ . From (1.18a),

$$q_{tt} = (\Delta q)_t - (v \cdot w)_t.$$

Given the regularity already established for  $q$ ,  $q_t$ ,  $\Psi$ , and  $\Psi_t$ , we need to establish the regularity for  $w_t = \Psi_{tt}$ . Since  $h_{tt} \in C_t^0 H^{0.5}(\Gamma)$ , then  $\Psi_{tt} \in C_t^0 H^1(\Omega)$ , and we find that  $q_{tt} \in C_t^0 L^2(\Omega)$ .

### 3. The Vanishing Surface Tension Limit

Local-in-time existence for the Stefan problem with surface tension has been studied in a variety of papers; see, for example, [23, 29, 30, 43]. For any  $(q_0^\sigma, h_0^\sigma) \in H^4(\Omega) \times H^{5.5}(\Omega)$  there exists a local-in-time classical solution  $(q, h)$  to the Stefan problem with surface tension in the harmonic gauge:

$$q_t - \Delta \Psi q = -v \cdot w \quad \text{in } \Omega \times (0, T], \quad (3.1a)$$

$$v^i + A_i^k q_{,k} = 0 \quad \text{in } \Omega \times (0, T], \quad (3.1b)$$

$$q = -\sigma \frac{\bar{\partial}^2 h}{(1 + |\bar{\partial} h|^2)^{\frac{3}{2}}} \quad \text{on } \Gamma \times [0, T], \quad (3.1c)$$

$$\Delta \Psi = 0 \quad \text{on } \Omega \times [0, T], \quad (3.1d)$$

$$\Psi = \text{Id} + h N \quad \text{on } \Gamma \times [0, T], \quad (3.1e)$$

$$\Psi = \text{Id} \quad \text{on } \partial\Omega_{\text{top}} \times [0, T], \quad (3.1f)$$

$$\Psi_t \cdot n(t) = -v \cdot n(t) \quad \text{on } \Gamma \times (0, T], \quad (3.1g)$$

$$v \cdot N = 0 \quad \text{on } \partial\Omega_{\text{top}} \times [0, T], \quad (3.1h)$$

$$\Psi(0, \cdot) = \Psi_0 \quad q(0, \cdot) = q_0^\sigma = p_0 \circ \Psi_0, \quad (3.1i)$$

With  $\sigma > 0$ , we can prove the following energy identities in the same way as Lemma 2.2.

**Lemma 3.1.** *Let  $(q, h)$  be a local-in-time solution to (3.1) defined on the time interval  $[0, T_\sigma]$ . Then we have the following energy identity:*

$$\begin{aligned} F^\sigma(q, \Psi)(t) &= \int_0^t \int_\Omega \{\mathcal{R}_1 + \mathcal{R}_3 + \mathcal{R}_5 + \mathcal{S}_1 + \mathcal{S}_3\} \\ &\quad + \int_0^t \int_\Gamma \{\mathcal{R}_2 + \mathcal{R}_4 + \mathcal{R}_6 + \mathcal{S}_2 + \mathcal{S}_4\} \\ &\quad + \int_0^t \int_\Gamma \{\mathcal{R}_2^\sigma + \mathcal{R}_4^\sigma + \mathcal{R}_6^\sigma + \mathcal{S}_2^\sigma + \mathcal{S}_4^\sigma\}, \end{aligned}$$

where

$$\begin{aligned} F^\sigma &:= \mathcal{F} + \frac{\sigma}{2} \sum_{a+2b \leq 4} \left| |\bar{\partial} \Psi|^{-3/2} J^{-1/2} \bar{\partial}^{a+1} \partial_t^b h \right|_{L_t^\infty L_x^2}^2 \\ &\quad + \sigma \sum_{a+2b \leq 3} \left| |\bar{\partial} \Psi|^{-3/2} J^{-1/2} \bar{\partial}^{a+1} \partial_t^b h_t \right|_{L_t^2 L_x^2}^2, \end{aligned} \quad (3.2)$$

with the energy  $\mathcal{F}$  and error terms  $\mathcal{R}_i, i = 1, \dots, 6, \mathcal{S}_i, i = 1, \dots, 4$  given by (2.24) and Lemma 2.2 respectively, wherein we drop the  $\kappa$ -dependent terms. Furthermore,

$$\begin{aligned} \mathcal{R}_2^\sigma &:= -\sigma \bar{\partial} \left( \frac{\bar{\partial}^2 h}{|\bar{\partial} \Psi|^3} \right) \bar{\partial}^4 \Psi \cdot A \cdot \bar{\partial}^4 v \cdot A^2 \\ &\quad + \sigma \{ -\bar{\partial}^5 h \cdot \bar{\partial} (-\bar{\partial}^4 h_t |\bar{\partial} \Psi|^{-3}) + \bar{\partial}^5 h h_t |\bar{\partial} \Psi|^{-3} \} \\ &\quad + \frac{\sigma}{2} |\bar{\partial}^5 h|^2 \partial_t \left( |\bar{\partial} \Psi|^{-3} J^{-1} \right) + \sigma \bar{\partial}^4 w \cdot A^2 \left[ \bar{\partial}^4 \left( \frac{\bar{\partial}^2 h}{|\bar{\partial} \Psi|^3} \right) - \bar{\partial}^6 h |\bar{\partial} \Psi|^{-3} \right] \\ &\quad + \sigma \sum_{l=1}^4 a_l \bar{\partial}^{4-l} (w + v) \cdot \bar{\partial}^l A^2 \bar{\partial}^4 \left( \frac{\bar{\partial}^2 h}{|\bar{\partial} \Psi|^3} \right) \end{aligned} \quad (3.3)$$

$$\mathcal{R}_4^\sigma := -\sigma \left( \frac{\bar{\partial}^2 h}{|\bar{\partial} \Psi|^3} \right)_{,1} \bar{\partial}^2 \partial_t \Psi \cdot A \cdot \bar{\partial}^2 \partial_t v \cdot A^2$$

$$\begin{aligned}
 & + \frac{\sigma}{2} |\bar{\partial}^3 \partial_t h|^2 \partial_t (|\bar{\partial} \Psi|^{-3} J^{-1}) + \sigma \bar{\partial}^3 \partial_t w \cdot A_{\bullet}^2 \left[ \bar{\partial} \partial_t \left( \frac{\bar{\partial}^2 h}{|\bar{\partial} \Psi|^3} \right) - \frac{\bar{\partial}^3 h_t}{|\bar{\partial} \Psi|^3} \right] \\
 & + \sigma \sum_{l+l' \geq 1} a_{l,l'} \bar{\partial}^2 \partial_t \left( \frac{\bar{\partial}^2 h}{|\bar{\partial} \Psi|^2} \right) \bar{\partial}^l \partial_t' (w+v) \cdot \bar{\partial}^{2-l} \partial_t^{1-l'} A_{\bullet}^2; \tag{3.4}
 \end{aligned}$$

$$\begin{aligned}
 \mathcal{R}_6^{\sigma} & := -\sigma \left( \frac{\bar{\partial}^2 h}{|\bar{\partial} \Psi|^3} \right)_{,1} \Psi_{tt} \cdot A_{\bullet}^1 \partial_{tt} v \cdot A_{\bullet}^2 \\
 & + \frac{\sigma}{2} |\bar{\partial} h_{tt}|^2 \left( |\bar{\partial} \Psi|^{-3} J^{-1} \right)_t - \sigma \bar{\partial} h_{tt} h_{ttt} \bar{\partial} \left( |\bar{\partial} \Psi|^{-3} \right) \\
 & + \sigma w_{tt} \left( \partial_{tt} \left( \frac{\bar{\partial}^2 h}{|\bar{\partial} \Psi|^3} \right) - A_{\bullet}^2 \frac{\bar{\partial}^2 h_{tt}}{|\bar{\partial} \Psi|^3} \right) \\
 & + \sigma \sum_{l=0}^1 a_l \partial_{tt} \frac{\bar{\partial}^2 h}{|\bar{\partial} \Psi|} \partial_t^l (w+v) \cdot \partial_t^{2-l} A_{\bullet}^2; \tag{3.5}
 \end{aligned}$$

$$\begin{aligned}
 \mathcal{S}_2^{\sigma} & := \sigma \sum_{l=1}^3 a_l \bar{\partial}^{3-l} (w+v) \cdot \bar{\partial}^l A_{\bullet}^2 \bar{\partial}^3 \partial_t \bar{\partial} \left( \frac{\bar{\partial} h}{|\bar{\partial} \Psi|} \right) \\
 & + \sigma \sum_{\substack{a+b \leq 4 \\ a \leq 3, b \leq 1}} \bar{\partial}^4 w \cdot A_{\bullet}^2 \bar{\partial}^{a+1} \partial_t^b h \bar{\partial}^{3-a} \partial_t^{1-b} \left( |\bar{\partial} \Psi|^{-1} \right) \tag{3.6}
 \end{aligned}$$

$$\begin{aligned}
 \mathcal{S}_4^{\sigma} & := \sigma \sum_{\substack{l+l' \leq 2 \\ l, l' \leq 1}} a_{l,l'} \bar{\partial}^{1-l} \partial_t^{1-l'} (w+v) \cdot \bar{\partial}^l \partial_t' A_{\bullet}^2 \bar{\partial} \partial_{tt} \bar{\partial} \left( \frac{\bar{\partial} h}{|\bar{\partial} \Psi|} \right) \\
 & + \sigma \sum_{\substack{l+l' \leq 3 \\ l \leq 1, l' \leq 2}} b_{l,l'} \bar{\partial} \partial_t w \cdot A_{\bullet}^2 \bar{\partial}^{l+1} \partial_t^{l'} h \bar{\partial}^{1-l} \partial_t^{2-l'} \left( |\bar{\partial} \Psi|^{-1} \right). \tag{3.7}
 \end{aligned}$$

**Remark 9.** The higher-order energy function  $F^{\sigma}$  is obtained by proceeding in the same way as in the derivation of the energy function  $\mathcal{F}_{\kappa}$  in Sect. 2.6. The essential difference is the nontrivial trace of the term  $\bar{\partial}^4 q$  on the boundary  $\Gamma$ . Since  $q = \sigma \mathcal{H}$  on  $\Gamma$  an integration by parts with respect to  $x_k$  in the integral

$$\int_{\Omega} A_i^k \bar{\partial}^4 q_{,k} \bar{\partial}^4 v^i$$

leads to an additional  $\sigma$ -dependent energy term in (3.2).

### 3.1. Nonlinear Energy Estimates

In the following proposition we prove the basic energy estimate in analogy to Proposition 2.6. Most importantly, we establish a nonlinear polynomial inequality for the energy  $\mathcal{E}_{\sigma}$  with  $\sigma$ -independent coefficients. As a consequence, we show that under the assumptions of Theorem 1.2, the time interval  $T_{\sigma}$  is independent of  $\sigma$ .

**Proposition 3.2.** *Let  $(q_0^{\sigma}, h_0^{\sigma})_{\sigma \geq 0}$  be a given family of well-prepared initial conditions in the sense of Definition 1. There exists a constant  $C$  independent of  $\sigma$  and a*

universal polynomial  $P$  such that for any  $t \in [0, T^\sigma]$  the following bound holds:

$$\mathcal{E}^\sigma(t) \leq C\mathcal{E}^\sigma(0) + C(t + \sqrt{t})P(\mathcal{E}^\sigma). \quad (3.8)$$

In particular, there exists a time  $T > 0$  independent of  $\sigma$ , a constant  $C^* > 0$  and the solution  $(q^\sigma, \Psi^\sigma)$  to the Stefan problem with surface tension defined on  $[0, T]$  satisfying the bound

$$\mathcal{E}^\sigma(q^\sigma, \Psi^\sigma)(t) \leq C^*, \quad 0 \leq \sigma \leq 1, \quad t \in [0, T].$$

**Proof.** In comparison to the estimates for the classical Stefan problem carried over in Section 2.7 the only new error terms to estimate are the terms  $\mathcal{R}_2^\sigma, \mathcal{R}_4^\sigma, \mathcal{R}_6^\sigma, \mathcal{S}_2^\sigma, \mathcal{S}_4^\sigma$  given in the statement of Lemma 3.1.

*Estimating  $\int_0^t \int_\Gamma \mathcal{R}_2^\sigma$  defined by (3.3)* We start by bounding the first term on the right-hand side of (3.3):

$$\begin{aligned} \sigma \left| \int_0^t \int_\Gamma \bar{\partial}^2 \left( \frac{\bar{\partial} h}{|\bar{\partial} \Psi \bar{\partial} h|} \right) \bar{\partial}^4 \Psi \cdot A_\bullet^1 \bar{\partial}^4 v \cdot A_\bullet^2 \right| &\lesssim \int_0^t P(|\sqrt{\sigma} \bar{\partial} h|_4) |\sqrt{\sigma} \Psi|_5 |v|_3 \\ &\lesssim P(|\sqrt{\sigma} \bar{\partial} h|_{L^\infty H^4}) |\sqrt{\sigma} \Psi|_{L^\infty H^5} \sqrt{t} |v|_{L^2 H^3} \lesssim \sqrt{t} P(\mathcal{E}^\sigma \bar{\partial} h). \end{aligned}$$

The second and the third terms on the right-hand side of (3.3) are estimated analogously and rely on the standard  $L^\infty - L^2 - L^2$  estimates. As for the fourth term on the right-hand side of (3.3), note that, due to (2.3),

$$\begin{aligned} \sigma \left| \int_0^t \int_\Gamma J^{-1} \bar{\partial}^4 h_t \left( \bar{\partial}^4 \left( \frac{\bar{\partial}^2 h}{|\bar{\partial} \Psi|^3} \right) - \frac{\bar{\partial}^6 h}{|\bar{\partial} \Psi|^3} \right) \right| \\ \lesssim \int_0^t |\sqrt{\sigma} \bar{\partial}^3 h_t|_0 |\sqrt{\sigma}| \bar{\partial}^4 \left( \frac{\bar{\partial}^2 h}{|\bar{\partial} \Psi|^3} \right) - \frac{\bar{\partial}^6 h}{|\bar{\partial} \Psi|^3} \Big|_1 \\ \lesssim \sqrt{t} P(\mathcal{E}^\sigma), \end{aligned}$$

where the last estimate follows in the standard way; terms with less derivatives are bounded in the  $L^\infty$ -norm and then by the Sobolev embedding theorem. In the last term on the right-hand side of (3.3), the hardest case to deal with is  $l = 4$ . Note that

$$\begin{aligned} \sigma \int_0^t \int_\Gamma (v + w) \cdot \bar{\partial}^4 A_\bullet^2 \bar{\partial}^4 \left( \frac{\bar{\partial}^2 h}{|\bar{\partial} \Psi|} \right) &= \sigma \int_0^t \int_\Gamma (v + w) \cdot \bar{\partial}^4 A_\bullet^2 (\bar{\partial}^6 h |\bar{\partial} \Psi|^{-3}) \\ &+ \sigma \int_0^t \int_\Gamma (v + w) \cdot \bar{\partial}^4 A_\bullet^2 \sum_{l'=1}^4 a_{l'} \bar{\partial}^{6-l'} h \bar{\partial}^{l'} (|\bar{\partial} \Psi|^{-2}) =: I + II. \end{aligned}$$

The more challenging term to estimate is term  $I$ . Since  $A_\bullet^2 = J^{-1}(\bar{\partial} h, -1)$  we have the identity

$$\begin{aligned} I &= \sigma \int_0^t \int_\Gamma (v + w) \cdot \bar{\partial}^4 \left( J^{-1}(\bar{\partial} h, -1) \right) (\bar{\partial}^6 h |\bar{\partial} \Psi|^{-3}) \\ &= \sigma \int_0^t \int_\Gamma J^{-1}(v + w) \cdot (\bar{\partial}^5 h, 0) (\bar{\partial}^6 h |\bar{\partial} \Psi|^{-3}) \end{aligned}$$

$$\begin{aligned}
 & + \sigma \sum_{m=1}^3 c_m \int_0^t \int_{\Gamma} \bar{\partial}^m \left( J^{-1} \right) (v + w) \cdot \bar{\partial}^{4-m} (\bar{\partial} h, -1) (\bar{\partial}^6 h |\bar{\partial} \Psi|^{-3}) \\
 & =: I_A + I_B
 \end{aligned}$$

for some universal constants  $c_m \in \mathbb{R}$ . Note that when  $m = 4$  term  $(v + w) \cdot \bar{\partial}^{4-m} (\bar{\partial} h, -1)$  vanishes since  $(\bar{\partial} h, -1)$  is parallel to  $\mathbf{n}$  and  $v \cdot \mathbf{n} = -w \cdot \mathbf{n}$  by (3.1g).

$$\begin{aligned}
 |I_A| & = \sigma \left| \int_0^t \int_{\Gamma} J^{-1} (v^1 - w^1) \bar{\partial}^5 h \bar{\partial}^6 h |\bar{\partial} \Psi|^{-3} \right| \\
 & = \frac{1}{2} \sigma \left| \int_0^t \int_{\Gamma} J^{-1} \bar{\partial} (|\bar{\partial}^5 h|^2) (v^1 - w^1) |\bar{\partial} \Psi|^{-3} \right| \\
 & = \frac{1}{2} \sigma \left| \int_0^t \int_{\Gamma} |\bar{\partial}^5 h|^2 \bar{\partial} \left( J^{-1} (v^1 - w^1) |\bar{\partial} \Psi|^{-3} \right) \right| \lesssim t P(\mathcal{E}^\sigma),
 \end{aligned}$$

where we have used the parametric representation of  $\Psi$  in terms of  $h$  and integrated by parts. The last inequality is rather standard and follows by estimating  $\bar{\partial} \left( (v^1 - w^1) |\bar{\partial} \Psi|^{-3} \right)$  in  $L^\infty$  norm and further via Sobolev inequality, where we also use  $\sigma |\bar{\partial}^5 h|_{L_t^\infty L_x^2} \lesssim \mathcal{E}^\sigma$ . Terms  $I_B$  and  $II$  are easily estimated via the standard energy  $L^\infty - L^2 - L^2$  bounds and Sobolev imbedding, and the same applies to the remaining cases  $l = 1, 2, 3$ . When estimating the fourth term on the right-hand side of (3.3) first integrate by parts so to remove one  $\bar{\partial}$ -derivative from  $\bar{\partial}^6 \Psi$  term and then apply the standard energy estimates.

*Estimating  $\int_0^t \int_{\Gamma} \mathcal{R}_4^\sigma$  defined by (3.4)* The estimates are completely analogous to the ones for  $\mathcal{R}_2^\sigma$ .

*Estimating  $\int_0^t \int_{\Gamma} \mathcal{R}_6^\sigma$  defined by (3.5)* The first term on the right-hand side of (3.5) is estimated analogously to the first term on the right-hand side of (3.3). Note that

$$\begin{aligned}
 \left| \frac{\sigma}{2} \int_0^t \int_{\Gamma} |\bar{\partial} h_{tt}|^2 (|\bar{\partial} \Psi|^{-3} J^{-1})_t \right| & \lesssim (|\bar{\partial} \Psi|^{-3} J^{-1})_t |_\infty \sigma \int_0^t |\bar{\partial} h_{tt}|_2^2 \\
 & \lesssim t |\bar{\partial} h|_\infty |\bar{\partial} h_{kt}|_\infty \mathcal{E}^\sigma \lesssim t P(\mathcal{E}^\sigma),
 \end{aligned}$$

where we use Sobolev inequality and the definition of  $\mathcal{E}^\sigma$  to infer

$$|\bar{\partial} h_{kt}|_\infty^2 \lesssim |\bar{\partial} h_{kt}|_1^2 \lesssim \int_{\Gamma} (-q, 2) |\bar{\partial}^2 h_{kt}|^2 \lesssim \mathcal{E}^\sigma,$$

and similarly that

$$|\bar{\partial} h|_\infty^2 \lesssim \int_{\Gamma} (-q, 2) |\bar{\partial}^2 h|^2 \lesssim \mathcal{E}^\sigma. \quad (3.9)$$

Space-time integrals of the third and fourth term on the right-hand side of (3.5) are bounded in the usual way by  $P(\mathcal{E}^\sigma)$ . To bound the last term on the right-hand side of (3.5) we distinguish the cases  $l = 0$  and  $l = 1$ . If  $l = 1$ , by the Leibniz rule, expand

$$\partial_{tt} \frac{\bar{\partial}^2 h}{|\bar{\partial} \Psi|} = \bar{\partial}^2 h_{tt} |\bar{\partial} \Psi|^{-1} + 2 \bar{\partial}^2 h_t (|\bar{\partial} \Psi|^{-1})_t + \bar{\partial}^2 h (|\bar{\partial} \Psi|^{-1})_{tt}.$$

For the first two terms above, integrate by parts to move one  $\bar{\partial}$  derivative away from  $\bar{\partial}^2 h_{tt}$  and  $\bar{\partial}^2 h_t$ . Then use the standard  $L^\infty - L^2 - L^2$  type estimates as well as the bound  $|\bar{\partial} v_t|_{L_t^\infty L_x^2} \lesssim \|q_t\|_{L_t^\infty H_x^{2.5}}$  to get the desired estimate. For the third term on right-hand side above we have

$$\begin{aligned} & \sigma \left| \int_0^t \int_\Gamma (\bar{\partial}^2 h (|\bar{\partial} \Psi|^{-1})_{tt}) \partial_t (w + v) \cdot \partial_t A_\bullet^2 \right| \\ & \leq |\bar{\partial}^2 h|_\infty \sqrt{t} |\sqrt{\sigma} (|\bar{\partial} \Psi|^{-1})_{tt}|_{L_t^2 L_x^2} |\partial_t (v + w)|_{L_t^\infty L_x^2} |\partial_t A_\bullet^2|_{L_t^\infty L_x^\infty} \lesssim \sqrt{t} P(\mathcal{E}^\sigma). \end{aligned}$$

*Estimating  $S_2^\sigma$  and  $S_4^\sigma$  defined by (3.6) and (3.7) respectively* The estimates are straightforward and follow the same principle: terms with least amount of derivatives are bounded via Sobolev embedding by the  $\sigma$ -independent energy  $\mathcal{E}(q^\sigma, h^\sigma)$ .

Summing up the above estimates we prove the first inequality in the proposition. The existence of a  $\sigma$ -independent time  $T$  follows from the standard continuity argument and the fact that constant  $C$  in (3.8) is  $\sigma$ -independent. Since  $\mathcal{E}^\sigma(0) \rightarrow \mathcal{E}(0)$  as  $\sigma \rightarrow 0$ , due to our assumption on initial data, the last statement of the proposition follows.  $\square$

*Proof of Theorem 1.2.* Recall the definition (1.38) of  $\|(q, h)\|_{C_t^1 C_x^0 \cap C_t^0 C_x^2}$ . Assume that  $\|(q^\sigma, h^\sigma) - (q^0, h^0)\|_{C_t^1 C_x^0 \cap C_t^0 C_x^2}$  does not converge to 0 as  $\sigma \rightarrow 0$ . Then there exists an  $\epsilon > 0$  and a subsequence  $(\sigma_n)_{n \in \mathbb{N}}$ ,  $\sigma_n \rightarrow 0$  as  $n \rightarrow \infty$ , such that

$$\|(q^{\sigma_n}, h^{\sigma_n}) - (q^0, h^0)\|_{C_t^1 C_x^0 \cap C_t^0 C_x^2} \geq \epsilon \quad \forall n \in \mathbb{N}. \quad (3.10)$$

Since  $\mathcal{E}(q^{\sigma_n}, h^{\sigma_n}) \leq C$ , there exists a subsequence of  $(q^{\sigma_n}, h^{\sigma_n})_n$  (without loss of generality indexed again by  $(\sigma_n)$ ) and  $(\bar{q}, \bar{h}) \in \mathcal{S}$  such that

$$(q^{\sigma_n}, h^{\sigma_n}) \rightharpoonup (\bar{q}, \bar{h}), \quad \text{weakly in } \mathcal{S},$$

where we recall that  $\mathcal{S}$  is defined in (1.27). Note that the injection operator  $I : \mathcal{S} \rightarrow C_t^1 C_x^0 \cap C_t^0 C_x^2$  is compact. Hence  $(q^{\sigma_n}, h^{\sigma_n}) \rightarrow (\bar{q}, \bar{h})$  in  $C_t^1 C_x^0 \cap C_t^0 C_x^2$  where  $(\bar{q}, \bar{h})|_{t=0} = (q_0, h_0)$  due to the property 3) in Definition 1 of the well-prepared initial data. Since  $\sigma_n \rightarrow 0$  as  $n \rightarrow \infty$ ,  $(\bar{q}, \bar{h})$  solves the classical Stefan problem with those initial conditions. From the uniqueness statement of Theorem 1.1, we conclude that  $(\bar{q}, \bar{h}) = (q, h)$ . Thus  $(q^{\sigma_n}, h^{\sigma_n}) \rightarrow (q, h)$  in  $C_t^1 C_x^0 \cap C_t^0 C_x^2$  contradicting (3.10).

#### 4. The Three-Dimensional Case

In this section, we briefly sketch how to adapt the analysis of the previous sections to prove theorems analogous to Theorem 1.1 and 1.2 in the *three* dimensional setting. We assume now that  $\Omega(t)$  is an evolving phase inside the reference domain

$$\Omega := \mathbb{T}^2 \times (0, 1),$$

where  $\mathbb{T}^2$  is the 2-torus. Initially at  $t = 0$  the moving boundary

$$\Gamma_0 = \mathbb{T}^2 \times \{x^3 = h_0(x)\}$$

is parametrized as a graph over  $\Gamma = \mathbb{T}^2 \times \{x^3 = 0\}$  by the height function  $h_0$ . The top boundary  $\partial\Omega_{\text{top}} = \mathbb{T}^2 \times \{x^3 = 1\}$  is fixed and the temperature  $p$  satisfies the homogeneous Neumann boundary condition on  $\partial\Omega$  just like in (1.4). We parametrize boundary as a graph over  $\Gamma$  with the height function  $h(t, x')$ , where  $x' := (x^1, x^2)$ . Using the harmonic coordinates we can change variables as in (1.13) to obtain a fixed boundary problem given by (1.18). The associated energy is given by

$$\begin{aligned}
 \mathcal{E}^{3D}(t) = \mathcal{E}^{3D}(q, h)(t) &:= \sum_{|\alpha|+2b \leq 5} \|\bar{\nabla}^\alpha \partial_t^b v\|_{L_t^2 L_x^2}^2 + \frac{1}{2} \sum_{|\alpha|+2b \leq 4} \|\bar{\nabla}^\alpha \partial_t^b v\|_{L_t^\infty L_x^2}^2 \\
 &+ \frac{1}{2} \sum_{|\alpha|+2b \leq 5} |\sqrt{-q_{,2}} \bar{\nabla}^\alpha \partial_t^b h|_{L_t^\infty L_x^2}^2 + \sum_{|\alpha|+2b \leq 4} |\sqrt{-q_{,2}} \bar{\nabla}^\alpha \partial_t^b h_t|_{L_t^2 L_x^2}^2 \\
 &+ \frac{1}{2} \sum_{|\alpha|+2b \leq 5} \|\bar{\nabla}^\alpha \partial_t^b q + \bar{\nabla}^\alpha \partial_t^b \Psi \cdot v\|_{L_t^\infty L_x^2}^2 \\
 &+ \sum_{|\alpha|+2b \leq 4; } \|\bar{\nabla}^\alpha \partial_t^b q_t + \bar{\nabla}^\alpha \partial_t^b \Psi_t \cdot v\|_{L_t^2 L_x^2}^2. \tag{4.1}
 \end{aligned}$$

In the above definition,  $\alpha = (\alpha_1, \alpha_2)$  is a multi-index of order  $|\alpha| = \alpha_1 + \alpha_2$ , whereby  $\alpha_1, \alpha_2$  are non-negative integers. Symbol  $\bar{\nabla}$  refers to differentiation in tangential directions, i.e.  $\bar{\nabla}^\alpha := \partial_{x^1}^{\alpha_1} \partial_{x^2}^{\alpha_2}$ . The three-dimensional Taylor sign condition for a function  $q$  reads:

$$\min_{x' \in \Gamma} (q_{,3})(t, x', 0) > 0. \tag{4.2}$$

The following theorem holds:

**Theorem 4.1.** *Let the initial conditions  $(q_0, h_0)$  be such that  $\mathcal{E}^{3D}(q_0, h_0) < \infty$  and let  $q_0$  satisfy the Taylor sign condition (4.2). Then the three-dimensional one-phase classical Stefan problem is locally-in-time well-posed, i.e. there is a  $T > 0$  such that there exists a unique solution  $(q, h)$  with the initial data  $(q_0, h_0)$  on the time interval  $[0, T]$ . In addition it satisfies the bound:*

$$\mathcal{E}^{3D}(q, h) \leq 2\mathcal{E}^{3D}(q_0, h_0).$$

Furthermore, let  $(q_0^\sigma, \Psi_0^\sigma)_{\sigma \geq 0}$  be a given family of well-prepared initial conditions in the sense of Definition 1. Assume that it satisfies the Taylor sign condition (4.2) and the corresponding compatibility conditions. By  $(q^\sigma, h^\sigma)_{\sigma \geq 0}$  we denote the associated family of solutions to the problem (1.18). There exists a  $\sigma$ -independent time  $T > 0$  and a constant  $C$  depending only on  $(q_0, h_0)$  such that

$$\mathcal{E}^{3D, \sigma}(q^\sigma, h^\sigma)(T) \leq C \quad \sigma \geq 0$$

for all  $\sigma \geq 0$ . As a consequence, sequence  $(q^\sigma, h^\sigma)$  converges to the unique solution  $(q, h)$  of the classical Stefan problem (1.18) with  $\sigma = 0$  in  $C_t^1 C_x^0 \cap C_t^0 C_x^2$ -norm.

**Remark 10.** Note that the definition of  $\mathcal{E}^{3D}$  contains time derivatives. Thus, to make sense out of the assumption  $\mathcal{E}^{3D}(q_0, h_0) < \infty$ , we express the time derivatives  $\partial_t q_0$  and  $\partial_t h_0$  in terms of the spatial derivatives as explained in Remark 4.

## Appendix A. Modifications of Our Analysis For a More General Initial Domain

In this section we explain how to construct a smooth reference interface for a general graph  $\Gamma_0 = \{\mathbf{x} \mid \mathbf{x} = (x, h_0(x))\} \subset \mathbb{T}^1 \times [0, 1]$ , where the size of  $|h|_{4.5}$  is not necessarily small. For any  $\varepsilon > 0$  we define

$$h_0^\varepsilon(x) = \int_{\mathbb{T}^1} h(y) \rho_\varepsilon(x - y), \quad x \in \mathbb{T}^1, \quad \Gamma_0^\varepsilon = \{\mathbf{x} \mid \mathbf{x} = (x, h_0^\varepsilon(x))\} \subset \mathbb{T}^1 \times [0, 1],$$

and set

$$\Omega_0^\varepsilon := \{(x, y) \in \mathbb{T}^1 \times [0, 1] \mid x \in \mathbb{T}^1, h_0^\varepsilon(x) < y < 1\}.$$

Here  $\rho_\varepsilon$  is the standard mollifier defined in Definition 2 and the domain  $\Omega_0^\varepsilon$  will be our reference domain. Clearly  $h_0^\varepsilon \in C^\infty(\Gamma)$  and for  $\varepsilon$  sufficiently small we can parametrize the evolving surface  $\Gamma(t)$  as a graph over  $\Gamma_0^\varepsilon$  using the outward-pointing unit normal vector field  $N^\varepsilon$  to  $\Gamma_0^\varepsilon$ :

$$\Gamma(t) = \{\mathbf{x} \mid \mathbf{x} = (x, h_0^\varepsilon(x)) + h(t, x)N^\varepsilon(x)\}, \quad N^\varepsilon(x) = \frac{(\bar{\partial}h_0^\varepsilon, -1)}{\sqrt{1 + |\bar{\partial}h_0^\varepsilon|^2}}.$$

Note that  $|h_\varepsilon - h_0|_{4.5} \rightarrow 0$  as  $\varepsilon \rightarrow 0$ . The construction of the harmonic diffeomorphic extension  $\Psi : \Omega_0^\varepsilon \rightarrow \Omega(t)$  of the boundary data

$$\Psi(t, x, h_0^\varepsilon(x)) = (x, h_0^\varepsilon(x)) + h(t, x)N^\varepsilon(x), \quad \Psi(t, x, 1) = (x, 1)$$

is a simple consequence of the existence theory for the Dirichlet boundary value problems for systems of elliptic partial differential equations, since for small  $\varepsilon$  and small times  $t \geq 0$  we have

$$\|\Psi - \text{Id}\|_{4.5} \lesssim \varepsilon \ll 1.$$

Using the argument in (1.7) the trace estimate (1.8) is true. Fixing an  $\varepsilon > 0$  sufficiently small we drop the  $\varepsilon$ -notation and refer to the reference curve  $\Gamma_0^\varepsilon$  as  $\Gamma$ , the reference domain  $\Omega_0^\varepsilon$  as  $\Omega$ , the reference unit normal  $N^\varepsilon$  as  $N$ , and the reference height  $h_0^\varepsilon$  as  $\tilde{h}$ . In the harmonic gauge, the Stefan problem takes nearly the same form (1.18):

$$q_t - A_i^j(A_i^k q_{,k}),_{j} = -v \cdot w \quad \text{in } \Omega, \quad (\text{A.1a})$$

$$v^i + A_i^k q_{,k} = 0 \quad \text{in } \Omega, \quad (\text{A.1b})$$

$$q = 0 \quad \text{on } \Gamma, \quad (\text{A.1c})$$

$$\Psi_t \cdot n(t) = -v \cdot n(t) \quad \text{on } \Gamma, \quad (\text{A.1d})$$

$$v \cdot N = 0 \quad \text{on } \partial\Omega_{\text{top}}, \quad (\text{A.1e})$$

$$q(0, \cdot) = q_0 = p_0 \circ \Psi; \quad \Psi(0, \cdot) = \Psi_0, \quad (\text{A.1f})$$

where

$$\|\Psi_0 - \text{Id}\|_{H^5} \lesssim \varepsilon$$



and the local coordinate realization of the unit normal  $n(t, x)$  takes the more general form:

$$n(t, x) = \frac{(1 - h\mathcal{H})\sqrt{1 + (\bar{\partial}\tilde{h})^2} N - \bar{\partial}hT}{\sqrt{(1 + (\bar{\partial}\tilde{h})^2)(1 - h\mathcal{H}_0)^2 + (\bar{\partial}h)^2}}, \quad x \in \mathbb{T}^1,$$

where

$$\mathcal{H} = -\frac{\bar{\partial}^2\tilde{h}}{(1 + (\bar{\partial}\tilde{h})^2)^{3/2}}, \quad T = \frac{(1, \bar{\partial}\tilde{h})}{\sqrt{1 + (\bar{\partial}\tilde{h})^2}}$$

stand for the mean curvature and the unit tangent to the reference surface  $\Gamma$ , respectively. The proof of Theorem 1.1 applies to (A.1) in an analogous manner; it is simply more technical. The main technical novelty is that the tangential vector-fields to the reference surface  $\Gamma$  are not given by  $\bar{\partial} = \partial_x$ , as  $\Gamma$  may have a nontrivial curvature in general. Therefore, in the neighborhood of  $\Gamma$  for any  $C^1$  function  $f : \Omega \rightarrow \mathbb{R}$  we define the tangential derivative

$$\bar{\partial}f = \nabla f \cdot T,$$

where  $T$  is a local extension of the unit tangent vector field  $T$  into the domain  $\Omega$ . Choosing a smooth cut-off function  $\mu : \Omega \rightarrow [0, 1]$  defined to be 1 in a neighborhood of  $\Gamma$  and 0 in a neighborhood of  $\partial\Omega_{\text{top}}$ , we can replace the operator  $\bar{\partial}$  in Lemma 2.2 by the operator

$$\mu\bar{\partial} + (1 - \mu)\partial_i, \quad i = 1, 2.$$

The ensuing energy identities, energy estimates, and the proof of Theorem 1.1 follow in an analogous way.

## Appendix B. Auxiliary Lemmas

We collect some auxiliary estimates in this section that have been used in the proof of the energy estimates. The following commutator estimate is used in the proof of Proposition 2.6.

**Lemma B.1.** (Lemma 5.1 in [20]) *For  $F \in W^{1,\infty}(\Gamma)$  and  $G, \bar{\partial}G \in L^2(\Gamma)$ , there is a generic constant  $C$  independent of  $\kappa$  such that*

$$|\Lambda_\kappa(F\bar{\partial}G) - f\Lambda_\kappa\bar{\partial}G| \leq C|F|_{W^{1,\infty}(\Gamma)}|G|_0,$$

where  $W^{1,\infty}(\Gamma)$  denotes the Sobolev space of functions  $h \in L^\infty(\Gamma)$  with weak derivative  $\bar{\partial}h \in L^\infty(\Gamma)$ .

Similarly, the following bound is used in estimating some top-order terms in the energy estimates.

**Lemma B.2.** (Lemma 8.5 in [20]) Let  $H^{\frac{1}{2}}(\Omega)'$  denote the dual space of  $H^{\frac{1}{2}}(\Omega)$ . Then there exists a positive constant  $C > 0$  such that

$$\|\bar{\partial}F\|_{H^{\frac{1}{2}}(\Omega)'} \leq C\|F\|_{H^{\frac{1}{2}}(\Omega)}.$$

**Proof.** The proof is a simple consequence of an interpolation estimate between  $L^2(\Omega)$  and  $H^1(\Omega)'$ -spaces. The details are given in [20].  $\square$

*Acknowledgements.* MH was supported by the National Science Foundation under Grant CMG-0530862 and the European Research Council under Grant ERC-240385. SS was supported by the National Science Foundation under Grants DMS-1001850 and DMS-1301380, and by the Royal Society Wolfson Merit Award.

## References

1. ALMGREN, F., WANG, L.: Mathematical existence of crystal growth with Gibbs–Thomson curvature effects. *J. Geom. Anal.* **10**(1), 1–100 (2000)
2. AMBROSE, D.M., MASMOUDI, N.: The zero surface tension limit of three-dimensional water waves. *Indiana U. Math. J.* **58**, 479–522 (2009)
3. AMBROSE, D.M., MASMOUDI, N.: The zero surface tension limit of two-dimensional water waves. *Comm. Pure Appl. Math.* **58**, 1287–1315 (2005)
4. ATHANASOPOULOS, I., CAFFARELLI, L. A., SALSA, S.: Regularity of the free-boundary in parabolic phase-transition problems. *Acta Math.* **176**, 245–282 (1996)
5. ATHANASOPOULOS, I., CAFFARELLI, L. A., SALSA, S.: Phase transition problems of parabolic type: flat free-boundaries are smooth. *Comm. Pure Appl. Math.* **51**, 77–112 (1998)
6. CAFFARELLI, L.A.: Some aspects of the one-phase Stefan problem. *Indiana Univ. Math. J.* **27**, 73–77 (1978)
7. CAFFARELLI, L.A., EVANS, L.C.: Continuity of the temperature in the two-phase Stefan problem. *Arch. Rational Mech. Anal.* **81**, 199–220 (1983)
8. CAFFARELLI, L.A., SALSA, S.: *A geometric approach to free-boundary problems*. American Mathematical Society, Providence, RI, 2005
9. CHENG, C.H.A., COUTAND, D., SHKOLLER, S.: Global existence and decay for solutions of the Hele-Shaw flow with injection. *Interfaces Free Bound.* **16**, 297–338 (2014)
10. CHENG, C.H.A., GRANERO-BELINCHÓN, R., SHKOLLER, S.: Well-posedness of the Muskat problem with  $H^2$  initial data, *Adv. Math.* **286**, 32–104 (2016)
11. CHENG, C.H.A., SHKOLLER, S.: Solvability and regularity for an elliptic system prescribing the curl, divergence, and partial trace of a vector field on Sobolev-class domains. <http://arxiv.org/abs/1408.2469> (2014)
12. CHOI, S., KIM, I.: The two-phase Stefan problem: regularization near Lipschitz initial data by phase dynamics. *Anal. PDE*, **5**(5), 1063–1103 (2012)
13. CHOI, S., KIM, I.: Regularity of one-phase Stefan problem near Lipschitz initial data. *Am. J. Math.*, **132**(6), 1693–1727 (2010)
14. CONSTANTIN, P., CÓRDOBA, D., GANCEDO, F., STRAIN, R. M.: On the global existence for the Muskat problem. *J. Eur. Math. Soc.* **15**(1), 201–227 (2013)
15. CÓRDOBA, A., CÓRDOBA, D., GANCEDO, F.: Interface evolution: the Hele-Shaw and Muskat problems. *Annals of Math.* **173**(1), 477–542 (2011)
16. CÓRDOBA, A., CÓRDOBA, D., GANCEDO, F.: Porous media: the Muskat problem in 3D. *Analysis & PDE*, **6**(2), 447–497 (2013)
17. COUTAND, D., HOLE, J., SHKOLLER, S.: Well-posedness of the free-boundary compressible 3-D Euler equations with surface tension and the zero surface tension limit. *SIAM J. Math. Anal.*, **45**, 3690–3767 (2013)

18. COUTAND, D., SHKOLLER, S.: On the interaction between quasilinear elastodynamics and the Navier-Stokes equations. *Arch. Rational Mech. Anal.* **179**(3), 303–352 (2006)
19. COUTAND, D., SHKOLLER, S.: Well-posedness of the free-surface incompressible Euler equations with or without surface tension. *J. Am. Math. Soc.* **20**(3), 829–930 (2007)
20. COUTAND, D., SHKOLLER, S.: A simple proof of well-posedness for the free surface incompressible Euler equations. *Discr. Cont. Dyn. Systems, Series S* **3**(3), 429–449 (2010)
21. COUTAND, D., SHKOLLER, S.: On the finite-time splash and splat singularities for the 3-D free-surface Euler equations. *Commun. Math. Phys.*, **325**, 143–183 (2014)
22. DE GIORGI, E.:  $\Gamma$ -convergenza e  $G$ -convergenza. *Boll. Un. Mat. Ital.* **5-B**, 213–220 (1977)
23. ESCHER, J., PRÜSS, J., SIMONETT, G.: Analytic solutions for a Stefan problem with Gibbs–Thomson correction. *J. Reine Angew. Math.* **563**, 1–52 (2003)
24. FRIEDMAN, A.: *Variational Principles and free-boundary problems*. Wiley, New York, 1982
25. FRIEDMAN, A.: The Stefan problem for a hyperbolic heat equation. *J. Math. Anal. Appl.* **138**, 249–279 (1989)
26. FRIEDMAN, A., KINDERLEHRER, D.: A one phase Stefan problem. *Indiana Univ. Math. J.* **25** 1005–1035 (1975)
27. FRIEDMAN, A., REITICH, F.: The Stefan problem with small surface tension. *Trans. Amer. Math. Soc.* **328**, 465–515 (1991)
28. FROLOVA, E.V., SOLONNIKOV, V.A.:  $L_p$ -theory for the Stefan problem. *J. Math. Sci.* **99**(1), 989–1006 (2000)
29. HADŽIĆ, M.: Orthogonality conditions and asymptotic stability in the Stefan problem with surface tension. *Arch. Rational Mech. Anal.*, **203** (3), 719–745 (2012)
30. HADŽIĆ, M., GUO, Y.: Stability in the Stefan problem with surface tension (I). *Commun. Partial Diff. Eqns.* **35** (2), 201–244 (2010)
31. HADŽIĆ, M., SHKOLLER, S.: Global stability and decay for the classical Stefan problem. *Comm. Pure Appl. Math.*, **68**, 689–757 (2015)
32. HADŽIĆ, M., SHKOLLER, S.: Global stability and decay for the classical Stefan problem for general boundary shapes. *Philosophical Transactions of Royal Society A*, **373**, pp. 2050, 2015
33. HADŽIĆ, M., NAVARRO, G., SHKOLLER, S.: Local well-posedness and global stability of the two-phase stefan problem (**Preprint**)
34. HANZAWA, E.I.: Classical solution of the Stefan problem. *Tohoku Math. J.* **33**, 297–335 (1981)
35. KAMENOMOSTSKAYA, S.L.: On the Stefan problem. *Mat. Sb.* **53**, 489–514 (1961)
36. KIM, I.: Uniqueness and existence of Hele-Shaw and Stefan problem. *Arch. Rat. Mech. Anal.* **168**, 299–328 (2003)
37. KIM, I., POŽAR, N.: Viscosity solutions for the two-phase Stefan problem. *Comm. PDE* **36**(1), 42–66 (2011)
38. LADYŽENSKAJA, O.A., SOLONNIKOV, V.A., URALŌCEVA, N.N.: Linear and quasilinear equations of parabolic type. *Trans. Math. Monographs* **23**, Am. Math. Soc., Providence, RI (1968), Russian edition: Nauka, Moscow 1967.
39. LUCKHAUS, S.: Solutions for the two-phase Stefan problem with the Gibbs–Thomson law for the melting temperature. *Eur. J. Appl. Math.* **1**, 101–111 (1990)
40. MEIRMANOV, A.M.: *The Stefan Problem*. De Gruyter Expositions in Mathematics. 3 (1992)
41. PRÜSS, J., SAAL, J., SIMONETT, G.: Existence of analytic solutions for the classical Stefan problem. *Math. Ann.* **338**, 703–755 (2007)
42. PRÜSS, J., SIMONETT, G., ZACHER, R.: Qualitative behavior of solutions for thermodynamically consistent Stefan problems with surface tension. *Arch. Ration. Mech. Anal.* **207**, 611–667 (2013)
43. RADKEVICH, E.V.: Gibbs-Thomson law and existence of the classical solution of the modified Stefan problem. *Soviet Dokl. Acad. Sci.* **316**, 1311–1315 (1991)

44. RAYLEIGH, L.: On the instability of jets. *Proc. London Math. Soc.* **1** s1–10, 4–13, 1878
45. RÖGER, M.: Solutions for the Stefan problem with Gibbs–Thomson law by a local minimisation. *Interfaces Free Bound.* **6**, 105–133 (2004)
46. TAYLOR, G.: The instability of liquid surfaces when accelerated in a direction perpendicular to their planes. I *Proc. R. Soc. Lond. A* **201** no. 1065, 192–196 (1950)
47. TAYLOR, M.E.: Partial differential equations. III. Nonlinear equations. Corrected reprint of the 1996 original. *Applied Mathematical Sciences*, **117**, Springer-Verlag, New York, 1997
48. VISINTIN, A.: *Models of phase transitions*. Progr. Nonlin. Diff. Equ. Appl. **28**. Birkhauser, Boston, 1996
49. VISINTIN, A.: *Introduction to Stefan-type problems*. Handbook of differential equations, evolutionary equations, **4**, 377–484, Elsevier B.V., North-Holland, 2008
50. WU, S.: Well-posedness in Sobolev spaces of the full water wave problem in 2-D. *Invent. Math.* **130**(1), 39–72 (1997)

Mahir Hadžić and Steve Shkoller  
Department of Mathematics,  
King's College London,  
London,  
United Kingdom.  
e-mail: mahir.hadzic@kcl.ac.uk

and

Department of Mathematics,  
University of California,  
Davis,  
CA,  
USA.

(Received August 23, 2012 / Accepted August 1, 2016)  
© Springer-Verlag Berlin Heidelberg (2016)