

QUANTUM MECHANICAL ALGORITHMS FOR THE  
NONABELIAN HIDDEN SUBGROUP PROBLEM

MICHELANGELO GRIGNI\*, LEONARD J. SCHULMAN†,  
MONICA VAZIRANI‡, UMESH VAZIRANI§

Received September 14, 2000

We provide positive and negative results concerning the “standard method” of identifying a hidden subgroup of a nonabelian group using a quantum computer.

## 1. Introduction

### 1.1. Overview

The hidden subgroup problem is at present the keystone problem in quantum computation. We are given a function  $f: G \rightarrow S$ , with the property that  $f$  is constant on cosets of an unknown subgroup  $H \subseteq G$ , and distinct on distinct cosets. Here  $f$  is given as an oracle or as an efficient classical program, and  $S$  is an arbitrary set. The problem is to determine the hidden subgroup  $H$ .

The difficulty of the task depends on the type of group  $G$ . The abelian case can be effectively computed with a quantum computer by repetition of coset state preparation and Fourier sampling – the “standard method”

---

*Mathematics Subject Classification (2000):* 81P68, 68Q17

\* Supported in part by NSF grants CCR-9820931 and CCR-0208929.

† Supported in part by NSF CAREER grant 0049092, the NSF Institute for Quantum Information, the Charles Lee Powell Foundation, and the Mathematical Sciences Research Institute.

‡ Supported in part by an NSF Mathematical Sciences Postdoctoral Research Fellowship and NSF grant DMS-0301320.

§ Supported in part by DARPA QUIST grant F30602-01-2-0524, ARO grant DAAD19-03-1-0082, and NSF ITR grant CCR-0121555.

developed by Simon [15] and Shor [14]. In particular this method is the heart of Shor’s solution of the discrete logarithm and factoring problems. In [9] Kitaev formulated the “abelian stabilizer” problem, which he solved by his related “phase estimation” technique. Stabilizer problems are a special case of hidden subgroup problems, but include the key examples.

The status of the nonabelian hidden subgroup problem is one of the most fundamental open problems in quantum algorithms. In particular, the graph automorphism and isomorphism problems may be formulated as hidden subgroup problems over the symmetric group  $S_n$  (see [10]). It is natural to generalize the standard method for the abelian hidden subgroup problem to nonabelian groups. Fourier transforms over nonabelian groups are defined in terms of the irreducible complex representations of the group. There are efficient quantum circuits for computing these transforms for some groups of interest such as the symmetric group (see for example [2, 5, 11]). However, since the dimension of these irreducible representations is in general greater than one, the Fourier transform is not unique, and is defined only up to a unitary change of basis for each irreducible. The Fourier sampling step in the standard method now yields the name of an irreducible representation  $\rho$ , together with the indices  $i, j$  of the entry within that irreducible. The main question, then, is whether the statistics of a sample from the Fourier transform of a coset state reveal sufficient information about the hidden subgroup, to allow for efficient reconstruction. One would hope that this approach is robust, in the sense that the answer to this question should not depend on the arbitrary choice of basis within each irreducible. Our main result is that with respect to a random choice of basis, the Fourier sampling statistics reveal, in general, an exponentially small amount of information about the hidden subgroup. It is still possible that a clever choice of basis within each irreducible can solve the hidden subgroup problem.

Given how algebraically arbitrary this basis choice is, this seems somewhat unlikely. Ideally, one might hope to go beyond the standard method, which is the basis of almost all exponential speedups of quantum algorithms over their classical counterparts. A recent exception to this rule is [16].

Our lower bound on the runtime of the standard method, for subgroups of a group  $G$ , depends upon two parameters: the size of the hidden subgroup  $H$  (naturally the problem becomes easy if  $H$  is very large), and  $c(G)$ , the number of conjugacy classes in  $G$ . We give a lower bound showing that approximately  $\left(\frac{\sqrt{|G|}}{|H|\sqrt{c(G)}}\right)^{1/3}$  rounds of Fourier sampling are required before the standard method can identify  $H$ .

For the special case of hidden subgroups of order 2 in  $S_n$ , this yields a lower bound of approximately  $(k!)^{1/6}$  repetitions of Fourier sampling in order to determine the correct non-identity element of  $H$ , where  $k$  is the number of transpositions in this element. Hallgren, Russell and Ta-Shma [8] independently obtained a similar bound for the weak form of the standard method, where only the name of the irreducible representation  $\rho$  is measured, and the indices  $i, j$  are ignored.

On the positive side, Hallgren, Russell and Ta-Shma [8] showed that the weak form of the standard method for abelian groups efficiently finds hidden normal subgroups in nonabelian groups. We consider a measure of nonabelianness of a group  $G$  – the size of  $\kappa(G)$ , the intersection of all normalizer subgroups. We say that the group is almost abelian if the index of  $\kappa(G)$  in  $G$  is small, and we show that there is a polynomial time algorithm (no longer just Fourier sampling once) for the HSP for any almost abelian group. The new class of groups for which there is an efficient quantum algorithm for the HSP includes the particular example of the semidirect product  $C_3 \rtimes C_m$  for large  $m$  (here  $C_k$  is the cyclic group with  $k$  elements).

Some other interesting previous work on the positive side concerns query complexity. Ettinger, Høyer and Knill [7] show that for any group there exists a sequence of polynomially many queries, from which, with exponentially many measurements, we can reconstruct the hidden subgroup. For the special case of the dihedral group  $D_n$  Ettinger and Høyer [6] showed how to obtain sufficient statistical information about the hidden subgroup using polynomially many queries and polynomially many measurements; leaving open the question of whether there is an efficient reconstruction algorithm using that data. The dihedral group is interesting because by some measures it is not far from abelian, for instance none of its irreps have dimension greater than 2; on the other hand by our measure defined above, it is highly nonabelian, since  $|\kappa(D_n)| \leq 2$ .

## 1.2. The Fourier transform and the standard method for hidden subgroup computation

We first recall some basic group representation theory [13]. Given a group  $G$ , a matrix representation is a group homomorphism  $\rho: G \rightarrow GL(d_\rho, \mathbb{C})$ , where  $GL(d, \mathbb{C})$  is the group of invertible  $d \times d$  complex matrices. A finite group  $G$  has a finite list of inequivalent irreducible representations  $\{\rho\}$ , which we henceforth call its *irreps*. Without loss of generality we may assume the irreps are unitary. The sum of the squares of irrep dimensions  $\sum_\rho d_\rho^2$  equals  $|G|$ , the order of the group.

To every group element  $g$  we associate a complex vector of dimension  $|G|$ , indexed by triples  $\rho, i, j$  where  $\rho$  is an irrep and  $1 \leq i, j \leq d_\rho$  indicate an entry of the matrix  $\rho$ . The vector associated with  $g$  has value  $\frac{\sqrt{d_\rho} \rho_{ij}(g)}{\sqrt{|G|}}$  in the  $\rho, i, j$  entry.

The Fourier transform over  $G$  is the extension of this mapping by linearity to the vector space  $\mathbb{C}^G$  of complex linear combinations of group elements. This linear mapping (whose matrix we will denote  $F$ ) is unitary; this fact is a consequence of the orthogonality relations for group representations.

The trivial representation is the 1-dimensional homomorphism which assigns to every group element the number 1. For a subset  $S$  of  $G$ , define  $|S\rangle = \frac{1}{\sqrt{|S|}} \sum_{g \in S} |g\rangle$  and  $\rho(S) = \rho(|S\rangle) = \frac{1}{\sqrt{|S|}} \sum_{g \in S} \rho(g)$ . The orthogonality relations imply that  $\rho(G)$  is  $\sqrt{|G|}$  when  $\rho$  is the trivial representation, and a zero matrix otherwise. (As mentioned above, the Fourier transform has a scalar factor  $\sqrt{d_\rho/|G|}$ , so this corresponds to the fact that the Fourier transform of the unit norm uniform superposition on  $G$ , is 1 on the trivial representation and 0 elsewhere.)

$\mathbb{C}^G$  has an additional structure beyond its vector space structure: it is also an algebra over  $\mathbb{C}$ , using the product which is the extension of the group product by linearity. This structure is preserved by the Fourier transform, simply because each irrep is a group homomorphism. This is what is often known, for abelian groups (where each irrep is 1-dimensional), as the “convolution-multiplication” property of the Fourier transform.

In the “standard method” for the hidden subgroup problem we begin by forming the uniform superposition over a random coset  $gH$  of the hidden subgroup  $H$ : in other words, we form<sup>1</sup> the uniform distribution over vectors  $|gH\rangle$ . First suppose that we know  $g$  (or at least  $gH$ ), then we have the pure superposition  $|gH\rangle$ . We then apply the Fourier transform to this superposition, obtaining the vector

$$\frac{1}{\sqrt{|G||H|}} \sum_{\rho, i, j} \sqrt{d_\rho} \sum_{h \in H} \rho_{ij}(gh) |\rho, i, j\rangle.$$

---

<sup>1</sup> To form this mixture of superpositions, we first form the uniform-amplitudes superposition  $\frac{1}{\sqrt{|G|}} \sum_{g \in G} |g, 0\rangle$ , and then compute  $f$ , obtaining the superposition  $\frac{1}{\sqrt{|G|}} \sum_{g \in G} |g, f(g)\rangle$ . We then measure  $f(g)$ , which determines the coset  $gH$ . The result is the superposition  $\frac{1}{\sqrt{|H|}} \sum_{h \in H} |gh\rangle$  for a uniformly random  $g$ .

By not using  $f(g)$  to affect the subsequent computation, we are discarding some potentially useful information. No proposal exists, however, for taking advantage of this information.

This gives rise to the probability distribution

$$P_{gH}(|\rho, i, j\rangle) = \frac{d_\rho}{|G||H|} \left| \sum_{h \in H} \rho_{ij}(gh) \right|^2 = \frac{d_\rho}{|G|} |\rho(gH)_{ij}|^2.$$

Since we actually do not know  $g$ , and  $g$  is distributed uniformly, we sample  $\rho, i, j$  with the probability

$$P_H(|\rho, i, j\rangle) = \frac{1}{|G|} \sum_{g \in G} P_{gH}(|\rho, i, j\rangle).$$

The success of this method depends on how much statistical information about  $H$  is present in this distribution. In particular: do a polynomial number of samples suffice to identify  $H$  with high probability? In the following  $\chi_\rho(g)$  denotes the *character* of  $\rho$  at  $g$ , which is simply the trace of  $\rho(g)$ .

**Lemma 1.**  $\rho(H) = \frac{1}{\sqrt{|H|}} \sum_{h \in H} \rho(h)$  is  $\sqrt{|H|}$  times a projection matrix, and  $\text{rank}(\rho(H)) = \frac{1}{|H|} \sum_{h \in H} \chi_\rho(h)$ .

**Proof.** Restricted to  $H$ ,  $\rho$  decomposes into the direct sum of several irreps  $\sigma_1, \dots, \sigma_k$ .  $\rho(H)$  is the direct sum of  $\sigma_i(H)$ ; as discussed above  $\sigma_i(H)$  is  $\sqrt{|H|}$  if  $\sigma_i$  is the trivial representation of  $H$ , and zero otherwise. ■

A certain amount of information about  $H$  is given just by sampling  $\rho$ , and ignoring the matrix indices  $i$  and  $j$ . We refer to this as the “weak form” of the standard method. In the normal case this more limited information is already enough, and in fact no further information is available in the indices. For general subgroups further information is present in the indices, and in the “strong form” of the method, these are sampled as well; we will discuss this issue below. First we show that, when we Fourier sample the unit norm uniform superposition on  $gH$ , (i.e. sample from the probability distribution defined by the Fourier transform of this superposition), the probability  $P_{gH}(|\rho\rangle)$  of sampling  $\rho$  is independent of  $g$ .

**Lemma 2.** <sup>2</sup>The probability of measuring  $\rho$  is the same for the uniform superposition on the coset  $gH$  (or  $Hg$ ), as for the superposition on  $H$ .

**Proof.**  $\rho(gH) = \rho(g)\rho(H)$  and  $\rho(g)$  is unitary. ■

**Corollary 3.**  $P_{gH}(|\rho\rangle) = P_H(|\rho\rangle) = \frac{d_\rho}{|G|} \sum_{h \in H} \chi_\rho(h) = \frac{|H|d_\rho}{|G|} \text{rank}(\rho(H))$ .

**Corollary 4.** The probability of sampling  $\rho$  is the same for the subgroup  $H$  as it is for a conjugate subgroup  $g^{-1}Hg$ .

<sup>2</sup> For methods that measure  $\rho$  but discard  $i$  and  $j$ , Lemma 2 implies that there is no loss in discarding  $f(g)$  as well. In particular we may discard  $f(g)$  when  $G$  is commutative.

## 2. Normal $H$

Hallgren, Russell, and Ta-Shma [8] showed that the weak form of the standard method quickly obtains enough information to identify hidden normal subgroups. This section briefly describes how.

Recall that in the standard method we sample from the Fourier transform of the uniform superposition over a random coset  $gH$  of the hidden subgroup  $H$ . In the weak form of this method, we just sample the name of the irreducible  $\rho$  that results from this transform, and let  $N = \bigcap_{\rho} \ker(\rho)$  for a sequence of  $O(\log |G|)$  such samples.

**Theorem 5.** [8] *The intersection of  $\ker(\rho)$  from  $O(\log |G|)$  repetitions of Fourier sampling is with high probability equal to the largest normal subgroup of the hidden subgroup.*

We first show that when we restrict attention to normal subgroups, all the information about  $H$  is present in the label of the sampled irrep. By lemma 2, the probability of sampling  $\rho$  is independent of the particular coset  $gH$ : so we will examine the uniform superposition on  $H$ .

**Lemma 6.** *If  $H$  is a normal subgroup of  $G$  and  $\rho$  is an irrep of  $G$ ,  $\rho(H)$  is a nonnegative scalar multiple of the identity  $I$ , nonzero if and only if  $H \subseteq \ker(\rho)$ .*

**Proof.** Let  $\sigma_1, \dots, \sigma_k$  be the decomposition of  $\rho$  for  $H$ . We claim that if  $\sigma_1$  is trivial, so are all the rest.

Let  $W$  be the space  $\rho$  acts on. Let  $V$  be the 1-dimensional subspace of  $W$  which  $\sigma_1$  acts on. Since  $\rho$  is irreducible over  $G$ , the elements  $g$  of  $G$  carry  $V$  to a set of subspaces spanning  $W$ . Since  $H = gHg^{-1}$  for every  $g$ , each of the images  $gV$  is invariant for  $H$ . ■

To prove theorem 5, it suffices to show that if  $N$  is the current intersection of the kernels, but  $N \not\subseteq H$ , then with probability at most  $1/2$ , the next Fourier sampling will yield an irrep  $\rho$  such that  $N \subseteq \ker(\rho)$ . This probability is given by:

$$\sum_{\rho: N \subseteq \ker(\rho)} \frac{d_{\rho} |H| \text{rank}(\rho(H))}{|G|}$$

Observe that since  $N$  is normal in  $G$ , Fourier sampling the superposition  $|NH\rangle$  (where  $NH$  is the set of ordered products of elements of  $N$  and  $H$ ) yields only irreps whose kernels contain  $N$ . Again since  $N$  is normal in  $G$ ,

$NH$  is a group, so we can write

$$1 = \sum_{\rho: N \subseteq \ker(\rho)} \frac{d_\rho |NH| \text{rank}(\rho(NH))}{|G|}.$$

Next observe that  $\rho(NH) = \rho(N)\rho(H)$ , which is a nonzero scalar multiple of  $\rho(H)$  for any  $\rho$  whose kernel contains  $N$ . Hence when we Fourier sample from the superposition  $|H\rangle$ , the probability of obtaining an irrep  $\rho$  whose kernel contains  $N$  is

$$\begin{aligned} \sum_{\rho: N \subseteq \ker(\rho)} \frac{d_\rho |H| \text{rank}(\rho(H))}{|G|} &= \sum_{\rho: N \subseteq \ker(\rho)} \frac{d_\rho |H| \text{rank}(\rho(NH))}{|G|} \\ &\leq \frac{1}{2} \sum_{\rho: N \subseteq \ker(\rho)} \frac{d_\rho |NH| \text{rank}(\rho(NH))}{|G|} \\ &= \frac{1}{2}. \end{aligned} \quad \blacksquare$$

### 3. “Almost Abelian” Groups

#### 3.1. Algorithm

The case of normal subgroups was one way of extending the standard method beyond the abelian case. Another extension is to consider the case in which  $G$  is nearly abelian in some measure. We consider the *Baer norm* [1] of  $G$ , which is the intersection of the normalizers of all subgroups of  $G$ . Following Bryce [3], we denote this intersection by  $\kappa(G)$ . (Thus  $\kappa(G) = \bigcap_H N(H)$  where the intersection ranges over subgroups  $H$  of  $G$ .) For abelian groups, of course,  $\kappa(G) = G$ . In order for our algorithm to run in polynomial time (in  $n = \log |G|$ ), the index  $[G : \kappa(G)]$  should be  $\exp(O(\log^{1/2} n))$ .

The basic method to identify the unknown  $H$  for “almost abelian”  $G$  begins again with the observation that  $\kappa(G) \subseteq N(H) \subseteq G$ .

**Lemma 7.**  $\kappa(G)$  is normal in  $G$ .

**Proof.** Let  $g \in \kappa(G)$  and  $a \in G$ . We wish to show that  $aga^{-1} \in N(H) \forall H$ . Fix  $H$ , then for any  $h \in H$ ,

$$(aga^{-1})h(aga^{-1})^{-1} = aga^{-1}hag^{-1}a^{-1}.$$

Now  $a^{-1}ha \in a^{-1}Ha$ , and since  $g \in \kappa(G) \subseteq N(a^{-1}Ha)$ , it follows that  $ga^{-1}hag^{-1} \in a^{-1}Ha$ . But then, as desired,

$$aga^{-1}hag^{-1}a^{-1} \in H. \quad \blacksquare$$

The algorithm to determine  $H$  is this: for each subgroup  $J$  of  $G$  containing  $\kappa(G)$ , run the normal-subgroups algorithm on  $J$  and determine (with high probability) its hidden subgroup  $H_J$ . Then output the union of the  $H_J$ .

**Theorem 8.** *With high probability  $H = \bigcup_J H_J$ . If  $[G : \kappa(G)] \in \exp(O(\log^{1/2} n))$ , the algorithm runs in polynomial time.*

**Proof.** Although  $N(H)$  is unknown,  $N(H)/\kappa(G)$  is a subgroup of  $G/\kappa(G)$ , and the algorithm examines all possibilities. A group of order  $a$  has at most  $2^{lg^2 a}$  subgroups. Thus a bound of  $\exp(O(\log^{1/2} n))$  on  $[G : \kappa(G)]$  guarantees that we only need consider polynomially many subgroups.

All of the hidden subgroups of the various  $J$  are subgroups of  $H$ ; at least one of them is equal to  $H$ . The guarantees for the normal hidden subgroup algorithm ensure that there is only a small probability that any  $H_J$  produced by the algorithm differs from the hidden subgroup of  $J$ . ▀

### 3.2. Example: extensions of groups

One way to construct an almost abelian group is by extending one abelian group  $A$  by another  $B$ . We say  $G$  is an extension of  $A$  by  $B$  if  $A$  is normal in  $G$  and  $G/A \simeq B$ .

Here we'll consider the special case when  $G$  is the semidirect product of  $A$  by  $B$ , written  $G = A \rtimes B$ . In other words,  $A$  is a normal subgroup of  $G$ ,  $B$  is isomorphic to a subgroup of  $G$ ,  $AB = G$  and  $A \cap B = \{1\}$ . The representation theory of  $G = A \rtimes B$  is well understood in terms of that of  $A$  and  $B$ .

To define the semidirect product, we need a homomorphism  $\theta : B \rightarrow \text{Aut}(A)$ . Then the group structure of  $G = AB$  is defined by the identity

$$bab^{-1} = (\theta(b))(a).$$

(Since construction of  $G$  from  $A$  and  $B$  requires specification of  $\theta$ , one can more carefully write  $G = A \rtimes_{\theta} B$ . This is unnecessary when  $A$  and  $B$  are specified as particular subgroups of a given  $G$ .)

We remark that  $\theta$  need not be injective or surjective. In fact, it will be convenient for us to have  $\ker(\theta)$  be large, because  $\ker(\theta) \subseteq Z(G) \subseteq \kappa(G)$  so this provides us with a large  $\kappa(G)$ . (Here  $Z(G)$  denotes the center of  $G$ .)



A basic example of a semidirect product is the dihedral group  $D_n = \langle x, y \mid x^2 = y^n = 1; xyx^{-1} = y^{-1} \rangle = C_n \rtimes C_2$  (where  $C_n$  denotes a cyclic group of order  $n$ ). The homomorphism  $\theta$  sends the nontrivial element  $x \in C_2$  to the map  $y \mapsto y^{-1}$ , because  $xyx^{-1} = y^{-1}$ .

In the context of “almost abelian” groups we are interested in the following example:  $G = C_3 \rtimes C_m$  where  $m$  is a power of two. Let  $a$  be a generator for the  $C_3$  subgroup, and  $b$  a generator for the  $C_m$  subgroup. We have  $bab^{-1} = a^2$ .

Let  $\omega$  be a primitive 3’rd root of unity and  $\zeta$  a primitive  $m$ ’th root of unity.

$G$  has  $m$  one-dimensional representations. These correspond to the trivial character of  $C_3$ . Each is indexed by  $0 \leq k < m$ , and sends  $(a^i, b^j)$  to  $\zeta^{kj}$ .

$G$  has  $m/2$  two-dimensional representations. These correspond to the character  $\chi$  of  $C_3$  for which  $\chi(a) = \omega$ . The  $m/2$  representations, indexed by  $0 \leq k < m/2$ , are

$$\begin{aligned} \Phi_k(a) &= \begin{pmatrix} \omega & 0 \\ 0 & \omega^2 \end{pmatrix} \\ \Phi_k(b) &= \begin{pmatrix} 0 & \zeta^k \\ \zeta^k & 0 \end{pmatrix}. \end{aligned}$$

It is easy to modify the standard quantum Fourier transform circuits to compute the Fourier transform for  $G$ . Given a group element  $(a^i, b^j)$ , represented as the pair  $|i, j\rangle$ , begin with a Fourier transform over the group  $C_3$  on the first index,  $i$ . Conditional on the new value of the first index,  $i'$ , being 0, perform a quantum Fourier transform over the group  $C_m$  on  $j$ : this yields the superposition on the one-dimensional irreps. Conditional on  $i'$  being 1 or 2, separate  $j$  into its low-order bit  $j_0$  and the high order bits  $j_h$ . Perform a quantum Fourier transform over the group  $C_{m/2}$  on  $j_h$ . The result of the last transform indexes the irrep, while  $i'$  and  $j_0$  index the entry within the irrep.

### 4. General $H$

Up to now we have focussed on the extent to which information about  $H$  can be detected from measuring just the name of the irrep,  $\rho$ . (The “weak” standard method.) Of course we can actually measure more, namely the row index  $i$  and column index  $j$  within  $\rho$ . (The “strong” standard method.) It is possible that this contributes substantially to our power. In particular, conjugate subgroups give rise to identical distributions on irreps and so cannot be told apart without measuring the matrix indices within the irreps. In

this section, we establish limits on what further information can be obtained from the row and column labels.

#### 4.1. Row indices provide no information

In this section we show that – due to the fact that in the standard method we average over random cosets  $gH$  – there is no point in measuring the row index  $i$ . (Whether row or column depends on whether the group acts on the left or right. Here we suppose the group acts on the left.) This is because, conditional on measuring  $\rho$  and  $j$ , the distribution on  $i$  is independent of  $H$  (actually it's always uniform). Even more, given that we measure the column index  $j$ , there is no measurement within the row space that will reveal any further information about the hidden subgroup. We describe an arbitrary measurement in the row space by an arbitrary unitary action  $U$ , followed by measurement of the row index  $i$ .

We now show, for any  $U$ , the uniformity of the distribution on  $i$ , conditional on  $j$ . For a particular coset  $gH$ , the probability of sampling entry  $i, j$  is proportional to the norm squared of  $(U\rho(gH))_{ij}$ . Thus the probability of sampling entry  $i, j$  is proportional to the norm squared of the  $|G|$ -dimensional vector  $((U\rho(gH))_{ij})_{g \in G}$  with entries indexed by  $g$ . Since  $\rho(gH) = \rho(g)\rho(H)$ , this vector is a linear combination of the  $d_\rho$  vectors  $((U\rho(g))_{ik})_{g \in G}$  (each  $|G|$ -dimensional), with coefficients  $\rho(H)_{kj}$ . By the orthogonality relations, the  $d_\rho^2$  vectors  $(\rho(g)_{\ell k})_{g \in G}$  are orthonormal; by unitarity of  $U$ , so are the vectors  $((U\rho(g))_{\ell k})_{g \in G}$ . Therefore the norm squared of the  $|G|$ -dimensional vector  $((U\rho(gH))_{ij})_{g \in G}$  is equal to the norm squared of the  $j$ -th column of  $\rho(H)$ , and independent of  $i$ .

If we keep track of the leading constants, this argument shows that (even after applying any unitary transformation to the row space),

**Theorem 9.**  $P_H(|\rho, i, j\rangle) = \frac{1}{|G|} |\rho(H)_j|_2^2$ . ■

The fact that no operation on the row space can change the measurement statistics implies that the underlying density matrix is a tensor product of the identity density matrix for the row space, with the density matrix describing the column space. To be specific, let  $\Delta_\rho$  be the density matrix in the Fourier basis, restricted to the irrep  $\rho$ . The entries of  $\Delta_\rho$  are labelled by two indices  $i_1, i_2$  (for rows of  $\rho$ ) and two indices  $j_1, j_2$  (for columns of  $\rho$ ). Let  $I$  be the  $d_\rho \times d_\rho$  identity matrix with rows indexed by  $i_1$  and columns indexed by  $i_2$ . Let  $\sigma$  be the  $d_\rho \times d_\rho$  matrix with rows indexed by  $j_1$  and columns indexed by  $j_2$ , defined by  $\sigma_{j_1, j_2} = \sum_k \rho(H)_{k, j_1} \rho(H)_{k, j_2}^*$ . Then

**Theorem 10.**  $\Delta_\rho = \frac{1}{|G|} I \otimes \sigma$ . ■

### 4.2. Random basis

The Fourier transform is uniquely defined only up to a change of basis within each irrep; for abelian groups all irreps are one-dimensional so there is no ambiguity in the definition of the transform, but for nonabelian groups there is an arbitrary choice of basis to be made within each irrep. The amount of additional statistical information available from measuring the matrix indices  $i, j$ , in addition to the irrep  $\rho$ , may in general depend on the basis. In this section, we show that if we choose a random basis for each irrep, then the additional information available is negligible, provided that the subgroup  $H$  is sufficiently small and the group  $G$  is sufficiently nonabelian.

Given an irrep  $\rho$  in a particular basis, the probability of sampling the  $j$ -th column of  $\rho$  is  $\frac{d_\rho}{|G|} |\rho(H)_j|_2^2$  (where  $\rho(H)_j$  is the  $j$ -th column of  $\rho(H)$ ). Thanks to the previous section, the row index  $i$  is uniformly random and therefore can be ignored. Suppose we now choose a different basis for  $\rho$ , which we do by replacing  $\rho$  by the isomorphic irrep  $A^{-1}\rho A$  for a unitary  $A$ . Then the probability of measuring the  $j$ -th column in this modified irrep is  $\frac{d_\rho}{|G|} |\rho(H)A_j|_2^2$ .

What we consider here is the effect of choosing  $A$  from the Haar distribution in the unitary group. The expected value of the probability of measuring the  $j$ -th column is the same for all  $j$ , since each  $A_j$  is uniformly distributed on the unit sphere. So the averaged distribution on columns is the uniform distribution  $\overline{P_H} = \overline{P_H}(|\rho, j\rangle) = \int P_H(|A^{-1}\rho A, j\rangle) dA = \frac{1}{d_\rho} P_H(|\rho\rangle)$ .

Let  $\alpha = \frac{\sqrt{|G|}}{|H|\sqrt{c(G)}}$  where  $c(G)$  is the number of conjugacy classes in  $G$ . This parameter reflects the apparent difficulty of the hidden subgroup problem that is due to the small size of  $H$  and the degree of nonabelianness of  $G$ .

**Theorem 11.** *Let  $\varepsilon = \left(\frac{1}{\alpha} \frac{54}{2-\sqrt{3}} \ln \frac{4|G|}{\delta}\right)^{1/3}$ . Then with probability at least  $1-\delta$  (over the choice of random basis for the Fourier transform),  $\left|P_H - \overline{P_H}\right|_1 \leq \varepsilon$ .*

**Proof.** Our task is to show the following for sufficiently large  $\alpha$ . If, in each irrep,  $A$  is chosen from the Haar measure on the unitary group, then, for  $\rho$  sampled from the distribution  $P_H(|\rho\rangle)$ , almost certainly the probability  $\frac{d_\rho}{|G|} |\rho(H)A_j|_2^2$  of measuring the  $j$ -th column is close to its expectation  $\frac{1}{d_\rho} P_H(|\rho\rangle)$ . This amounts to bounding the  $L_1$  distance between the vectors  $P_H$  and  $\overline{P_H}$ . We consider separately irreps  $\rho$  according to whether  $\text{rank}(\rho(H))$  is higher or lower than the threshold  $T = \varepsilon\alpha/3$ .

**Case I.** For the high rank case, we show that with probability at least  $1-\delta$ , for all  $\rho$  and all  $j$ ,  $|\rho(H)A_j|_2^2$  deviates from its expectation by at most a  $2\varepsilon/3$  fraction. (Since we are concerned here only with fractional error we have suppressed the leading scale factor of the projection  $\frac{1}{\sqrt{|H|}}\rho(H)$ .)

What we are considering is the following process: a unit vector is chosen uniformly in  $\mathbb{C}^{d_\rho}$ , then projected onto a fixed subspace of dimension  $t > T$ ; by appropriate change of basis we can without loss of generality suppose that the subspace is spanned by the first  $t$  basis vectors of  $\mathbb{C}^{d_\rho}$ . Let  $s$  be the probability that the squared length of the projected vector differs from its expectation  $t/d_\rho$  by a fraction greater than  $2\varepsilon/3$ . Since we will apply a union bound over all  $\rho$  and  $j$ , it suffices to show that  $s \leq \delta/|G|$ . To begin with, note that, due to the isometric correspondence between the unit spheres in  $\mathbb{C}^{d_\rho}$  and  $\mathbb{R}^{2d_\rho}$ , the problem is equivalent to the same problem in real spaces of twice the dimensions, namely projection of the unit sphere in  $\mathbb{R}^{2d_\rho}$  onto a  $2t$ -dimensional subspace. Let  $M$  denote the projection matrix; in the appropriate basis it is diagonal, with  $2t$  1's on the diagonal.

We analyze the uniform sampling from the unit sphere indirectly, approximating it by the process of sampling a vector  $v$  from the spherically symmetric,  $2d_\rho$ -dimensional unit variance Gaussian distribution. Let the projection of  $v$  be  $v' = Mv$ . (Note that  $v'$  is distributed according to a  $2t$ -dimensional Gaussian distribution of variance  $t/d_\rho$ .) Then  $\frac{v'}{|v|_2}$  has the same distribution as  $\frac{1}{\sqrt{|H|}}\rho(H)A_j$  (with the understanding that pairs of real coordinates in the first vector form individual complex coordinates in the second). The probability  $s$  that  $\left|\frac{v'}{|v|_2}\right|_2$  deviates from its expectation by fraction  $2\varepsilon/3$  is bounded by the sum of the probabilities that  $|v|_2$  and  $|v'|_2$  deviate from their expectations by fraction  $\varepsilon/3$ .

We use the following large deviation bound: if  $a_1, \dots, a_\tau$  are independent Gaussian random variables each with unit standard deviation, then

$$P\left(\left|\frac{1}{\tau}\sum(a_i^2 - 1)\right| > \varepsilon\right) < 2\left[(1 + \varepsilon)^{1/2}e^{-\varepsilon/2}\right]^\tau.$$

For  $\varepsilon \leq 2$ ,  $(1 + \varepsilon)^{1/2}e^{-\varepsilon/2} \leq \exp\left(-\varepsilon^2\frac{2-\sqrt{3}}{4}\right)$ , and therefore

$$P\left(\left|\frac{1}{\tau}\sum(a_i^2 - 1)\right| > \varepsilon\right) < 2\exp\left(-\tau\varepsilon^2\frac{2-\sqrt{3}}{4}\right).$$

Since this bound is decreasing in  $\tau$ , and we are applying it with  $\tau = 2t > 2T$ , we conclude that

$$s < 4 \exp \left( -2T(\varepsilon/3)^2 \frac{2 - \sqrt{3}}{4} \right) = 4 \exp \left( -\alpha\varepsilon^3 \frac{2 - \sqrt{3}}{54} \right).$$

In order to ensure that  $s \leq \delta/|G|$  it suffices therefore that

$$\alpha\varepsilon^3 \geq \frac{54}{2 - \sqrt{3}} \ln \frac{4|G|}{\delta}$$

as assumed. Therefore the  $L_1$  distance between  $P_H$  and  $\overline{P_H}$  due to high rank irreps is at most  $2\varepsilon/3$ .

**Case II.** In the case that the rank of  $\rho$  is low,  $\text{rank}(\rho(H)) \leq T$ , we can no longer obtain a strong concentration bound on the probability of sampling each column. Instead we will show that Fourier sampling picks such an irrep with probability  $p_T \leq \varepsilon/3$ .

Let  $K = \sum_{\rho} d_{\rho}$ . We upper bound  $K$  by applying the Cauchy–Schwartz inequality to the two vectors  $(1)_{\rho}$  and  $(d_{\rho})_{\rho}$ . The norm squared of the first vector is simply  $c(G)$ , the number of irreps of  $G$ . Now

$$K \leq \left( \sum_{\rho} 1 \right)^{1/2} \left( \sum_{\rho} d_{\rho}^2 \right)^{1/2} = c(G)^{1/2} |G|^{1/2}.$$

By [Corollary 3](#),  $P_H(|\rho\rangle) = \frac{|H|d_{\rho}}{|G|} \text{rank}(\rho(H))$ . So

$$\begin{aligned} p_T &= \sum_{\rho: \text{rank}(\rho(H)) \leq T} P_H(|\rho\rangle) \\ &\leq \frac{|H|T}{|G|} \sum_{\rho: \text{rank}(\rho(H)) \leq T} d_{\rho} \\ &\leq \frac{|H|T}{|G|} \sum_{\rho} d_{\rho} \\ &= \frac{|H|TK}{|G|} \end{aligned}$$

which from the preceding argument is bounded above by  $\frac{|H|T\sqrt{c(G)}}{\sqrt{|G|}} = T/\alpha$ .

Since we chose  $T = \varepsilon\alpha/3$ , this gives  $p_T \leq \varepsilon/3$ . ■

**Corollary 12.** *With probability at least  $1 - \delta$  (over the choice of random basis for the Fourier transform),  $\Omega\left(\left(\frac{\alpha}{\log(|G|/\delta)}\right)^{1/3}\right)$  repetitions of Fourier sampling are required in order to achieve constant bias in distinguishing any two (a priori equally probable) conjugate subgroups  $H$  and  $H'$ .*

**Proof.** The Hellinger distance  $D_H(\vec{p}, \vec{q}) = \sum (p^{1/2} - q^{1/2})^2$  between two distributions  $\vec{p}$  and  $\vec{q}$  is additive across independent samples, and obeys the inequalities

$$|\vec{p} - \vec{q}|_1^2 / 4 \leq D_H(\vec{p}, \vec{q}) \leq |\vec{p} - \vec{q}|_1. \quad \blacksquare$$

As an example consider the symmetric group  $G = S_n$ ; we know that  $c(G) = \exp(\Theta(\sqrt{n}))$ . If  $|H| \leq |G|^{1/2-\gamma}$  for a fixed  $\gamma > 0$  then we need exponentially many samples to gain useful information from  $j$ .

### 4.3. Distinguishing $|H|=2$ from $|H|=1$

The graph automorphism problem reduces, via polynomial time reductions [10], to determining the size of the automorphism group in the special case where that is known to be either 1 or 2. So, although the weak form of the standard method (which ignores row and column indices) cannot distinguish conjugate subgroups, one may still hope that it is useful for the graph automorphism problem.

In [theorem 13](#) we show, however, that the weak form of the method is not useful for this task. Taken in conjunction with the previous section, this means that even the strong form of the method (which measures row and column indices) implemented with random bases, cannot solve the graph automorphism problem efficiently.

Consider the problem of distinguishing  $H = \{e, s\}$  from  $H' = \{e\}$ . Let  $C(s)$  be the conjugacy class of  $s$ .

**Theorem 13.** *The  $L_1$  distance between the distributions on irreps obtained by Fourier sampling  $|H\rangle$  and  $|H'\rangle$ , is at most  $1/\sqrt{|C(s)|}$ .*

The equation  $P_H(|\rho\rangle) = \frac{d_\rho}{|G|} \sum_{h \in H} \chi_\rho(h)$  implies that for  $H'$ ,  $\rho$  is sampled with probability  $d_\rho^2/|G|$ ; while for  $H$ ,  $\rho$  is sampled with probability  $d_\rho(d_\rho + \chi_\rho(s))/|G|$ . So the  $L_1$  distance between the distributions is  $\frac{1}{|G|} \sum_\rho d_\rho |\chi_\rho(s)|$ .

We upper bound this using the Cauchy–Schwartz inequality and the following identities:

1.  $\sum_\rho d_\rho^2 = |G|$ .
2.  $|Z(s)| \cdot |C(s)| = |G|$ .

$$3. \sum_{\rho} |\chi_{\rho}(s)|^2 = |Z(s)|.$$

Here  $Z(s)$  is the centralizer of  $s$ .

(1) is basic. (2) follows by considering the action of  $G$  on itself by conjugation, since under this action  $Z(s)$  is the stabilizer of  $s$  and  $C(s)$  is the orbit of  $s$ . To see (3), which generalizes (1), recall that the unitary character table of  $G$  has conjugacy classes labeling columns, irreps labeling rows, and that the  $(\rho, s)$  entry is  $\sqrt{\frac{|C(s)|}{|G|}} \chi_{\rho}(s)$ . Since each column is unit norm,

$$1 = \sum_{\rho} \frac{|C(s)|}{|G|} |\chi_{\rho}(s)|^2.$$

With (2) this shows (3). Now we can apply Cauchy–Schwartz.

$$\begin{aligned} \left| |F| |H\rangle \right|^2 - |F| |H'\rangle \right|^2 \Big|_1 &= \frac{1}{|G|} \sum_{\rho} d_{\rho} |\chi_{\rho}(s)| \\ &\leq \frac{1}{|G|} \left[ \sum_{\rho} d_{\rho}^2 \right]^{1/2} \left[ \sum_{\rho} |\chi_{\rho}(s)|^2 \right]^{1/2} \\ &= (|Z(s)|/|G|)^{1/2} \\ &= |C(s)|^{-1/2}. \end{aligned}$$

■

For strong Fourier sampling, [theorems 11 and 13](#) give us:

**Corollary 14.** *Let  $\varepsilon$  and  $\delta$  be as in [theorem 11](#). With probability at least  $1 - \delta$  (relative to the random choice of basis), the  $L_1$  distance between the distributions on irreps, rows and columns obtained from strong Fourier sampling  $|H\rangle$  and  $|H'\rangle$ , is at most  $\varepsilon$ .*

There are examples in which it is challenging to compute  $s$  even though the conjugacy class  $C(s)$  is known. Observe that in such cases this quantum algorithm has at most a quadratic advantage over the simple probabilistic strategy of checking whether  $f(s') = f(e)$  for a random conjugate  $s'$ .

We apply [Theorem 13](#) in the case that  $s$  is an involution in  $G = S_n$ , i.e.  $s$  is a product of some  $k$  disjoint transpositions. In this case  $|C(s)| = \frac{n!}{2^k k! (n-2k)!}$ ; as a convenient lower bound on this quantity, count only those conjugates which transpose odd elements with even elements, of which there are  $\binom{\lceil n/2 \rceil}{k} \binom{\lfloor n/2 \rfloor}{k} k! \geq k!$ . So the  $L_1$  distance between the distributions on irreps is at most  $(k!)^{-1/2}$ . In the graph automorphism application  $k$  can be

proportional to  $n$ , in which case this is exponentially small. A similar bound was independently obtained by Hallgren, Russell and Ta-Shma [8].

Finally we combine this bound with [Theorem 11](#). Note that  $\alpha = \frac{\sqrt{n!}}{2 \cdot 2^{e(\sqrt{n})}} \in \exp\left(\frac{1}{2}n \ln n - O(n)\right)$ , so:

**Corollary 15.** *If we apply the strong form of the standard method, using a random basis, to the graph automorphism problem, then with probability at least  $1 - \delta$  the  $L_1$  distance between the Fourier sampling distribution given that the automorphism group is trivial, and the Fourier sampling distribution given that the automorphism group is of size 2 and contains an involution with  $k$  transpositions, is at most  $\exp\left(-\frac{1}{6}n \ln n + O(n)\right) \log^{\frac{1}{3}} \frac{1}{\delta} + (k!)^{-1/2}$ .*

Just as in [Corollary 12](#), the upper bound on  $L_1$  distance implies a lower bound on the number of samples which must be collected in order to distinguish the hypotheses reliably.

## 5. Discussion

It is natural to propose an even stronger lower bound on the Fourier sampling method than that shown in [section 4.2](#), namely that there is no unitary choice of basis for the irreps that allows information the hidden subgroup  $H$  to be efficiently reconstructed.

A concrete approach to this problem would be to show that, for any fixed basis for an irrep  $\rho$ , if a subgroup  $H$  is selected at random from some distribution then  $P_H$  is with high probability close to uniform. It would be necessary to make assumptions (as in [theorem 11](#)) about the nonabelianness of  $G$ , the size of  $H$  and the rank of  $\rho(|H\rangle)$ ; while a natural choice for the distribution, is the uniform distribution on the conjugates of a subgroup possessing many conjugates. (The number of such conjugates is itself one quantification of the nonabelianness of the group.)

We already know that there is no information about the hidden subgroup in the row distributions. The question we face is this: a fixed vector  $v$  in  $\mathbb{C}^{d_\rho}$  is mapped by a uniformly chosen element of  $G$  to the vector  $\rho(g)(v)$ . The objective is to show a large deviation result for the norm squared of the projection by  $\rho(|H\rangle)$  of this random vector. More abstractly, we ask whether it is possible to make a statement of the following nature: let the projection operator  $\rho(|H\rangle)$  be of sufficiently high rank, and let  $v$  be an arbitrary vector. Prove a sharp concentration result for the norm-squares of the projection of  $\rho(g)(v)$  on  $\rho(|H\rangle)$ , for  $g$  chosen uniformly in  $G$ .



This is a very natural geometric conjecture about representations. Proving it (with sufficiently strong parameters) would yield as an immediate consequence that Fourier sampling cannot efficiently tell apart conjugate subgroups. Even more, the conjecture would imply that, for example, Fourier sampling cannot efficiently tell apart the trivial subgroup from a random subgroup of order 2, since averaging over conjugate subgroups gives the uniform distribution on columns, which is the same as that induced by the trivial subgroup. In particular, it would show that for  $G = S_n$ , even the decision problem for graph automorphism cannot be solved by Fourier sampling in any basis.

## References

- [1] R. BAER: Gruppen mit Hamiltonischem Kern, *Compositio Math.* **2** (1935), 241–246.
- [2] ROBERT BEALS: Quantum computation of Fourier transforms over symmetric groups, In *Proceedings of the 29th Annual ACM Symposium on Theory of Computing* (1997), 48–53.
- [3] R. A. BRYCE: The subgroups of Baer and Hughes, *Arch. Math.* **61** (1993), 305–312.
- [4] MICHAEL CLAUSEN: Fast generalized Fourier transforms, *Theor. Comp. Sci.* **67** (1989), 55–63.
- [5] PERSI DIACONIS and DANIEL ROCKMORE: Efficient computation of the Fourier transform of finite groups, *J. Amer. Math. Soc.* **3(2)** (1990), 297–332.
- [6] MARK ETTINGER and PETER HØYER: On quantum algorithms for noncommutative hidden subgroups, *Advances in Applied Mathematics* **25(3)** (2000), 239–251. (LANL preprint quant-ph/9807029, 1998.)
- [7] MARK ETTINGER, PETER HØYER and EMANUEL KNILL: Hidden subgroup states are almost orthogonal, LANL preprint quant-ph/9901034.
- [8] SEAN HALLGREN, ALEXANDER RUSSELL and AMNON TA-SHMA: Normal subgroup reconstruction and quantum computation using group representations, In *Proceedings of the 32nd Annual ACM Symposium on Theory of Computing* (2000), 627–635.
- [9] ALEXEI KITAEV: Quantum measurements and the abelian stabilizer problem, ECC Report TR96-003, 1996.
- [10] JOHANNES KÖBLER, UWE SCHÖNING and JACOBO TORÁN: *The Graph Isomorphism Problem: Its Structural Complexity*, Birkhäuser, 1993.
- [11] MARKUS PÜSCHEL, MARTIN RÖTTELER and THOMAS BETH: Fast quantum Fourier transforms for a class of non-abelian groups, LANL e-preprint quant-ph/9807064, 1998.
- [12] DANIEL ROCKMORE: Computation of Fourier Transforms on the Symmetric Group, in: E. Kaltofen, S. M. Watt (eds.), *Computers and Mathematics*, Springer, 1989.
- [13] JEAN-PIERRE SERRE: *Linear Representations of Finite Groups*, Springer-Verlag, 1977.
- [14] PETER SHOR: Polynomial-time algorithms for prime factorization and discrete logarithms on a quantum computer. *SIAM Journal on Computing* **26(5)** (1997), 1484–1509.
- [15] DANIEL SIMON: On the power of quantum computation, *SIAM Journal on Computing* **26(5)** (1997), 1474–1483.

- [16] WIM VAN DAM and SEAN HALLGREN: Efficient Quantum Algorithms for Shifted Quadratic Character Problems, LANL e-preprint quant-ph/0011067, 2000.

Michelangelo Grigni

*Mathematics and Computer Science*

*Emory University*

*400 Dowman Dr., Suite 401*

*Atlanta GA 30322*

*USA*

[mic@mathcs.emory.edu](mailto:mic@mathcs.emory.edu)

Leonard J. Schulman

*Computer Science*

*Caltech*

*1200 E. California Blvd., MC 258-80*

*Pasadena CA 91125*

*USA*

[schulman@caltech.edu](mailto:schulman@caltech.edu)

Monica Vazirani

*Mathematics*

*U. C. Davis*

*One Shields Ave.*

*Davis CA 95616-8633*

*USA*

[vazirani@math.ucdavis.edu](mailto:vazirani@math.ucdavis.edu)

Umesh Vazirani

*Computer Science Division*

*U. C. Berkeley*

*Berkeley CA 94720*

*USA*

[vazirani@cs.berkeley.edu](mailto:vazirani@cs.berkeley.edu)