

## Homework 4

due March 3

**Problem 1.**

(1) Recall from class that

$$h_n = \sum_{\lambda \vdash n} \frac{1}{z_\lambda} p_\lambda,$$

where  $z_\lambda = \prod_i i^{m_i} m_i!$  for  $\lambda = (1^{m_1}, 2^{m_2}, \dots)$ . Show that this is equivalent to Newton's determinant formula

$$h_n = \frac{1}{n!} \det \begin{pmatrix} p_1 & -1 & 0 & \dots & 0 \\ p_2 & p_1 & -2 & \dots & 0 \\ \vdots & \vdots & \vdots & & \vdots \\ p_{n-1} & p_{n-2} & \cdot & \dots & -(n-1) \\ p_n & p_{n-1} & \cdot & \dots & p_1 \end{pmatrix}$$

(2) Show that  $e_n$  is given by the same determinant without the minus signs.

**Problem 2.** Prove the identity

$$s_{(n-1, n-2, \dots, 1)}(x_1, \dots, x_n) = \prod_{1 \leq i < j \leq n} (x_i + x_j).$$

**Problem 3.** Permutation  $\pi = x_1 x_2 \cdots x_n$  has a descent at index  $i$  if  $x_i > x_{i+1}$ . The corresponding descent set is

$$\text{Des } \pi = \{i \mid i \text{ is a descent of } \pi\}.$$

Similarly,  $i$  is a descent for standard tableau  $T$  if  $i+1$  is in a lower row than  $i$  in  $T$  (in English notation). The descent set of  $T$  is

$$\text{Des } T = \{i \mid i \text{ is a descent of } T\}.$$

- (1) Show that if, by Robinson-Schensted,  $Q(\pi) = Q$ , then  $\text{Des } \pi = \text{Des } Q$ .
- (2) Let  $\lambda \vdash n$ ,  $S = \{n_1 < n_2 < \cdots < n_k\} \subset \{1, 2, \dots, n-1\}$ , and  $\mu = (n_1, n_2 - n_1, \dots, n - n_k)$ . Then

$$|\{\pi \in S_n \mid \text{shape}(Q(\pi)) = \lambda, \text{Des } \pi \subset S\}| = f^\lambda K_{\lambda\mu}.$$

**Problem 4.** Let  $T$  be a standard Young tableau of shape  $\lambda$ , and let  $b$  be its corner box occupied by entry 1. Define

$$\Delta(T) = \text{jdt}_b(\tilde{T})$$

where  $\tilde{T}$  is the skew standard tableau of shape  $\lambda/(1)$  obtained from  $T$  by removing the box  $b$  and subsequently decreasing all the remaining entries by one. The evacuation tableau  $\text{evac}(T)$  is the standard tableau of shape  $\lambda$  encoded by the sequence of shapes of

$$\emptyset, \Delta^{n-1}(T), \Delta^{n-2}(T), \dots, \Delta^2(T), \Delta(T), T.$$

For a permutation  $\pi = x_1 x_2 \cdots x_n \in S_n$  define

$$w^\# = (n+1-x_n)(n+1-x_{n-1}) \cdots (n+1-x_1).$$

- (1) Show: If  $w$  is mapped to  $(P, Q)$  under RSK, then  $w^\#$  is mapped to  $(\text{evac}(P), \text{evac}(Q))$  under RSK.
- (2) Show that  $\text{evac}$  is an involution.