

## Homework Set 4: Exercises on Linear Maps

**Directions:** Please work on all of the following problems! Hand in the Computational Problems 1 and 2, and the Proof-Writing Problems 6 and 7 at the **beginning** of lecture on February 2, 2007.

As usual, we are using  $\mathbb{F}$  to denote either  $\mathbb{R}$  or  $\mathbb{C}$ .

1. Define the map  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  by  $T(x, y) = (x + y, x)$ .
  - (a) Show that  $T$  is linear.
  - (b) Show that  $T$  is surjective.
  - (c) Find  $\dim \text{null}T$ .
  - (d) Find the matrix for  $T$  with respect to the canonical basis of  $\mathbb{R}^2$ .
  - (e) Show that the map  $F : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  given by  $F(x, y) = (x + y, x + 1)$  is not linear.
2. Consider the complex vector spaces  $\mathbb{C}^2$  and  $\mathbb{C}^3$  with their canonical bases. Let  $S : \mathbb{C}^3 \rightarrow \mathbb{C}^2$  be defined by the matrix

$$M(S) = A = \begin{pmatrix} i & 1 & 1 \\ 2i & -1 & -1 \end{pmatrix}.$$

Find a basis for  $\text{null}S$ .

3. Give an example of a function  $f : \mathbb{R}^2 \rightarrow \mathbb{R}$  having the property that

$$\forall a \in \mathbb{R}, \forall v \in \mathbb{R}, f(av) = af(v)$$

but such that  $f$  is not a linear map.

4. Let  $V$  and  $W$  be vector spaces over  $\mathbb{F}$  with  $V$  finite-dimensional, and let  $U$  be any subspace of  $V$ . Given a linear map  $S \in \mathcal{L}(U, W)$ , prove that there exists a linear map  $T \in \mathcal{L}(V, W)$  such that, for every  $u \in U$ ,  $S(u) = T(u)$ .
5. Let  $V$  and  $W$  be vector spaces over  $\mathbb{F}$ , and suppose that  $T \in \mathcal{L}(V, W)$  is injective. Given a linearly independent list  $(v_1, \dots, v_n)$  of vectors in  $V$ , prove that the list  $(T(v_1), \dots, T(v_n))$  is linearly independent in  $W$ .

6. Let  $U$ ,  $V$ , and  $W$  be vector spaces over  $\mathbb{F}$ , and suppose that the linear maps  $S \in \mathcal{L}(U, V)$  and  $T \in \mathcal{L}(V, W)$  are both injective. Prove that the composition map  $T \circ S$  is also injective.
7. Let  $V$  and  $W$  be vector spaces over  $\mathbb{F}$ , and suppose that  $T \in \mathcal{L}(V, W)$  is surjective. Given a spanning list  $(v_1, \dots, v_n)$  for  $V$ , prove that  $\text{span}(T(v_1), \dots, T(v_n)) = W$ .
8. Let  $V$  and  $W$  be vector spaces over  $\mathbb{F}$  with  $V$  finite-dimensional. Given  $T \in \mathcal{L}(V, W)$ , prove that there is a subspace  $U$  of  $V$  such that

$$U \cap \text{null}(T) = \{0\} \quad \text{and} \quad \text{range}(T) = \{T(u) \mid u \in U\}.$$

9. Show that the linear map  $T : \mathbb{F}^4 \rightarrow \mathbb{F}^2$  is surjective if

$$\text{null}(T) = \{(x_1, x_2, x_3, x_4) \in \mathbb{F}^4 \mid x_1 = 5x_2, x_3 = 7x_4\}.$$

10. Show that no linear map  $T : \mathbb{F}^5 \rightarrow \mathbb{F}^2$  can have as its null space the set

$$\{(x_1, x_2, x_3, x_4, x_5) \in \mathbb{F}^5 \mid x_1 = 3x_2, x_3 = x_4 = x_5\}.$$

11. Let  $V$  be a vector spaces over  $\mathbb{F}$ , and suppose that there is a linear map  $T \in \mathcal{L}(V, V)$  such that both  $\text{null}(T)$  and  $\text{range}(T)$  are finite-dimensional subspaces of  $V$ . Prove that  $V$  must also be finite-dimensional.