University of California Davis Abstract Linear Algebra MAT 67	Name (Print): Student ID (Print):	
Practice Midterm Examination Time Limit: 50 Minutes		April 26 2024

This examination document contains 7 pages, including this cover page, and 4 problems. You must verify whether there any pages missing, in which case you should let the instructor know. Fill in all the requested information on the top of this page, and put your initials on the top of every page, in case the pages become separated.

You may *not* use your books, notes, or any calculator on this exam.

You are required to show your work on each problem on this exam. The following rules apply:

- (A) If you use a lemma, proposition or theorem which we have seen in the class or in the book, you must indicate this and explain why the theorem may be applied.
- (B) Organize your work, in a reasonably neat and coherent way, in the space provided. Work scattered all over the page without a clear ordering will receive little credit.
- (C) Mysterious or unsupported answers will not receive full credit. A correct answer, unsupported by calculations, explanation, or algebraic work will receive little credit; an incorrect answer supported by substantially correct calculations and explanations will receive
- this.

	partial	credit.					
(D)	If you	need m	ore space	e, use	the	back	of the
	pages;	clearly	indicate	when	you	have	done

Do not write in the table to the right.

1. (25 points) Let  $V = \mathbb{R}^4$  and consider the two subsets

$$U_1 := \{(x_1, x_2, x_3, x_4) \in \mathbb{R}^4 : x_1 + x_2 + 2x_3 + x_4 = 0\},\$$

$$U_2 := \{(x_1, x_2, x_3, x_4) \in \mathbb{R}^4 : 3x_1 - x_2 + 8x_3 = 0, x_4 = 0\}.$$

(a) (5 points) Show that  $U_1 \subseteq V$  is a vector subspace.

(b) (10 points) Show that  $V = U_1 + U_2$ .

(c) (5 points) Find a vector  $v \in V$  such that  $U_1 \cap U_2 = \operatorname{span}(v)$ .

(d) (5 points) Prove that  $V \neq U_1 \oplus U_2$ .

2. (25 points) Consider the vector space  $V = \mathbb{R}^3$  and the vectors

$$v_1 = (3, 4, -1), \quad v_2 = (0, 2, 5), \quad v_3 = (-6, -2, 17).$$

(a) (10 points) Decide whether  $v_3$  is a linear combination of  $\{v_1, v_2\}$ .

(b) (5 points) Is  $v_1$  a linear combination of  $\{v_2, v_3\}$ ?

(c) (5 points) Show that (1,0,0) is *not* a linear combination of  $\{v_1,v_2,v_3\}$ .

(d) (5 points) Find a vector  $w \in V$  such that  $V = \operatorname{span}(v_1, v_2) \oplus \operatorname{span}(w)$ .

- 3. (25 points) Consider  $V = \mathbb{R}^n$  for some  $n \in \mathbb{N}$ . Solve the following parts.
  - (a) (20 points) Let  $f: \mathbb{R}^n \longrightarrow \mathbb{R}$  be a linear function which is not equal to the zero function. Show that the subset

$$U_f := \{ v \in V : f(v) = 0 \}$$

is a vector subspace.

(b) (5 points) Show that there exists a  $w \in V$  such that  $V = U \oplus \text{span}(w)$ .

- 4. (25 points) For each of the sentences below, circle whether they are **true** or **false**. (You do *not* need to justify your answer.)
  - (a) (5 points) Let  $V = \mathbb{R}[x]$ . Then  $U = \{p(x) \in V : p(0) = 1\}$  is a vector subspace.
    - (1) True.

- (2) False.
- (b) (5 points) Let  $V = \mathbb{R}[x]$ . Then  $U = \{p(x) \in V : p(0) = p(1)\}$  is a vector subspace.
  - (1) True.

- (2) False.
- (c) (5 points) If  $v \in V$  is a vector, there exist different vectors  $w_1, w_2 \in V$  such that  $v + w_1 = 0$  and  $v + w_2 = 0$ .
  - (1) True.

- (2) False.
- (d) (5 points) The function  $f: \mathbb{R}^3 \longrightarrow \mathbb{R}^2$  given by

$$f(x_1, x_2, x_3) = (x_1 - 2x_2 + 6x_3, x_1 + 4x_2 - 1)$$

is a linear function.

(1) True.

- (2) False.
- (e) (5 points) The function  $f: \mathbb{R}^2 \longrightarrow \mathbb{R}^4$  given by

$$f(x_1, x_2) = (x_1, x_2, 3x_1x_2, x_1 + x_2^4)$$

is a linear function.

(1) True.

(2) False.