

1.) Do detailed graphing (See instruction sheet from class.) for each function

a.)  $y = x(x - 4)$  on the interval  $[0, 5]$

b.)  $y = x(x - 5)^4$

c.)  $f(x) = \frac{3x^2}{x - 4}$

d.)  $f(x) = 4\sqrt{x} - x$

2.) Consider the function  $f(x) = 1 - x^{2/3}$  on the interval  $[-1, 1]$ . Show that  $f(1) = f(-1) = 0$  but that  $f'(x)$  is never zero on the interval  $[-1, 1]$ . Explain how this is possible, in view of the Mean Value Theorem.

3.) Let  $f(x) = \begin{cases} -x^2, & \text{if } -1 \leq x \leq 0 \\ x^2(x - 1), & \text{if } 0 < x \leq 2. \end{cases}$

a.) Sketch the graph of  $f$ .

b.) Show that  $f$  satisfies the conditions of the Mean Value Theorem (MVT) over the interval  $[-1, 2]$ , including special attention at  $x = 0$ , and determine all values of  $c$  guaranteed by the MVT.

4.) Use a linearization to estimate the value of

a.)  $\sqrt{150}$

b.)  $e^{0.1}$

5.) The radius of a circle is measured with absolute percentage error of at most 3%. Use differentials to estimate the maximum absolute percentage error in computing the circle's

a.) circumference.

b.) area.

(RECALL: For a circle : circumference  $C = 2\pi r$  and area  $A = \pi r^2$ .)

6.) The radius of a sphere is measured with absolute percentage error of at most 4%. Use differentials to estimate the maximum absolute percentage error in computing the sphere's

a.) surface area.

b.) volume.

(RECALL: For a sphere : surface area  $S = 4\pi r^2$  and volume  $V = (4/3)\pi r^3$ .)

+++++

The following problem is for recreational purposes only.

7.) Find a hidden pattern and determine the next number in the sequence :

$0, 1, 3, 7, 14, 25, 41, 63, \dots$