

Math 21C

Kouba

Complex Numbers and Taylor Series

Recall : 1.) $\cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} + \dots$ for all x -values

2.) $\sin x = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!} + \dots$ for all x -values

3.) $e^x = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \frac{x^4}{4!} + \dots$ for all x -values

Question : Is there a mathematical connection between e^x and the trig functions $\cos x$ and $\sin x$? Yes ...

Consider the complex number $z = i\theta$ where θ is a real number. Then

$$\begin{aligned} e^z &= 1 + z + \frac{z^2}{2!} + \frac{z^3}{3!} + \frac{z^4}{4!} + \dots \\ &= 1 + (i\theta) + \frac{(i\theta)^2}{2!} + \frac{(i\theta)^3}{3!} + \frac{(i\theta)^4}{4!} + \dots \\ &= 1 + i\theta + (i^2)\frac{\theta^2}{2!} + \frac{(i^3)\theta^3}{3!} + \frac{(i^4)\theta^4}{4!} + \frac{(i^5)\theta^5}{5!} + \dots \\ &= 1 + i\theta - \frac{\theta^2}{2!} - i\frac{\theta^3}{3!} + \frac{\theta^4}{4!} + i\frac{\theta^5}{5!} - \dots \\ &= \left(1 - \frac{\theta^2}{2!} + \frac{\theta^4}{4!} - \frac{\theta^6}{6!} + \dots\right) \\ &\quad + i\left(\theta - \frac{\theta^3}{3!} + \frac{\theta^5}{5!} - \frac{\theta^7}{7!} + \dots\right) \\ &= \cos \theta + i \sin \theta, \text{ i.e.,} \end{aligned}$$

$$e^{i\theta} = \cos \theta + i \sin \theta$$

for all real numbers θ .

Ex: Simplify each expression.

$$\begin{aligned} 1.) 4e^{i\frac{\pi}{6}} &= 4 \left(\cos \frac{\pi}{6} + i \sin \frac{\pi}{6} \right) \\ &= 4 \left(\frac{\sqrt{3}}{2} + i \cdot \frac{1}{2} \right) = 2\sqrt{3} + 2i \end{aligned}$$

$$\begin{aligned} 2.) (e^{i\frac{\pi}{4}})^7 &= e^{i \cdot \frac{7\pi}{4}} = \cos \frac{7\pi}{4} + i \sin \frac{7\pi}{4} \\ &= \frac{\sqrt{2}}{2} - i \cdot \frac{\sqrt{2}}{2} \end{aligned}$$

$$\begin{aligned} 3.) e^{i+\ln 3} &= e^i e^{\ln 3} = 3(\cos 1 + i \sin 1) \\ &= 3 \cos 1 + 3i \sin 1 \end{aligned}$$

$$4.) e^{i\pi} = \cos \pi + i \sin \pi = -1 + i(0) = -1$$

$$5.) e^{2\pi i} = \cos 2\pi + i \sin 2\pi = 1 + i(0) = 1$$

$$6.) e^{4\pi i} = \cos 4\pi + i \sin 4\pi = 1 + i(0) = 1$$

$$7.) e^{-6\pi i} = \cos(-6\pi) + i \sin(-6\pi) = 1 + i(0) = 1$$

Ex: Find all complex numbers z which solve $e^z = 1+i$: Let $z = a+bi$. Then

$$e^z = 1+i \Rightarrow e^{a+bi} = 1+i \Rightarrow$$

$$e^a e^{bi} = 1+i \Rightarrow e^a (\cos b + i \sin b) = 1+i ;$$

(stop and think) $|1+i| = \sqrt{1^2+1^2} = \sqrt{2}$ and

$$1+i = \sqrt{2} \cdot \frac{1+i}{\sqrt{2}} = \sqrt{2} \left(\frac{1}{\sqrt{2}} + i \cdot \frac{1}{\sqrt{2}} \right), \text{ so that}$$

$$e^a (\cos b + i \sin b) = \sqrt{2} \left(\frac{1}{\sqrt{2}} + i \cdot \frac{1}{\sqrt{2}} \right) \Rightarrow e^a = \sqrt{2}$$

$$\Rightarrow \boxed{a = \ln \sqrt{2}} \text{ and } \cos b = \frac{1}{\sqrt{2}}, \sin b = \frac{1}{\sqrt{2}} \Rightarrow$$

$$\boxed{b = \frac{\pi}{4} \pm 2\pi n}$$

$$\text{for } n=0,1,2,3,\dots, \text{ i.e., } \boxed{z = \ln \sqrt{2} + i \left(\frac{\pi}{4} \pm 2\pi n \right)}$$

$$\text{for } n=0,1,2,3,\dots$$