

Homework Set No. 9 – Probability Theory (235A), Fall 2011

Due: 11/29/09

1. (a) If X is a r.v., show that $\operatorname{Re}(\varphi_X)$ (the real part of φ_X) and $|\varphi_X|^2 = \varphi_X \overline{\varphi_X}$ are also characteristic functions (i.e., construct r.v.'s Y and Z such that $\varphi_Y(t) = \operatorname{Re}(\varphi_X(t))$, $\varphi_Z(t) = |\varphi_X(t)|^2$).

(b) Show that X is equal in distribution to $-X$ if and only if φ_X is a real-valued function.

2. (a) Let Z_1, Z_2, \dots be a sequence of independent r.v.'s such that the random series $X = \sum_{n=1}^{\infty} Z_n$ converges a.s. Prove that

$$\varphi_X(t) = \prod_{n=1}^{\infty} \varphi_{Z_n}(t), \quad (t \in \mathbb{R}).$$

(b) Let X be a uniform r.v. in $(0, 1)$, and let Y_1, Y_2, \dots be the (random) bits in its binary expansion, i.e. each Y_n is either 0 or 1, and the equation

$$X = \sum_{n=1}^{\infty} \frac{Y_n}{2^n} \tag{1}$$

holds. Show that Y_1, Y_2, \dots are i.i.d. unbiased coin tosses (i.e., taking values 0, 1 with probabilities $1/2, 1/2$).

(c) Compute the characteristic function φ_Z of $Z = 2X - 1$ (which is uniform in $(-1, 1)$). Use (1) to represent this in terms of the characteristic functions of the Y_n 's (note that the series (1) converges absolutely, so here there is no need to worry about almost sure convergence). Deduce the infinite product identity

$$\frac{\sin(t)}{t} = \prod_{n=1}^{\infty} \cos\left(\frac{t}{2^n}\right), \quad (t \in \mathbb{R}). \tag{2}$$

(d) Substitute $t = \pi/2$ in (2) to get the identity

$$\frac{2}{\pi} = \frac{\sqrt{2}}{2} \cdot \frac{\sqrt{2 + \sqrt{2}}}{2} \cdot \frac{\sqrt{2 + \sqrt{2 + \sqrt{2}}}}{2} \cdot \dots$$

3. Let X be a r.v. From the inversion formula, it follows without much difficulty (see p. 95 in Durrett's book, 3rd or 4th eds.), that if φ_X is integrable, then X has a density f_X , and

the density and characteristic function are related by

$$\begin{aligned}\varphi_X(t) &= \int_{-\infty}^{\infty} f_X(x)e^{itx} dx, \\ f_X(x) &= \frac{1}{2\pi} \int_{-\infty}^{\infty} \varphi_X(t)e^{-itx} dt\end{aligned}$$

(this shows the duality between the Fourier transform and its inverse). Use this and the answer to question 4(e) in homework assignment #8 to conclude that if X is a r.v. with the Cauchy distribution (i.e., X has density $f_X(x) = 1/\pi(1 + x^2)$) then its characteristic function is given by

$$\varphi_X(t) = e^{-|t|}.$$

Deduce from this that if X, Y are independent Cauchy r.v.'s then any weighted average $\lambda X + (1 - \lambda)Y$, where $0 \leq \lambda \leq 1$, is also a Cauchy r.v. (As a special case, it follows by induction that if X_1, \dots, X_n are i.i.d. Cauchy r.v.'s, then their average $(X_1 + \dots + X_n)/n$ is also a Cauchy r.v., which was a claim we made without proof earlier in the course.)

Happy Thanksgiving!