


Lecture 4: The Isoperimetric Problem

Note Title

★ The Problem of Queen Dido

See the reference page for its rich history.



Find a curve $y = y(x)$ with $y(0) = y(1) = 0$ s.t.

the enclosed area

$$I = \int_0^1 y \, dx \rightarrow \max!$$

$$\text{subj. to } C = \int_0^1 \sqrt{1 + y'^2} \, dx = \text{const.}, \text{ say } \frac{\pi}{2}.$$

The difference from the previous variational problem is the constraint (not in the form of B.C.). See also P. Lax's nice short article based on non-variational approach!

A more general form is:

Find a fn $y = y(x)$ s.t.

$$(*) \begin{cases} \text{B.C. : } y(x_i) = y_i, \quad i=1,2. \\ \text{fncl : } I = \int_{x_1}^{x_2} f(x, y, y') \, dx \rightarrow \max (\text{or min}) \\ \text{constr : } C = \int_{x_1}^{x_2} g(x, y, y') \, dx = L (\text{const.}) \end{cases}$$

★ The E-L Eqn. for the Isoperimetric Problem in One Indep. Variable

again, we consider the solution $y = y^*(x)$ to (*) and its variation. However, unlike the previous case, $y_\varepsilon(x) = y^*(x) + \varepsilon \eta(x)$, $\eta(x_i) = 0$, $i=1,2$ does not work, $\leftarrow C$ must be kept const. $\equiv L$. Adding $\varepsilon \eta$, C would change from L .

⇒ We need a two-param. variation to counter-balance this, i.e., consider

$$\begin{cases} y = y^*(x) + \varepsilon_1 \eta_1(x) + \varepsilon_2 \eta_2(x) \\ \eta_i(x_j) = 0, \quad i, j = 1, 2. \end{cases}$$

Substitute this into $I = \int f$ & $C = \int g$ to get:

$$(**) \begin{cases} I(\varepsilon_1, \varepsilon_2) |_{\varepsilon_1 = \varepsilon_2 = 0} = I(0, 0) = E \text{ (extreme val.)} \\ C(\varepsilon_1, \varepsilon_2) |_{\varepsilon_1 = \varepsilon_2 = 0} = C(0, 0) = L \end{cases}$$

For E to be an extreme value, the Jacobian must vanish at $(\varepsilon_1, \varepsilon_2) = (0, 0)$, i.e.,

$$\Delta := \frac{\partial(I, C)}{\partial(\varepsilon_1, \varepsilon_2)} \Big|_{\varepsilon_1 = \varepsilon_2 = 0} = \begin{vmatrix} \frac{\partial I}{\partial \varepsilon_1}(0, 0) & \frac{\partial I}{\partial \varepsilon_2}(0, 0) \\ \frac{\partial C}{\partial \varepsilon_1}(0, 0) & \frac{\partial C}{\partial \varepsilon_2}(0, 0) \end{vmatrix} = 0$$

Why? If $\Delta \neq 0$ at $(\varepsilon_1, \varepsilon_2) = (0, 0)$, then by the Implicit Fcn Thm, we could solve (**) in the neighborhood of $(\varepsilon_1, \varepsilon_2) = (0, 0)$ s.t.

$\varepsilon_i = \varepsilon_i(E, L)$, $i = 1, 2$, which are continuous.

Now, if E is an extreme val. (i.e., min or max) of I , then we'll get $\varepsilon_1 = \varepsilon_2 = 0$ by assumption.

But, since $\varepsilon_i = \varepsilon_i(E, L)$, $i = 1, 2$, we can get some $(\varepsilon_1, \varepsilon_2)$ for (E', L) with $E' < E$ or $E' > E$.

⇒ E cannot be an extreme val. #

Now, let's compute Δ !

Similarly to the previous variational problems, we can get:

$$\begin{cases} \frac{\partial I}{\partial \varepsilon_i}(0, 0) = \int_{x_1}^{x_2} \eta_i \left(\frac{\partial f}{\partial y} - \frac{d}{dx} \frac{\partial f}{\partial y'} \right) dx \\ \frac{\partial C}{\partial \varepsilon_i}(0, 0) = \int_{x_1}^{x_2} \eta_i \left(\frac{\partial g}{\partial y} - \frac{d}{dx} \frac{\partial g}{\partial y'} \right) dx \end{cases} \quad i = 1, 2.$$

Now, we assume that $y^*(x)$ is **not** a sol. of

$$\frac{\partial g}{\partial y} - \frac{d}{dx} \frac{\partial g}{\partial y'} = 0$$

\Rightarrow This assumpt. is reasonable since y^* maximizes (or minimizes) I and it's unlikely to make C extreme simultaneously.

\Rightarrow So, we can choose any admissible fcn $\zeta_2(x)$ s.t. $\int_{x_1}^{x_2} \zeta_2 \left(\frac{\partial g}{\partial y} - \frac{d}{dx} \frac{\partial g}{\partial y'} \right) dx \neq 0$.

Once we choose such ζ_2 , let

$$\int_{x_1}^{x_2} \zeta_2 \left(\frac{\partial g}{\partial y} - \frac{d}{dx} \frac{\partial g}{\partial y'} \right) dx =: \lambda_1 \neq 0$$

$$\int_{x_1}^{x_2} \zeta_1 \left(\frac{\partial f}{\partial y} - \frac{d}{dx} \frac{\partial f}{\partial y'} \right) dx =: \lambda_2$$

Then,

$$\Delta = \begin{vmatrix} \int_{x_1}^{x_2} \zeta_1 \left(\frac{\partial f}{\partial y} - \frac{d}{dx} \frac{\partial f}{\partial y'} \right) dx & \lambda_2 \\ \int_{x_1}^{x_2} \zeta_2 \left(\frac{\partial g}{\partial y} - \frac{d}{dx} \frac{\partial g}{\partial y'} \right) dx & \lambda_1 \end{vmatrix}$$

$$= \lambda_1 \int_{x_1}^{x_2} \zeta_1 \left(\frac{\partial f}{\partial y} - \frac{d}{dx} \frac{\partial f}{\partial y'} \right) dx - \lambda_2 \int_{x_1}^{x_2} \zeta_2 \left(\frac{\partial g}{\partial y} - \frac{d}{dx} \frac{\partial g}{\partial y'} \right) dx = 0$$

By setting $\lambda := -\lambda_2/\lambda_1$, we have

$$\int_{x_1}^{x_2} \zeta_1 \left[\frac{\partial f}{\partial y} - \frac{d}{dx} \frac{\partial f}{\partial y'} + \lambda \left(\frac{\partial g}{\partial y} - \frac{d}{dx} \frac{\partial g}{\partial y'} \right) \right] dx = 0$$

Using the F.L.C.V., we have the necessary cond. for the sol. to the isoperimetric problem (x) :

$$\frac{\partial}{\partial y} (f + \lambda g) - \frac{d}{dx} \frac{\partial}{\partial y'} (f + \lambda g) = 0 \quad \text{or}$$

$$\frac{\partial h}{\partial y} - \frac{d}{dx} \frac{\partial h}{\partial y'} = 0 \quad \text{with } h = f + \lambda g$$

This is a form of the **Lagrange multiplier**!
 λ is, at this point, an arbitrary parameter.
 Yet, λ must be used to satisfy the constraint
 $C = \int_{x_1}^{x_2} g \, dx = L$.

Let's check the Queen Dido problem.
 $f = y$, $g = \sqrt{1+y'^2} \Rightarrow h = y + \lambda \sqrt{1+y'^2}$

$$\frac{\partial h}{\partial y} = 1, \quad \frac{\partial h}{\partial y'} = \frac{\lambda y'}{\sqrt{1+y'^2}}$$


$$\text{So, } \frac{\partial h}{\partial y} - \frac{d}{dx} \frac{\partial h}{\partial y'} = 0 \Leftrightarrow 1 - \frac{d}{dx} \left(\frac{\lambda y'}{\sqrt{1+y'^2}} \right) = 0 \quad (***)$$

$$\text{Notice that } \frac{d}{dx} \left(\frac{y'}{\sqrt{1+y'^2}} \right) = \frac{y''}{(1+y'^2)^{3/2}} = \kappa$$

$$\text{So, } (***) \Leftrightarrow \lambda \kappa = 1 \text{ or } \kappa = \frac{1}{\lambda} = \text{const.}$$

curvature of $y(x)$.

This means that $y = y^*(x)$ is a **semicircle**
 with $L = \frac{\pi}{2}$.



$$y = y^*(x) = \sqrt{\frac{1}{4} - (x - \frac{1}{2})^2} \quad r = \frac{1}{2}$$

$$\kappa = \frac{1}{r} = 2, \text{ so } \lambda = \frac{1}{2} //$$

Exercise: What happens if $L \neq \frac{\pi}{2}$?