

A SIMPLE PROOF OF WELL-POSEDNESS FOR THE FREE-SURFACE INCOMPRESSIBLE EULER EQUATIONS

DANIEL COUTAND

Maxwell Institute for Mathematical Sciences and department of Mathematics
Heriot-Watt University, Edinburgh, EH14 4AS, UK

STEVE SHKOLLER

Department of Mathematics, University of California
Davis, CA 95616, USA

ABSTRACT. This paper contains the lectures presented by Shkoller at the Eleventh School on the Mathematical Theory in Fluid Mechanics at Kacov, Czech republic, in May, 2009, based on the well-posedness theory for the free-surface incompressible Euler equations developed in Coutand & Shkoller [3]. The purpose of those lectures and this paper is to present a new simple proof for the construction of solutions to the moving free-boundary incompressible 3-D Euler equations in vacuum. The method relies on the Lagrangian representation of the fluid, and the anisotropic smoothing operation that we call horizontal convolution-by-layers.

1. Introduction. This paper is founded upon the lectures presented by Shkoller at the Eleventh School on the Mathematical Theory in Fluid Mechanics at Kacov, Czech republic, in May, 2009 on the well-posedness of the free-surface incompressible Euler equations as developed in Coutand & Shkoller [3]. We are grateful to the organizers Josef Málek and Mirko Rokyta for both the opportunity to present these lectures, as well as for the exceptional summer school that they organized. We are also grateful to the many students who attended this summer school, and whose excitement for the subject matter motivated us greatly to provide a new and concise treatment of this well-posedness theory.

These lectures are focused on the incompressible Euler equations with moving free-surface. While we present the material for the case of three space dimensions, the same presentation covers the case of plane flow also. We will not consider the presence of surface tension in this paper, and we will not make any assumptions of irrotationality.

1.1. The Eulerian description. For $0 \leq t \leq T$, the evolution of a three-dimensional incompressible fluid with a moving free-surface is modeled by the incompressible

2000 *Mathematics Subject Classification.* 35L65, 35L70, 35L80, 35Q35, 35R35, 76B03.

Key words and phrases. Euler, water-waves, free boundary problems, vacuum.

Euler equations:

$$u_t + u \cdot Du + Dp = 0 \quad \text{in } \Omega(t), \quad (1a)$$

$$\operatorname{div} u = 0 \quad \text{in } \Omega(t), \quad (1b)$$

$$p = 0 \quad \text{on } \Gamma(t), \quad (1c)$$

$$\mathcal{V}(\Gamma(t)) = u \cdot n \quad (1d)$$

$$u = u_0 \quad \text{on } \Omega(0), \quad (1e)$$

$$\Omega(0) = \Omega. \quad (1f)$$

The open subset $\Omega(t) \subset \mathbb{R}^3$ denotes the changing volume occupied by the fluid, $\Gamma(t) := \partial\Omega(t)$ denotes the moving free-surface, $\mathcal{V}(\Gamma(t))$ denotes normal velocity of $\Gamma(t)$, and $n(t)$ denotes the exterior unit normal vector to the free-surface $\Gamma(t)$. The vector-field $u = (u_1, u_2, u_3)$ denotes the Eulerian velocity field, and p denotes the pressure function. We use the notation $D = (\partial_1, \partial_2, \partial_3)$ to denote the gradient operator. We have normalized the equations to have all physical constants equal to 1.

This is a free-boundary partial differential equation to determine the velocity and pressure in the fluid, as well as the location and smoothness of the a priori unknown free-surface. A recent explosion of interest in the analysis of the free-boundary incompressible Euler equations, particularly in irrotational form, has produced a number of different methodologies for obtaining a priori estimates, and the accompanying existence theories have mostly relied on the Nash-Moser iteration to deal with derivative loss in linearized equations when arbitrary domains are considered, or complex analysis tools for the irrotational problem with infinite depth. We refer the reader to [1], [3], [5], [6], [7], [8], [11], [12], [13], [14] for a partial list of papers on this topic.

The purpose of these lectures is to make our presentation in [3] more accessible, and to explain the basic interaction of the geometry of the free-surface with the motion of the fluid. As is often the case, with the passage of time, a more concise treatment can be presented, and this is indeed the case for the free-surface incompressible Euler problem.

To avoid the use of local coordinate charts necessary for arbitrary geometries, for simplicity, we will assume that the initial domain Ω at time $t = 0$ is given by

$$\Omega = \{(x_1, x_2, x_3) \in \mathbb{R}^3 \mid (x_1, x_2) \in \mathbb{T}^2, x_3 \in (0, 1)\}, \quad (2)$$

where \mathbb{T}^2 denotes the 2-torus, which can be thought of as the unit square $(0, 1)^2$ with periodic boundary conditions. This permits the use of *one* global Cartesian coordinate system. We only allow the *top* boundary

$$\Gamma = \{x_3 = 1\}$$

to move, while the *bottom* boundary is fixed with boundary condition

$$u_3 = 0 \quad \text{on } \{x_3 = 0\} \times [0, T].$$

We refer the reader to our paper [3] for the case that the initial domain is an arbitrary bounded, open subset of \mathbb{R}^3 with H^3 -class boundary.

1.2. Einstein's summation convention. Repeated Latin indices i, j, k, \dots , are summed from 1 to 3; for example, $F_{,ii} := \sum_{i=1,3} \frac{\partial^2}{\partial x_i \partial x_i}$.

1.3. The Lagrangian description. We transform the system (1) into Lagrangian variables. We let $\eta(x, t)$ denote the “position” of the fluid particle x at time t . Thus,

$$\partial_t \eta = u \circ \eta \text{ for } t > 0 \text{ and } \eta(x, 0) = x$$

where \circ denotes composition so that $[u \circ \eta](x, t) := u(\eta(x, t), t)$. We set

$$v = u \circ \eta \text{ (Lagrangian velocity),}$$

$$q = p \circ \eta \text{ (Lagrangian pressure),}$$

$$A = [D\eta]^{-1} \text{ (inverse of the deformation tensor),}$$

$$J = \det[D\eta] \text{ (Jacobian determinant of the deformation tensor),}$$

$$a = J A \text{ (cofactor of the deformation tensor).}$$

Since $\operatorname{div} u = 0$, we have that $\det D\eta = 1$, and hence the cofactor matrix of $D\eta$ is equal to $[D\eta]^{-1}$. Using Einstein’s summation convention, and using the notation $F_{,k}$ to denote $\frac{\partial F}{\partial x_k}$, the k th-partial derivative of F for $k = 1, 2, 3$, the Lagrangian version of equations (1) is given on the fixed reference domain Ω by

$$v_t^i + A_i^k q_{,k} = 0 \quad \text{in } \Omega \times (0, T], \quad (3a)$$

$$\operatorname{div}_\eta v = 0 \quad \text{in } \Omega \times (0, T], \quad (3b)$$

$$q = 0 \quad \text{on } \Gamma \times (0, T], \quad (3c)$$

$$v_3 = 0 \quad \text{on } \{x_3 = 0\} \times (0, T], \quad (3d)$$

$$(\eta, v) = (e, u_0) \quad \text{in } \Omega \times \{t = 0\}, \quad (3e)$$

where $e(x) = x$ denotes the identity map on Ω . Notice that the free surface $\Gamma(t)$ is given by

$$\Gamma(t) = \eta(t)(\Gamma).$$

The Lagrangian divergence $\operatorname{div}_\eta v = A_i^j v^i_{,j}$. Equation (3a) can be written in vector form as

$$v_t + A^T Dq = 0 \quad \text{in } \Omega \times (0, T], \quad (3a')$$

where A^T denotes the transpose of A . Solutions to (3) which are sufficiently smooth to ensure that $\eta(t)$ are diffeomorphisms, give solutions to (1) via the change of variables indicated above.

1.4. The Lagrangian vorticity equation. We make use of the permutation symbol

$$\varepsilon_{ijk} = \begin{cases} 1, & \text{even permutation of } \{1, 2, 3\}, \\ -1, & \text{odd permutation of } \{1, 2, 3\}, \\ 0, & \text{otherwise,} \end{cases}$$

and the basic identity regarding the i th component of the curl of a vector field u :

$$(\operatorname{curl} u)_i = \varepsilon_{ijk} u^k_{,j}.$$

Defining $\operatorname{curl}_\eta v = \operatorname{curl} u \circ \eta$, the chain rule shows, by taking the curl of the Euler equations (3a), that

$$(\operatorname{curl}_\eta v)_i = \varepsilon_{ijk} A_j^s v^k_{t,s} = 0. \quad (4)$$

2. Notation.

2.1. Differentiation and norms. For integers $k \geq 0$ and a smooth, open domain Ω of \mathbb{R}^3 , we define the Sobolev space $H^k(\Omega)$ ($H^k(\Omega; \mathbb{R}^3)$) to be the completion of $C^\infty(\Omega)$ ($C^\infty(\Omega; \mathbb{R}^3)$) in the norm

$$\|u\|_k^2 := \sum_{|a| \leq k} \int_{\Omega} |D^a u|^2 dx,$$

where D^a denotes all partial derivatives of order a . For real numbers $s \geq 0$, the Sobolev spaces $H^s(\Omega)$ and the norms $\|\cdot\|_s$ are defined by interpolation. We will write $H^s(\Omega)$ instead of $H^s(\Omega; \mathbb{R}^3)$.

We define the horizontal derivative by $\bar{\partial} = (\partial_1, \partial_2)$, and define the Sobolev space $H^k(\Gamma)$ to be the completion of $C^\infty(\Gamma)$ in the norm

$$|u|_k^2 := \sum_{|a| \leq k} \int_{\Gamma} |\bar{\partial}^a u|^2 dS,$$

where $\bar{\partial}^a$ denotes all horizontal partial derivatives of order a , and $dS = dx_1 dx_2$ denotes the ‘surface measure.’ For real $s \geq 0$, the Hilbert space $H^s(\Gamma)$ and the boundary norm $|\cdot|_s$ is defined by interpolation. The negative-order Sobolev spaces $H^{-s}(\Gamma)$ are defined via duality: for real $s \geq 0$, $H^{-s}(\Gamma) := [H^s(\Gamma)]'$.

2.2. The space of divergence-free vectors on Ω . In order to specify our initial velocity field, we introduce the following subspace of H^s vector-fields on Ω for $s \geq 0$:

Definition 2.1 (H^s -class divergence free vectors).

$$H_{\text{div}}^s(\Omega) = \{u \in H^s(\Omega; \mathbb{R}^3) : u^3 = 0 \text{ on } \{x_3 = 0\}, x_h \mapsto u(x_h, x_3) \text{ periodic} \\ \text{div } u = 0\}.$$

3. Properties of the cofactor matrix a , and a polynomial-type inequality.

3.1. Differentiating the inverse matrix A . Using that $D\eta A = \text{Id}$, we have the following identities

$$\bar{\partial} A_i^k = -A_i^s \bar{\partial} \eta^r{}_{,s} A_r^k, \quad (5)$$

$$D A_i^k = -A_i^s D \eta^r{}_{,s} A_r^k, \quad (6)$$

$$\partial_t A_i^k = -A_i^s v^r{}_{,s} A_r^k. \quad (7)$$

3.2. Relating the cofactor matrix and the unit normal $n(t)$. With $N = (0, 0, 1)$ the outward unit normal to Γ , we have the identity

$$n(\eta) = a^T N / |a^T N| \quad \text{or} \quad n_i(\eta) = a_i^3 / |a^3|.$$

For $\alpha, \beta = 1, 2$, we define the components of the *induced metric* on $\Gamma(t)$ by

$$g_{\alpha\beta}(x_1, x_2, t) = \eta_{,\alpha}(x_1, x_2, t) \cdot \eta_{,\beta}(x_1, x_2, t),$$

and let \sqrt{g} denote $\sqrt{\det(g_{\alpha\beta})}$. The metric g is a 2×2 matrix defined on Γ . It is an elementary computation (see [3]) to verify that

$$\sqrt{g} = |\eta_{,1} \times \eta_{,2}| = |a^3| \text{ on } \Gamma,$$

and hence that $a_i^3 = \sqrt{g} n_i(\eta)$.

$$A_i^3 = J^{-1} \sqrt{g} n_i(\eta) \text{ on } \Gamma. \quad (8)$$

3.3. A polynomial-type inequality. For a constant $M_0 \geq 0$, suppose that $f(t) \geq 0$, $t \mapsto f(t)$ is continuous, and

$$f(t) \leq M_0 + C t P(f(t)), \quad (9)$$

where P denotes a polynomial function, and C is a generic constant. Then for t taken sufficiently small, we have the bound

$$f(t) \leq 2M_0.$$

We use this type of inequality (see [3]) in place of nonlinear Gronwall-type of inequalities.

4. Trace estimates and the Hodge decomposition elliptic estimates. The normal trace theorem states that the existence of the normal trace $w \cdot N|_\Gamma$ of a velocity field $w \in L^2(\Omega)$ relies on the regularity of $\operatorname{div} w \in L^2(\Omega)$ (see, for example, [10]). If $\operatorname{div} w \in L^2(\Omega)$, then $w \cdot N$ exists in $H^{-0.5}(\Gamma)$. We will use the following variant:

$$\|\bar{\partial} w \cdot N\|_{H^{-0.5}(\Gamma)}^2 \leq C \left[\|\bar{\partial} w\|_{L^2(\Omega)}^2 + \|\operatorname{div} w\|_{L^2(\Omega)}^2 \right] \quad (10)$$

for some constant C independent of w .

The construction of our higher-order energy function is based on the following Hodge-type elliptic estimate:

Proposition 1. *For an H^r domain Ω with $\Gamma = \partial\Omega$, $r \geq 3$, if $F \in L^2(\Omega; \mathbb{R}^3)$ with $\operatorname{curl} F \in H^{s-1}(\Omega; \mathbb{R}^3)$, $\operatorname{div} F \in H^{s-1}(\Omega)$, and $\bar{\partial} F \cdot N|_\Gamma \in H^{s-\frac{3}{2}}(\Gamma)$ for $1 \leq s \leq r$, then there exists a constant $\bar{C} > 0$ depending only on Ω such that*

$$\begin{aligned} \|F\|_s &\leq \bar{C} \left(\|F\|_0 + \|\operatorname{curl} F\|_{s-1} + \|\operatorname{div} F\|_{s-1} + |\bar{\partial} F \cdot N|_{s-\frac{3}{2}} \right), \\ \|F\|_s &\leq \bar{C} \left(\|F\|_0 + \|\operatorname{curl} F\|_{s-1} + \|\operatorname{div} F\|_{s-1} + \sum_{\alpha=1}^2 |\bar{\partial} F \cdot T_\alpha|_{s-\frac{3}{2}} \right), \end{aligned} \quad (11)$$

where N denotes the outward unit-normal to Γ , and T_α , $\alpha = 1, 2$, denotes the two tangent vectors to Γ .

The first estimate is well-known and follows from the identity $-\Delta F = \operatorname{curl} \operatorname{curl} F - \operatorname{Ddiv} F$; a convenient reference is Taylor [9]. The second estimate immediately follows.

5. Horizontal convolution-by-layers and commutation estimates.

5.1. Horizontal convolution-by-layers. With $x_h = (x_1, x_2)$, we define $\rho \in C_0^\infty(\mathbb{R}^2)$ by $\rho(x_h) = C \exp\left(\frac{1}{|x_h|^2 - 1}\right)$ if $|x_h| < 1$ and $\rho(x_h) = 0$ if $|x_h| \geq 1$; we then select the constant C so that $\int_{\mathbb{R}^2} \rho dx_h = 1$. We define $\rho_\kappa(x_h) = \frac{1}{\kappa^2} \rho\left(\frac{x}{\kappa}\right)$. It follows that for $\kappa > 0$, $0 \leq \rho_\kappa \in C_0^\infty(\mathbb{R}^2)$ with $\operatorname{spt}(\rho_\kappa) \subset \overline{B(0, \kappa)}$. (Here, spt stands for support.) We define the operation of *horizontal convolution-by-layers* as follows:

$$\Lambda_\kappa f(x_h, x_3) = \int_{\mathbb{R}^2} \rho_\kappa(x_h - y_h) f(y_h, x_3) dy_h \quad \text{for } f \in L^1(\mathbb{R}^2).$$

We introduced this type of smoothing operation in Section 2 of [3], but in the setting of a general initial domain Ω , which required a partition of unity and a collection of local charts to define properly.

By standard properties of convolution, there exists a constant C which is independent of κ , such that for $s \geq 0$,

$$\|\Lambda_\kappa F\|_s \leq C \|F\|_s \quad \forall F \in H^s(\Omega),$$

and

$$|\Lambda_\kappa F|_s \leq C|F|_s \quad \forall F \in H^s(\Gamma).$$

Furthermore,

$$\kappa \|\bar{\partial} \Lambda_\kappa F\|_0 \leq C \|F\|_0 \quad \forall F \in L^2(\Omega). \quad (12)$$

5.2. Commutation estimates.

Lemma 5.1. *For $f \in W^{1,\infty}(\Gamma)$ and $g, \bar{\partial}g \in L^2(\Gamma)$, there is a generic constant C independent of κ such that*

$$|\Lambda_\kappa(f \bar{\partial}g) - f \Lambda_\kappa \bar{\partial}g|_0 \leq C |f|_{W^{1,\infty}(\Gamma)} |g|_0,$$

where $W^{1,\infty}(\Gamma)$ denotes the Sobolev space of functions $u \in L^\infty(\Gamma)$ with weak derivative $\bar{\partial}u \in L^\infty(\Gamma)$.

Proof. We have that

$$\begin{aligned} [\Lambda_\kappa(f \bar{\partial}g) - f \Lambda_\kappa \bar{\partial}g](x) &= \int_{B(x_h, \kappa)} \frac{1}{\kappa^2} \rho\left(\frac{x_h - y_h}{\kappa}\right) [f(y_h) - f(x_h)] \bar{\partial}g(y_h) dy_h \\ &= \underbrace{\int_{B(x_h, \kappa)} \frac{1}{\kappa^3} \bar{\partial}\rho\left(\frac{x_h - y_h}{\kappa}\right) [f(y_h) - f(x_h)] g(y_h) dy_h}_{\mathcal{I}_1} \\ &\quad - \underbrace{\int_{B(x_h, \kappa)} \rho_\kappa(x_h - y_h) \bar{\partial}f(y_h) g(y_h) dy_h}_{\mathcal{I}_2}, \end{aligned}$$

where we have used integration-by-parts in order to obtain the second equality. From Morrey's inequality, for all $y_h \in B(x_h, \kappa)$,

$$|f(x_h) - f(y_h)| \leq \kappa \|\bar{\partial}f\|_{L^\infty(\mathbb{T}^2)} \leq \kappa |f|_{W^{1,\infty}(\Gamma)},$$

so that with $\mathcal{K}(x_h) = \frac{1}{\kappa^2} |\bar{\partial}\rho(\frac{x_h}{\kappa})|$,

$$|\mathcal{I}_1| \leq |f|_{W^{1,\infty}(\Gamma)} \mathcal{K} * |g|,$$

so by Young's inequality for convolution integrals,

$$|\mathcal{I}_1|_0 \leq |f|_{W^{1,\infty}(\Gamma)} \|\mathcal{K}\|_{L^1(\mathbb{T}^2)} |g|_0.$$

Similarly,

$$|\mathcal{I}_2|_0 \leq |f|_{W^{1,\infty}(\Gamma)} \|\rho_\kappa\|_{L^1(\mathbb{T}^2)} |g|_0.$$

The assertion is proved, given that

$$\|\mathcal{K}\|_{L^1(\mathbb{T}^2)} = \int_{\mathbb{T}^2} \frac{1}{\kappa^2} |\bar{\partial}\rho(\frac{y_h}{\kappa})| dy_h = \int_{\mathbb{T}^2} |\bar{\partial}\rho(z)| dz < \infty.$$

□

Lemma 5.2. *For $\kappa > 0$, there exists $C > 0$ independent of κ , such that for any $g \in H^{\frac{1}{2}}(\Omega)$ and $f \in H^3(\Omega)$, we have that*

$$\|\Lambda_\kappa(fg) - f \Lambda_\kappa g\|_{\frac{1}{2}} \leq C \kappa \|g\|_{\frac{1}{2}} \|f\|_3 + C \kappa^{\frac{1}{2}} \|g\|_0 \|f\|_3.$$

Proof. Let $\Delta = \Lambda_\kappa(fg) - f \Lambda_\kappa g$. Then, we have that

$$\Delta(x) = \int_{B(x_h, \kappa)} \rho_\kappa(x_h - y_h) [f(y_h, x_3) - f(x_h, x_3)] g(y_h, x_3) dy_h.$$

Using the fact that $H^2(\Omega)$ is embedded in $L^\infty(\Omega)$, we have that

$$|\Delta(x)| \leq C\kappa \|f\|_3 \int_{B(x_h, \kappa)} \rho_\kappa(x_h - y_h) |g(y_h, x_3)| dy_h,$$

showing that

$$\begin{aligned} \|\Delta\|_0 &\leq C\kappa \|f\|_3 \|\Lambda_\kappa g\|_0 \\ &\leq C\kappa \|f\|_3 \|g\|_0. \end{aligned} \tag{13}$$

Now, for $j = 1, 2, 3$,

$$\Delta_{,j} = \Lambda_\kappa(fg_{,j}) - f \Lambda_\kappa g_{,j} + \Lambda_\kappa(f_{,j}g) - f_{,j} \Lambda_\kappa g.$$

The difference between the two first terms of the right-hand side of this identity can be treated in a similar fashion as (13), leading us to:

$$\begin{aligned} \|\Delta_{,j}\|_0 &\leq C\kappa \|f\|_3 \|g\|_1 + \|\Lambda_\kappa(f_{,j}g)\|_0 + \|f_{,j} \Lambda_\kappa g\|_0 \\ &\leq C\kappa \|f\|_3 \|g\|_1 + \|f_{,j}g\|_0 + \|f_{,j}\|_{L^\infty(\Omega)} \|\Lambda_\kappa g\|_0 \\ &\leq C\kappa \|f\|_3 \|g\|_1 + 2\|f_{,j}\|_{L^\infty(\Omega)} \|g\|_0 \\ &\leq C\kappa \|f\|_3 \|g\|_1 + C\|f\|_3 \|g\|_0. \end{aligned} \tag{14}$$

Consequently, we obtain by interpolation from (13) and (14):

$$\|\Delta\|_{\frac{1}{2}} \leq C\kappa \|f\|_3 \|g\|_{\frac{1}{2}} + C\kappa^{\frac{1}{2}} \|f\|_3 \|g\|_0.$$

□

Following the estimate (14), we can similarly obtain estimates for $\|\Delta_{,jk}\|_0$ and $\|\Delta_{,jkl}\|_0$, $j, k, l = 1, 2, 3$, and thus estimate the $H^2(\Omega)$ -norm as well as the $H^3(\Omega)$ -norm of Δ . Interpolation then yields the following:

Lemma 5.3. *For $\kappa > 0$ and $s = \frac{3}{2}$ or $\frac{5}{2}$, there exists $C > 0$ independent of κ , such that for any $g \in H^s(\Omega)$ and $f \in H^3(\Omega)$, we have that*

$$\|\Lambda_\kappa(fg) - f \Lambda_\kappa g\|_s \leq C\kappa \|g\|_s \|f\|_3 + C\kappa^{\frac{1}{2}} \|g\|_{s-\frac{1}{2}} \|f\|_3.$$

6. An asymptotically consistent κ -approximation of the Euler equations.

6.1. The smoothed flow map η_κ .

Definition 6.1 (The horizontally-smoothed flow map η_κ). Suppose the Lagrangian flow map η is in $C([0, T], H^3(\Omega))$. For $\kappa \in (0, \kappa_0)$, we set on $[0, T]$

$$\eta_\kappa = \Lambda_\kappa^2 \eta \quad \text{and} \quad v_\kappa = \partial_t \eta_\kappa.$$

By choosing $T > 0$ and κ_0 sufficiently small, we can ensure that

$$A_\kappa(x, t) = [D\eta_\kappa(x, t)]^{-1}$$

is well-defined on $[0, T]$. We then define $J_\kappa = \det[D\eta_\kappa]$, and $a_\kappa = J_\kappa A_\kappa$.

6.2. Smoothing the initial data. In order to construct solutions to (3), we will introduce an approximation scheme below, and it will be convenient to smooth the initial velocity field.

Let $0 \leq \varrho_\kappa \in C_0^\infty(\mathbb{R}^3)$ denote the standard family of mollifiers with $\text{spt}(\varrho_\kappa) \subset \overline{B(0, \kappa)}$, and let \mathcal{E}_Ω denote the Sobolev extension operator mapping $H^s(\Omega)$ to $H^s(\mathbb{R}^3)$ for $s \geq 0$.

We set $w_0^\kappa = \varrho_\kappa \mathcal{E}_\Omega(u_0)$, and define

$$u_0^\kappa = w_0^\kappa - Dr^\kappa,$$

where the scalar function r^κ is the solution to the elliptic problem

$$\Delta r^\kappa = \text{div } w_0^\kappa \quad \text{in } \Omega, \quad (15a)$$

$$x_h \mapsto r^\kappa(x_h, x_3) \text{ periodic} \quad x_3 \in [0, 1], \quad (15b)$$

$$r^\kappa = 0 \quad \text{on } \Gamma, \quad (15c)$$

$$\frac{\partial r^\kappa}{\partial x_3} = (w_0^\kappa)^3 \quad \text{on } \{x_3 = 0\}. \quad (15d)$$

Thus, given $u_0 \in H_{\text{div}}^4(\Omega)$, it follows that $u_0^\kappa \in H_{\text{div}}^s(\Omega)$ for all $s \geq 4$, and that $u_0^\kappa \rightarrow u_0$ in $H^4(\Omega)$ as $\kappa \rightarrow 0$.

6.3. The approximate κ -problem. For $\kappa \in (0, \kappa_0)$, we consider the following sequence of approximate problems:

$$v_t + A_\kappa^T Dq = 0 \quad \text{in } \Omega \times (0, T_\kappa], \quad (16a)$$

$$\text{div}_{\eta_\kappa} v = 0 \quad \text{in } \Omega \times (0, T_\kappa], \quad (16b)$$

$$q = 0 \quad \text{on } \Gamma \times (0, T_\kappa], \quad (16c)$$

$$v^3 = 0 \quad \text{on } \{x_3 = 0\} \times (0, T_\kappa], \quad (16d)$$

$$(v, \eta) = (u_0^\kappa, e_\kappa) \quad \text{on } \Omega \times \{t = 0\}, \quad (16e)$$

where the solution $\eta = \eta(\kappa)$ depends on the parameter κ , but for notational simplification, we do not explicitly write this dependence, and where $\text{div}_{\eta_\kappa} v = (A_\kappa)_i^j v^i_{,j}$, and $e_\kappa = \Lambda_\kappa^2 e$, where $e(x) = x$ denotes the identity map on Ω . We refer to the approximation (16) as the κ -problem. (Note that our solution is periodic in the x_1 and x_2 directions.)

7. Construction of smooth solutions to κ -approximate Euler equations.

Definition 7.1 (The manifold of volume-preserving embeddings). For $s \geq 3$, we let

$$\mathcal{D}_{\text{vol}}^s = \{ \eta \in H^s(\Omega; \mathbb{R}^3) : \eta(\{x_3 = 0\}) \subset \{x_3 = 0\}, x_h \mapsto \eta(x_h, x_3) \text{ periodic} \\ \det D\eta = 1, \eta^{-1} \in H^s(\eta(\Omega); \Omega) \}.$$

For $s \geq 3$, $\mathcal{D}_{\text{vol}}^s$ is a infinite-dimensional C^∞ Hilbert manifold (see, for example, [4]). We let $T\mathcal{D}_{\text{vol}}^s$ denote the tangent bundle over $\mathcal{D}_{\text{vol}}^s$, and $TT\mathcal{D}_{\text{vol}}^s$ the second tangent bundle. Locally, elements of $T\mathcal{D}_{\text{vol}}^s$ consist of pairs $(\eta, v) \in \mathcal{D}_{\text{vol}}^s \times H_{\text{div}}^s(\Omega) \circ \eta$, where

$$H_{\text{div}}^s(\Omega) \circ \eta = \{ u \circ \eta : u \in H_{\text{div}}^s(\Omega) \}.$$

Theorem 7.2. Suppose that for $s \geq 4$, $u_0^\kappa \in H^s(\Omega)$ with $\text{div } u_0 = 0$. Then for each $\kappa \in (0, \kappa_0)$, there exists a unique solution $(\eta(\kappa), v(\kappa)) \in C^\infty([0, T_\kappa]; T\mathcal{D}_{\text{vol}}^s)$ to (16) with $T_\kappa > 0$ depending on $\|u_0^\kappa\|_s$ and on $\kappa > 0$.

Proof. With η_κ and v_κ defined in Definition 6.1, we define

$$u = v \circ \eta_\kappa^{-1}, \quad u_\kappa = v_\kappa \circ \eta_\kappa^{-1}, \quad \text{and } p = q \circ \eta_\kappa^{-1}.$$

It follows from the chain-rule that (16) can be written on the time dependent domain $\eta_\kappa(t, \Omega)$ as

$$u_t + u_\kappa \cdot Du + Dp = 0 \quad \text{in } \eta_\kappa(t, \Omega), \quad (17a)$$

$$\operatorname{div} u = 0 \quad \text{in } \eta_\kappa(t, \Omega), \quad (17b)$$

$$p = 0 \quad \text{on } \eta_\kappa(t, \Gamma), \quad (17c)$$

$$(u, \eta) = (u_0^\kappa, e) \quad \text{on } \Omega \times \{t = 0\}, \quad (17d)$$

Taking the divergence of (17a), we see that $p(t)$ satisfies

$$-\Delta p = u^i_{,j} u^j_{\kappa,i} \quad \text{in } \eta_\kappa(t, \Omega), \quad (18a)$$

$$p = 0 \quad \text{on } \eta_\kappa(t, \Gamma), \quad (18b)$$

$$\frac{\partial p}{\partial N} = 0 \quad \text{on } \{x_3 = 0\}. \quad (18c)$$

The estimates for $p(t)$ can be easily obtained by transforming (18), set on the smoothed moving domain $\eta_\kappa(t, \Omega)$, to an elliptic equation on the fixed domain Ω . It is important to note that this transformation should not be made with the map $\eta_\kappa(t)$, but rather with a family of diffeomorphisms which inherits the smoothness of $\eta_\kappa|_\Gamma$. To this end, consider the solution to $\Delta\Phi(t) = 0$ in Ω with $\Phi(t) = \eta_\kappa(t)$ on Γ . It follows that

$$\|\Phi(t)\|_{s+1} \leq C|\eta_\kappa|_{s+1/2} \leq C(1 + \frac{1}{\kappa})|\eta|_{s-1/2} \leq C(1 + \frac{1}{\kappa})\|\eta\|_s, \quad (19)$$

where the first inequality is the standard elliptic estimate for the Dirichlet problem, the second follows from (12), and the third from the trace theorem. For κ and T_κ taken sufficiently small $\|\Phi(t) - e\|_s$ can be made arbitrarily small on $[0, T_\kappa]$, from which it follows that each such $\Phi(t) : \Omega \rightarrow \eta_\kappa(t, \Omega)$ is a diffeomorphism.

Next, define the matrix $B = [D\Phi]^{-1}$ and the pressure function $Q = p \circ \Phi$. Using the chain-rule, (18) is transformed to

$$-B^j_i [B^k_i Q_{,k}]_{,j} = (u^i_{,j} u^j_{\kappa,i}) \circ \Phi \quad \text{in } \Omega, \quad (20a)$$

$$Q = 0 \quad \text{on } \Gamma, \quad (20b)$$

$$Q_{,k} B^k_i B^3_i = 0 \quad \text{on } \{x_3\} = 0, \quad (20c)$$

for $0 \leq t \leq T_\kappa$. Elliptic estimates (in conjunction with the Sobolev embedding theorem) show that

$$\|Q(t)\|_{s+1} \leq CP(\|\Phi(t)\|_{s+1})\|u^i_{,j} u^j_{\kappa,i} \circ \Phi\|_{s-1}$$

where P is a polynomial function of its argument; hence, together with (19), we have the estimate

$$\|Q(t)\|_{s+1} \leq C_\kappa P(\|\eta(t)\|_s) \cdot \|v\|_s^2,$$

where the constant C_κ depends on $\kappa > 0$ and, in fact, blows-up as $\kappa \rightarrow 0$. Since $p = Q \circ \Phi^{-1}$, we see that

$$\|Dp(t) \circ \eta(t)\|_s \leq C_\kappa P(\|\eta(t)\|_s, \|v\|_s). \quad (21)$$

We define the function \mathcal{F} on $T\mathcal{D}_{\text{vol}}^s$ by

$$\mathcal{F}(\eta, v) = -Dp \circ \eta_\kappa,$$

and write (16a) as the coupled system of first-order ordinary differential equations written on $T\mathcal{D}_{\text{vol}}^s$:

$$\begin{aligned}\partial_t(\eta, v) &= (v, \mathcal{F}), \\ (\eta, v)|_{t=0} &= (e, u_0).\end{aligned}$$

According to the estimate (21), $(v, \mathcal{F}) : T\mathcal{D}_{\text{vol}}^s \rightarrow T\mathcal{D}_{\text{vol}}^s$ continuously; moreover, since composition on the right is a smooth operation,

$$(v, \mathcal{F}) : T\mathcal{D}_{\text{vol}}^s \rightarrow T\mathcal{D}_{\text{vol}}^s \text{ is a } C^\infty \text{ map.}$$

Thus, the fundamental theorem of ordinary differential equations (or Picard iteration) shows that there exists a time $T_\kappa > 0$ depending on the initial data (and, of course, $\kappa > 0$) such that

$$(\eta, v) = (\eta(\kappa), v(\kappa)) \in C^\infty([0, T_\kappa]; T\mathcal{D}_{\text{vol}}^s)$$

is a unique solution of (16). Since, by definition $Q = 0$ on Γ , and

$$q = Q \circ \Phi^{-1} \circ \eta_\kappa, \quad (22)$$

it follows that $q = 0$ on Γ as well. \square

8. Asymptotic estimates which are independent of the smooth parameter κ . According to Theorem 7.2, we have unique solutions to (16),

$$(\eta(\kappa), v(\kappa)) \in C^\infty([0, T_\kappa]; T\mathcal{D}_{\text{vol}}^s),$$

and $q(\kappa)$ given by (22). We will take $s \geq 6$, and for notational convenience we will denote $(\eta(\kappa), v(\kappa))$ by $(\tilde{\eta}, \tilde{v})$, and write \tilde{A} for $[D\tilde{\eta}]^{-1}$. We use the notation $\tilde{\eta}_\kappa$ to denote $\Lambda_\kappa^2 \tilde{\eta}$ and set $\tilde{A}_\kappa = [D\tilde{\eta}_\kappa]^{-1}$.

8.1. A continuous-in-time energy function appropriate for the asymptotic process $\kappa \rightarrow 0$.

Definition 8.1. We set on $[0, T_\kappa]$

$$E_\kappa(t) = 1 + \|\Lambda_\kappa \tilde{\eta}(t)\|_{4.5}^2 + \|\tilde{v}(t)\|_4^2 + \|\kappa \tilde{v}(t)\|_{4.5}^2 + \|\tilde{v}_t(t)\|_{3.5}^2 + \|\sqrt{\kappa} \tilde{v}_t(t)\|_4^2. \quad (23)$$

The function $E_\kappa(t)$ is the *higher-order energy function* which we will prove remains bounded on a time-interval which is independent of κ . Given $(\tilde{\eta}, \tilde{v}) \in T\mathcal{D}_{\text{vol}}^6$, the $E_\kappa(t)$ is continuous on $[0, T_\kappa]$.

Definition 8.2. We set $E(t) = E_{\kappa=0}(t)$.

Definition 8.3. We set the constant M_0 to be a polynomial function of $E(0)$ so that

$$M_0 = P(E(0), \|\text{curl } u_0^\kappa\|_{3.5}^2). \quad (24)$$

8.2. Statement of the main result. Given an initial velocity field $u_0 \in H_{\text{div}}^4$, we obtain the initial pressure function p_0 as the solution to the elliptic equation

$$\begin{aligned}-\Delta p_0 &= u_{0,j}^i u_{0,i}^j \text{ in } \Omega, \\ p_0 &= 0 \quad \text{on } \Gamma, \\ p_{0,3} &= 0 \quad \text{on } \{x_3 = 0\}.\end{aligned}$$

Theorem 8.4 (Main Result). *Given initial data $u_0 \in H_{\text{div}}^4(\Omega)$ with $\text{curl } u_0 \in H^{3.5}(\Omega)$ such that*

$$-\frac{\partial p_0}{\partial x_3}(x) \geq \lambda > 0 \quad \text{for } x \in \Gamma,$$

there exists a solution to (3) verifying

$$\sup_{t \in [0, T]} E(t) \leq P(E(0)).$$

If $u_0 \in H_{\text{div}}^5(\Omega)$ and $\text{curl } u_0 \in H^{4.5}(\Omega)$, then the solution is unique.

Remark 1. The same theorem and proof hold in the case that $\Omega \subset \mathbb{R}^2$. We refer the reader to our paper [3] for the case of a general initial domain Ω .

Remark 2. The regularity for the existence theory is not optimal. In fact, for our domain Ω , all that is necessary to establish existence and uniqueness of solutions to (3) is an initial velocity field $u_0 \in H_{\text{div}}^3(\Omega)$; nevertheless, the assumptions of Theorem 8.4 allow for the most transparent proof.

8.3. Conventions about constants. As noted above, Theorem 7.2 provides us with solutions $(\tilde{\eta}, \tilde{v}) \in T\mathcal{D}_{\text{vol}}^6$, and hence $\sup_{t \in [0, T_\kappa]} E_\kappa(t)$ is continuous.

We take $T_\kappa > 0$ sufficiently small so that, using the fundamental theorem of calculus, for constants c_1, c_2 and $t \in [0, T_\kappa]$,

$$\begin{aligned} -\tilde{q}_{,3}(t) &\geq \lambda/2, \\ c_1 \det \tilde{g}_\kappa(0) &\leq \det \tilde{g}_\kappa(t) \leq c_2 \det \tilde{g}_\kappa(0) \text{ on } \Gamma, \\ c_1 \det \tilde{J}_\kappa(0) &\leq \det \tilde{J}_\kappa(t) \leq c_2 \det \tilde{J}_\kappa(0) \text{ in } \Omega, \\ \|\tilde{\Lambda}_\kappa \eta(t)\|_4 &\leq \|\Lambda_\kappa e\|_4 + 1, \quad \|\tilde{q}(t)\|_4 \leq \|\tilde{q}(0)\|_4 + 1, \\ \|\tilde{v}(t)\|_{3.5} &\leq \|u_0\|_{3.5} + 1, \quad \|\tilde{v}_t(t)\|_3 \leq \|\tilde{v}_t(0)\|_3 + 1. \end{aligned}$$

The right-hand sides appearing in the last three inequalities shall be denoted by a generic constant C in the estimates that we will perform.

8.4. Curl and divergence estimates for $\tilde{\eta}$, \tilde{v} , and \tilde{v}_t .

Proposition 2. *For all $t \in (0, T)$, with $T \leq T_\kappa$,*

$$\begin{aligned} \|\text{curl } \Lambda_\kappa \tilde{\eta}(t)\|_{3.5}^2 + \|\text{curl } \tilde{v}(t)\|_3^2 + \|\kappa \text{curl } \tilde{v}(t)\|_{3.5}^2 + \|\sqrt{\kappa} \text{curl } \tilde{v}_t(t)\|_{2.5}^2 \\ \leq M_0 + C T P\left(\sup_{t \in [0, T]} E_\kappa(t)\right). \end{aligned} \quad (25)$$

Proof. By taking the curl of (16a), we have that

$$\text{curl}_{\eta_\kappa} \tilde{v}_t = 0.$$

From (4), it follows that $(\text{curl}_{\tilde{\eta}_\kappa} \tilde{v})_t^k = B(\tilde{A}_\kappa, D\tilde{v})$, where

$$B(\tilde{A}_\kappa, D\tilde{v}) = \varepsilon_{kji} (\tilde{A}_\kappa)_{tj}^s \tilde{v}^i_{,s} = \varepsilon_{kij} \tilde{v}^i_{,s} (\tilde{A}_\kappa)_p^s \tilde{v}_\kappa^p_{,l} (\tilde{A}_\kappa)_j^l;$$

hence,

$$\text{curl}_{\tilde{\eta}_\kappa} \tilde{v}(t) = \text{curl } u_0^\kappa + \int_0^t B(\tilde{A}_\kappa(t'), D\tilde{v}(t')) dt'. \quad (26)$$

Step 1. Estimate for $\text{curl } \Lambda_\kappa \tilde{\eta}$. Computing the gradient of (26) yields

$$\text{curl}_{\tilde{\eta}_\kappa} D\tilde{v}(t) = D \text{curl } u_0^\kappa - \varepsilon_{.ji} D(\tilde{A}_\kappa)_j^s \tilde{v}^i_{,s} + \int_0^t DB(\tilde{A}_\kappa(t'), D\tilde{v}(t')) dt'.$$

Applying the fundamental theorem of calculus once again, shows that

$$\begin{aligned} \operatorname{curl}_{\tilde{\eta}_\kappa} D\tilde{\eta}(t) &= tD \operatorname{curl} u_0^\kappa + \varepsilon_{.ji} \int_0^t [(\tilde{A}_\kappa)_{t_j}^s D\tilde{\eta}^i_{,s} - D(\tilde{A}_\kappa)_j^s \tilde{v}^i_{,s}] dt' \\ &\quad + \int_0^t \int_0^{t'} DB(\tilde{A}_\kappa(t''), D\tilde{v}(t'')) dt'' dt', \end{aligned}$$

and finally that

$$\begin{aligned} D \operatorname{curl} \tilde{\eta}(t) &= tD \operatorname{curl} u_0^\kappa - \varepsilon_{.ji} \int_0^t (\tilde{A}_\kappa)_{t_j}^s (t') dt' D\tilde{\eta}^i_{,s} \\ &\quad + \varepsilon_{.ji} \int_0^t [(\tilde{A}_\kappa)_{t_j}^s D\tilde{\eta}^i_{,s} - D(\tilde{A}_\kappa)_j^s \tilde{v}^i_{,s}] dt' + \int_0^t \int_0^{t'} DB(\tilde{A}_\kappa(t''), D\tilde{v}(t'')) dt'' dt'. \end{aligned} \quad (27)$$

Using the fact that $\partial_t(\tilde{A}_\kappa)_j^s = -(\tilde{A}_\kappa)_l^s \tilde{v}_{\kappa,p}^l (\tilde{A}_\kappa)_j^p$ and $D(\tilde{A}_\kappa)_j^s = -(\tilde{A}_\kappa)_l^s D\tilde{\eta}_{\kappa,p}^l (\tilde{A}_\kappa)_j^p$, we see that

$$\begin{aligned} DB(\tilde{A}, D\tilde{v}) &= -\varepsilon_{kji} [D\tilde{v}^i_{,s} (\tilde{A}_\kappa)_l^s \tilde{v}_{\kappa,p}^l (\tilde{A}_\kappa)_j^p + \tilde{v}^i_{,s} (\tilde{A}_\kappa)_l^s D\tilde{v}_{\kappa,p}^l (\tilde{A}_\kappa)_j^p \\ &\quad + \tilde{v}^i_{,s} \tilde{v}_{\kappa,p}^l D((\tilde{A}_\kappa)_l^s (\tilde{A}_\kappa)_j^p)]. \end{aligned}$$

The precise structure of the right-hand side is not very important; rather, the derivative count is the focus, and as such we write

$$DB(\tilde{A}, D\tilde{v}) \sim D^2 \tilde{v} D\tilde{v}_\kappa \tilde{A}_\kappa \tilde{A}_\kappa + D^2 \tilde{v}_\kappa D\tilde{v} \tilde{A}_\kappa \tilde{A}_\kappa + D^2 \tilde{\eta}_\kappa D\tilde{v} D\tilde{v}_\kappa \tilde{A}_\kappa \tilde{A}_\kappa.$$

Integrating by parts in time in the last term of the right-hand side of (27), we see that

$$\begin{aligned} \int_0^t \int_0^{t'} DB(\tilde{A}_\kappa, D\tilde{v}) dt'' dt' &\sim - \int_0^t \int_0^{t'} \left[D^2 \tilde{\eta} (D\tilde{v}_\kappa \tilde{A}_\kappa \tilde{A}_\kappa)_t + D^2 \tilde{\eta}_\kappa (D\tilde{v} \tilde{A}_\kappa \tilde{A}_\kappa)_t \right] dt'' dt' \\ &\quad + \int_0^t \int_0^{t'} D^2 \tilde{\eta}_\kappa D\tilde{v} D\tilde{v}_\kappa \tilde{A}_\kappa \tilde{A}_\kappa dt'' dt' \\ &\quad + \int_0^t \left[D^2 \tilde{\eta} D\tilde{v}_\kappa \tilde{A}_\kappa \tilde{A}_\kappa + D^2 \tilde{\eta}_\kappa D\tilde{v} \tilde{A}_\kappa \tilde{A}_\kappa \right] dt'. \end{aligned}$$

Thus, we can write

$$\begin{aligned} D \operatorname{curl} \tilde{\eta}(t) &\sim tD \operatorname{curl} u_0^\kappa + \underbrace{D^2 \tilde{\eta} \int_0^t D\tilde{v}_\kappa \tilde{A}_\kappa \tilde{A}_\kappa dt'}_{\mathcal{I}_1} + \underbrace{\int_0^t D^2 \tilde{\eta} D\tilde{v}_\kappa \tilde{A}_\kappa \tilde{A}_\kappa dt'}_{\mathcal{I}_2} \\ &\quad + \underbrace{\int_0^t D^2 \tilde{\eta}_\kappa D\tilde{v} \tilde{A}_\kappa \tilde{A}_\kappa dt'}_{\mathcal{I}_3} + \underbrace{\int_0^t \int_0^{t'} D^2 \tilde{\eta}_\kappa D\tilde{v} D\tilde{v}_\kappa \tilde{A}_\kappa \tilde{A}_\kappa dt'' dt'}_{\mathcal{I}_4} \\ &\quad + \underbrace{\int_0^t \int_0^{t'} D^2 \tilde{\eta}_\kappa (D\tilde{v} \tilde{A}_\kappa \tilde{A}_\kappa)_t dt'' dt'}_{\mathcal{I}_5} + \underbrace{\int_0^t \int_0^{t'} D^2 \tilde{\eta} (D\tilde{v}_\kappa \tilde{A}_\kappa \tilde{A}_\kappa)_t dt'' dt'}_{\mathcal{I}_6}. \end{aligned}$$

Our goal is to estimate $\|D \operatorname{curl} \Lambda_\kappa \eta\|_{2.5}^2$, which in turn requires us to estimate $\|\Lambda_\kappa \mathcal{I}_i\|_{2.5}^2$ for $i = 1, \dots, 6$. We begin with $i = 1$:

$$\begin{aligned} \|\Lambda_\kappa \mathcal{I}_1\|_{2.5}^2 &\leq \|D^2 \Lambda_\kappa \tilde{\eta} \int_0^t D \tilde{v}_\kappa \tilde{A}_\kappa \tilde{A}_\kappa dt'\|_{2.5}^2 \\ &\quad + \|\Lambda_\kappa \left(D^2 \tilde{\eta} \int_0^t D \tilde{v}_\kappa \tilde{A}_\kappa \tilde{A}_\kappa dt' \right) - \int_0^t D \tilde{v}_\kappa \tilde{A}_\kappa \tilde{A}_\kappa dt' \Lambda_\kappa D^2 \tilde{\eta} \|_{2.5}^2 \end{aligned}$$

It is easy to see that

$$\|D^2 \Lambda_\kappa \tilde{\eta} \int_0^t D \tilde{v}_\kappa \tilde{A}_\kappa \tilde{A}_\kappa dt'\|_{2.5}^2 \leq CTP \left(\sup_{t \in [0, T]} E_\kappa(t) \right),$$

and by Lemma 5.3

$$\begin{aligned} \|\Lambda_\kappa \left(D^2 \tilde{\eta} \int_0^t D \tilde{v}_\kappa \tilde{A}_\kappa \tilde{A}_\kappa dt' \right) - \int_0^t D \tilde{v}_\kappa \tilde{A}_\kappa \tilde{A}_\kappa dt' \Lambda_\kappa D^2 \tilde{\eta} \|_{2.5}^2 \\ \leq C\kappa^2 \|D^2 \tilde{\eta}\|_{2.5}^2 \left\| \int_0^t D \tilde{v}_\kappa \tilde{A}_\kappa \tilde{A}_\kappa dt' \right\|_3^2 + C\kappa \|D^2 \tilde{\eta}\|_2^2 \left\| \int_0^t D \tilde{v}_\kappa \tilde{A}_\kappa \tilde{A}_\kappa dt' \right\|_3^2 \\ \leq CTP \left(\sup_{t \in [0, T]} E_\kappa(t) \right) \end{aligned}$$

The same type of commutation estimate shows that

$$\|\Lambda_\kappa \mathcal{I}_2\|_{2.5}^2 \leq CTP \left(\sup_{t \in [0, T]} E_\kappa(t) \right).$$

For $i = 3, 4, 5$, we use that $\|\Lambda_\kappa \mathcal{I}_i\|_{2.5} \leq \|\mathcal{I}_i\|_{2.5}$ and as $H^{2.5}(\Omega)$ is a multiplicative algebra, we see that for $i = 3, 4, 5$,

$$\|\Lambda_\kappa \mathcal{I}_i\|_{2.5}^2 \leq CTP \left(\sup_{t \in [0, T]} E_\kappa(t) \right).$$

Finally, we consider the case that $i = 6$:

$$\begin{aligned} \|\Lambda_\kappa \mathcal{I}_6\|_{2.5}^2 &\leq \left\| \int_0^t \int_0^{t'} D^2 \Lambda_\kappa \tilde{\eta} (D \tilde{v}_\kappa \tilde{A}_\kappa \tilde{A}_\kappa)_t dt'' dt' \right\|_{2.5}^2 \\ &\quad + \left\| \int_0^t \int_0^{t'} \left(\Lambda_\kappa \left[D^2 \tilde{\eta} (D \tilde{v}_\kappa \tilde{A}_\kappa \tilde{A}_\kappa)_t \right] - (D \tilde{v}_\kappa \tilde{A}_\kappa \tilde{A}_\kappa)_t D^2 \Lambda_\kappa \tilde{\eta} \right) dt'' dt' \right\|_{2.5}^2 \\ &\leq CTP \left(\sup_{t \in [0, T]} E_\kappa(t) \right), \end{aligned}$$

where we have used Lemma 5.3 for the last inequality together with the fact that $\|\kappa^{\frac{1}{2}} \tilde{v}_t\|_4^2$ is contained in the energy function $E_\kappa(t)$. Therefore, we have proven that

$$\|D \operatorname{curl} \Lambda_\kappa \eta\|_{2.5}^2 \leq M_0 + CTP \left(\sup_{t \in [0, T]} E_\kappa(t) \right),$$

and hence with $\operatorname{curl}_{\tilde{\eta}_\kappa} \tilde{v}_t = 0$, that

$$\|\operatorname{curl} \Lambda_\kappa \eta\|_{3.5}^2 \leq M_0 + CTP \left(\sup_{t \in [0, T]} E_\kappa(t) \right).$$

Step 2. Estimate for $\operatorname{curl} \tilde{v}$. From (26),

$$\operatorname{curl} v(t) = \operatorname{curl} u_0^\kappa + \int_0^t B(\tilde{A}_\kappa(t'), D \tilde{v}(t')) dt' - \varepsilon_{jk} \tilde{v}^k{}_{,r} \int_0^t \tilde{\partial}_t (A_\kappa)_j^r dt'.$$

As $H^3(\Omega)$ is a multiplicative algebra, it follows that on $[0, T]$,

$$\|\operatorname{curl} \tilde{v}(t)\|_3^2 \leq M_0 + CTP \left(\sup_{t \in [0, T]} E_\kappa(t) \right).$$

Similarly,

$$\|\kappa \operatorname{curl} \tilde{v}(t)\|_{3.5}^2 \leq M_0 + CTP\left(\sup_{t \in [0, T]} E_\kappa(t)\right).$$

Step 3. Estimate for $\operatorname{curl} \tilde{v}_t$. From (26),

$$\operatorname{curl} \tilde{v}_t(t) = \varepsilon_{ikj} \int_0^t \partial_t (\tilde{A}_\kappa)_j^r(t') dt' \tilde{v}_{t,r}^i,$$

from which it follows that on $[0, T]$,

$$\|\sqrt{\kappa} \operatorname{curl} \tilde{v}_t(t)\|_3^2 \leq CTP\left(\sup_{t \in [0, T]} E_\kappa(t)\right).$$

□

Proposition 3. For all $t \in (0, T)$, with $T \leq T_\kappa$,

$$\begin{aligned} \|\operatorname{div} \Lambda_\kappa \tilde{\eta}(t)\|_{3.5}^2 + \|\operatorname{div} \tilde{v}(t)\|_3^2 + \|\kappa \operatorname{div} \tilde{v}(t)\|_{3.5}^2 + \|\sqrt{\kappa} \operatorname{div} \tilde{v}_t(t)\|_3^2 \\ \leq M_0 + CTP\left(\sup_{t \in [0, T]} E_\kappa(t)\right). \end{aligned} \quad (28)$$

Proof. Since $(\tilde{A}_\kappa)_i^j \tilde{v}^i_{,j} = 0$, we see that

$$(\tilde{A}_\kappa)_i^j D \tilde{v}^i_{,j} = -D(\tilde{A}_\kappa)_i^j \tilde{v}^i_{,j}. \quad (29)$$

Step 1. Estimate for $\operatorname{div} \Lambda_\kappa \tilde{\eta}$. It follows that

$$[(\tilde{A}_\kappa)_i^j D \tilde{\eta}^i_{,j}]_t = \partial_t (\tilde{A}_\kappa)_i^j D \tilde{\eta}^i_{,j} - D(\tilde{A}_\kappa)_i^j \tilde{v}^i_{,j}$$

so that

$$[(\tilde{A}_\kappa)_i^j D \tilde{\eta}^i_{,j}](t) = \int_0^t \left(\partial_t (\tilde{A}_\kappa)_i^j D \tilde{\eta}^i_{,j} - D(\tilde{A}_\kappa)_i^j \tilde{v}^i_{,j} \right) dt',$$

and hence

$$D \operatorname{div} \tilde{\eta}(t) = \underbrace{\int_0^t \partial_t (\tilde{A}_\kappa)_i^j D \tilde{\eta}^i_{,j} dt'}_{\mathcal{I}_1} - \underbrace{\int_0^t D(\tilde{A}_\kappa)_i^j \tilde{v}^i_{,j} dt'}_{\mathcal{I}_2} - \underbrace{\int_0^t \partial_t (\tilde{A}_\kappa)_i^j dt' D \tilde{\eta}^i_{,j}}_{\mathcal{I}_3}.$$

Thus,

$$\|D \operatorname{div} \Lambda_\kappa \tilde{\eta}(t)\|_{2.5}^2 \leq \sum_{i=1}^3 \|\Lambda_\kappa \mathcal{I}_i(t)\|_{2.5}^2.$$

Using Lemma 5.3 in the same fashion as was used for Proposition 2, we see that

$$\sum_{i=1}^3 \|\Lambda_\kappa \mathcal{I}_i(t)\|_{2.5}^2 \leq CTP\left(\sup_{t \in [0, T]} E_\kappa(t)\right),$$

from which it follows that

$$\|\operatorname{div} \Lambda_\kappa \tilde{\eta}(t)\|_{3.5}^2 \leq CTP\left(\sup_{t \in [0, T]} E_\kappa(t)\right).$$

Step 2. Estimate for $\operatorname{div} \tilde{v}$. From $(\tilde{A}_\kappa)_i^j \tilde{v}^i_{,j} = 0$, we see that

$$\operatorname{div} \tilde{v}(t) = - \int_0^t \partial_t (\tilde{A}_\kappa)_i^j dt' \tilde{v}^i_{,j}. \quad (30)$$

Hence, it is clear that

$$\|\operatorname{div} \tilde{v}(t)\|_3^2 + \|\kappa \operatorname{div} \tilde{v}(t)\|_{3.5}^2 \leq CTP\left(\sup_{t \in [0, T]} E_\kappa(t)\right).$$

Step 3. Estimate for $\operatorname{div} \tilde{v}_t$. From (30),

$$\operatorname{div} \tilde{v}_t(t) = -\partial_t(\tilde{A}_\kappa)_i^j \tilde{v}^i_{,j} - \int_0^t \partial_t(\tilde{A}_\kappa)_i^j dt' \tilde{v}^i_{,j} .$$

It follows that

$$\|\operatorname{div} \tilde{v}_t(t)\|_{2.5}^2 + \|\sqrt{\kappa} \operatorname{div} \tilde{v}_t(t)\|_3^2 \leq CTP(\sup_{t \in [0, T]} E_\kappa(t)) .$$

□

8.5. Pressure estimates. Letting $(\tilde{A}_\kappa)_i^j \frac{\partial}{\partial x_j}$ act on (16a), for $t \in [0, T_\kappa]$, the pressure function $q(x, t)$ satisfies the elliptic equation

$$\begin{aligned} -(\tilde{A}_\kappa)_i^j \left[(\tilde{A}_\kappa)_i^k \tilde{q}_{,k} \right]_{,j} &= \tilde{v}^i_{,j} (\tilde{A}_\kappa)_r^j \tilde{v}_{\kappa,s}^r (\tilde{A}_\kappa)_i^s \text{ in } \Omega , \\ \tilde{q} &= 0 \text{ on } \Gamma , \\ \tilde{q}_{,k} (\tilde{A}_\kappa)_i^k (\tilde{A}_\kappa)_i^3 &= \tilde{v} \cdot \partial_t (\tilde{A}_\kappa)_i^3 = 0 \text{ on } \{x_3 = 0\} . \end{aligned}$$

Using our conventions of Section 8.3 concerning the generic constant C , we have the standard elliptic estimate (see [3]) on $[0, T_\kappa]$

$$\|\tilde{q}(t)\|_{4.5} \leq C \|\Lambda_\kappa \tilde{\eta}(t)\|_{4.5} . \quad (31)$$

Similarly by time-differentiating the above elliptic equation for q , elliptic estimates show that

$$\|\tilde{q}_t(t)\|_4^2 \leq C (\|\Lambda_\kappa \tilde{\eta}(t)\|_{4.5}^2 + \|\tilde{v}(t)\|_4^2) .$$

8.6. Technical lemma. Our energy estimates require the use of the following

Lemma 8.5. *Let $H^{\frac{1}{2}}(\Omega)'$ denote the dual space of $H^{\frac{1}{2}}(\Omega)$. There exists a positive constant C such that*

$$\|\bar{\partial} F\|_{H^{\frac{1}{2}}(\Omega)'} \leq C \|F\|_{\frac{1}{2}} \quad \forall F \in H^{\frac{1}{2}}(\Omega) .$$

Proof. Integrating by parts with respect to the horizontal derivative yields for all $G \in H^1(\Omega)$,

$$\int_\Omega \bar{\partial} F G \, dx = - \int_\Omega F \bar{\partial} G \, dx \leq C \|F\|_0 \|G\|_1 ,$$

which shows that there exists $C > 0$ such that

$$\forall F \in L^2(\Omega), \quad \|\bar{\partial} F\|_{H^1(\Omega)'} \leq C \|F\|_0 . \quad (32)$$

Interpolating with the obvious inequality

$$\forall F \in H^1(\Omega), \quad \|\bar{\partial} F\|_{L^2(\Omega)} \leq C \|F\|_1$$

proves the lemma. □

8.7. Energy estimates. In this section, we take $T \in (0, T_\kappa)$.

Proposition 4. *For $t \in [0, T_\kappa]$,*

$$|\Lambda_\kappa \tilde{\eta}^3(t)|_4^2 + |\tilde{v}^3(t)|_{3.5}^2 \leq M_0 + CTP(\sup_{t \in [0, T]} E_\kappa(t)) . \quad (33)$$

Proof. Taking the $L^2(\Omega)$ inner-product $\bar{\partial}^4$ of (16a) with $\bar{\partial}^4 \tilde{v}^i$ yields

$$0 = \underbrace{\frac{1}{2} \frac{d}{dt} \|\bar{\partial}^4 \tilde{v}(t)\|_0^2}_{\mathcal{I}_1} + \underbrace{\int_{\Omega} \bar{\partial}^4 (\tilde{A}_{\kappa})_i^k \tilde{q}_{,k} \bar{\partial}^4 \tilde{v}^i dx}_{\mathcal{I}_2} + \underbrace{\int_{\Omega} (\tilde{A}_{\kappa})_i^k \bar{\partial}^4 \tilde{q}_{,k} \bar{\partial}^4 \tilde{v}^i dx}_{\mathcal{I}_3} + \mathcal{R}, \quad (34)$$

where \mathcal{R} denotes integrals consisting of lower-order terms which can easily be shown, via the Cauchy-Schwarz inequality, to satisfy

$$\int_0^T |\mathcal{R}(t)| dt \leq M_0 + C T P(\sup_{t \in [0, T]} E(t)).$$

Using the identity (5), we see that

$$\begin{aligned} \mathcal{I}_2 &= - \int_{\Omega} (\tilde{A}_{\kappa})_r^k \bar{\partial}^4 \tilde{\eta}_{\kappa, s}^r (\tilde{A}_{\kappa})_i^s \tilde{q}_{,k} \bar{\partial}^4 \tilde{v}^i dx + \mathcal{R} \\ &= - \underbrace{\int_{\Gamma} (\tilde{A}_{\kappa})_r^k \bar{\partial}^4 \tilde{\eta}_{\kappa}^r (\tilde{A}_{\kappa})_i^3 \tilde{q}_{,k} \bar{\partial}^4 \tilde{v}^i dx_h}_{\mathcal{I}_{2a}} + \underbrace{\int_{\Omega} (\tilde{A}_{\kappa})_r^k \bar{\partial}^4 \tilde{\eta}_{\kappa}^r (\tilde{A}_{\kappa})_i^s \tilde{q}_{,k} \bar{\partial}^4 \tilde{v}^i_{,s} dx}_{\mathcal{I}_{2b}} + \mathcal{R}. \end{aligned}$$

Since $\tilde{q} = 0$ on Γ , so that $\tilde{q}_{,k} = \tilde{q}_{,3}$, we see that

$$-\mathcal{I}_{2a} = \int_{\Gamma} (-\tilde{q}_{,3}) \bar{\partial}^4 \tilde{\eta}_{\kappa}^r (\tilde{A}_{\kappa})_r^3 \bar{\partial}^4 \tilde{v}^i (\tilde{A}_{\kappa})_i^3 dx_h,$$

Recalling that $\tilde{\eta}_{\kappa} = \Lambda_{\kappa} \Lambda_{\kappa} \tilde{\eta}$,

$$\begin{aligned} -\mathcal{I}_{2a} &= \underbrace{\int_{\Gamma} (-\tilde{q}_{,3}) \bar{\partial}^4 \Lambda_{\kappa} \tilde{\eta}^r (\tilde{A}_{\kappa})_r^3 \bar{\partial}^4 \Lambda_{\kappa} \tilde{v}^i (\tilde{A}_{\kappa})_i^3 dx_h}_{\mathcal{K}_1} \\ &\quad + \underbrace{\int_{\Gamma} \bar{\partial}^4 \Lambda_{\kappa} \tilde{\eta}^r \left[\Lambda_{\kappa} \left((-\tilde{q}_{,3}) (\tilde{A}_{\kappa})_i^3 (\tilde{A}_{\kappa})_r^3 \bar{\partial}^4 \tilde{v}^i \right) - (-\tilde{q}_{,3}) (\tilde{A}_{\kappa})_i^3 (\tilde{A}_{\kappa})_r^3 \Lambda_{\kappa} \bar{\partial}^4 \tilde{v}^i \right] dx_h}_{\mathcal{K}_2}, \end{aligned}$$

According to Lemma 5.1,

$$\begin{aligned} \mathcal{K}_2(t) &\leq |\bar{\partial}^4 \Lambda_{\kappa} \tilde{\eta}^r|_0 \left| \Lambda_{\kappa} \left((-\tilde{q}_{,3}) (\tilde{A}_{\kappa})_i^3 (\tilde{A}_{\kappa})_r^3 \bar{\partial}^4 \tilde{v}^i \right) - (-\tilde{q}_{,3}) (\tilde{A}_{\kappa})_i^3 (\tilde{A}_{\kappa})_r^3 \Lambda_{\kappa} \bar{\partial}^4 \tilde{v}^i \right|_0 \\ &\leq C |\bar{\partial}^4 \Lambda_{\kappa} \tilde{\eta}^r|_0 |\tilde{q}_{,3} (\tilde{A}_{\kappa})_i^3 (\tilde{A}_{\kappa})_r^3|_{W^{1,\infty}(\Gamma)} |\bar{\partial}^3 \tilde{v}^i|_0. \end{aligned}$$

By the Sobolev embedding theorem,

$$|\tilde{q}_{,3} (\tilde{A}_{\kappa})_i^3 (\tilde{A}_{\kappa})_r^3|_{W^{1,\infty}(\Gamma)} \leq C |\tilde{q}_{,3}|_{1.5} |(\tilde{A}_{\kappa})_i^3|_{1.5} |(\tilde{A}_{\kappa})_r^3|_{1.5},$$

so that

$$\int_0^T \mathcal{K}_2(t) dt \leq C T P(\sup_{t \in [0, T]} E_{\kappa}(t)).$$

To study the integral \mathcal{K}_1 , we define \tilde{n}_{κ} to be the outward unit normal to the moving surface $\tilde{\eta}_{\kappa}(t, \Gamma)$, given by $\tilde{n}_{\kappa} = (\tilde{a}_{\kappa})_i^3 / |(\tilde{a}_{\kappa})^3|$. From (8),

$$(\tilde{A}_{\kappa})^3 = \tilde{J}_{\kappa}^{-1} \sqrt{\tilde{g}_{\kappa}} n_{\kappa} \text{ on } \Gamma,$$

where

$$\sqrt{\tilde{g}_{\kappa}} = |\tilde{\eta}_{\kappa,1} \times \tilde{\eta}_{\kappa,2}| = |(\tilde{a}_{\kappa})^3| \text{ on } \Gamma.$$

It follows that

$$\begin{aligned} \mathcal{K}_1(t) &= \int_{\Gamma} (-\tilde{q}_{,3}) \bar{\partial}^4 \Lambda_{\kappa} \tilde{\eta} \cdot \tilde{n}_{\kappa} \bar{\partial}^4 \Lambda_{\kappa} \tilde{v} \cdot \tilde{n}_{\kappa} |\det \tilde{g}_{\kappa}| \tilde{J}_{\kappa}^{-2} dx_h \\ &= \frac{1}{2} \frac{d}{dt} \underbrace{\int_{\Gamma} (-\tilde{q}_{,3}) |\bar{\partial}^4 \Lambda_{\kappa} \tilde{\eta} \cdot \tilde{n}_{\kappa}|^2 |\det \tilde{g}_{\kappa}| \tilde{J}_{\kappa}^{-2} dx_h}_{\mathcal{K}_{1a}} \\ &\quad - \underbrace{\int_{\Gamma} \frac{1}{2} \bar{\partial}^4 \Lambda_{\kappa} \tilde{\eta}^i \bar{\partial}^4 \Lambda_{\kappa} \tilde{\eta}^j \partial_t [(\tilde{n}_{\kappa})_i (\tilde{n}_{\kappa})_j] |\det \tilde{g}_{\kappa}| \tilde{J}_{\kappa}^{-2} dx_h}_{\mathcal{K}_{1b}}. \end{aligned}$$

By the assumption of Section 8.3,

$$\left| \partial_t [(\tilde{n}_{\kappa})_i (\tilde{n}_{\kappa})_j] |\det \tilde{g}_{\kappa}| \tilde{J}_{\kappa}^{-2} \right|_{L^{\infty}(\Gamma)} \leq C,$$

from which it follows that

$$\int_0^T \mathcal{K}_{1b}(t) dt \leq C T P(\sup_{t \in [0, T]} E_{\kappa}(t)).$$

Using our assumed bounds for $-\tilde{q}_{,3}(t)$, $\det \tilde{g}_{\kappa}(t)$, \tilde{J}_{κ} on $[0, T]$, we see that

$$\bar{c} |\bar{\partial}^4 \Lambda_{\kappa} \tilde{\eta}(t) \cdot \tilde{n}_{\kappa}(t)|_0^2 \leq \int_0^T \mathcal{K}_1(t) dt + M_0 + C T P(\sup_{t \in [0, T]} E_{\kappa}(t)),$$

for a constant \bar{c} which depends on $\lambda, \tilde{g}_{\kappa}(0), \tilde{J}_{\kappa}(0)$.

Notice that $N = (0, 0, 1) = \tilde{n}_{\kappa}(t) - \int_0^t \partial_t \tilde{n}_{\kappa}(t') dt'$, and by our assumptions in Section 8.3, $|\partial_t \tilde{n}_{\kappa}(t)|_{L^{\infty}(\Gamma)} \leq C$, so that with $\tilde{\eta}^3 = \tilde{\eta} \cdot N$,

$$\bar{c} |\Lambda_{\kappa} \tilde{\eta}^3(t)|_4^2 \leq \int_0^T \mathcal{K}_1(t) dt + M_0 + C T P(\sup_{t \in [0, T]} E_{\kappa}(t)),$$

and hence

$$\bar{c} |\Lambda_{\kappa} \tilde{\eta}^3(t)|_4^2 \leq - \int_0^T \mathcal{I}_{2a}(t) dt + M_0 + C T P(\sup_{t \in [0, T]} E_{\kappa}(t)),$$

It remains to show that the integrals $\int_0^T \mathcal{I}_{2b}(t) dt$ and $\int_0^T \mathcal{I}_3(t) dt$ are both bounded by $C T P(\sup_{t \in [0, T]} E_{\kappa}(t))$. Using (16b),

$$\begin{aligned} \mathcal{I}_{2b}(t) &= - \int_{\Omega} (\tilde{A}_{\kappa})_r^k \bar{\partial}^4 \tilde{\eta}_{\kappa}^r \tilde{q}_{,k} \tilde{v}^i{}_{,s} \bar{\partial}^4 (\tilde{A}_{\kappa})_i^s dx + \mathcal{R} \\ &\leq C \|\bar{\partial}^4 \tilde{\eta}_{\kappa}(t)\|_{\frac{1}{2}} \|\bar{\partial}^4 \tilde{A}_{\kappa}(t)\|_{H^{\frac{1}{2}}(\Omega)} + \mathcal{R} \\ &\leq C \|\bar{\partial}^4 \tilde{\eta}_{\kappa}(t)\|_{\frac{1}{2}} \|\bar{\partial}^3 \tilde{A}_{\kappa}(t)\|_{H^{\frac{1}{2}}(\Omega)} + \mathcal{R} \\ &\leq C \sup_{t \in [0, T]} E(t) + \mathcal{R}, \end{aligned}$$

where we have used Lemma 8.5 for the second inequality.

Finally,

$$\begin{aligned} \mathcal{I}_3(t) &= - \int_{\Omega} \bar{\partial}^4 \tilde{q} \bar{\partial}^4 \tilde{v}^i{}_{,k} (\tilde{A}_{\kappa})_i^k dx = \int_{\Omega} \bar{\partial}^4 \tilde{q} \tilde{v}^i{}_{,k} \bar{\partial}^4 (\tilde{A}_{\kappa})_i^k dx + \mathcal{R} \\ &\leq C \|\bar{\partial}^3 \tilde{q}(t)\|_{\frac{1}{2}} \|\bar{\partial}^4 \tilde{A}_{\kappa}(t)\|_{H^{\frac{1}{2}}(\Omega)'} + \mathcal{R} \\ &\leq C \sup_{t \in [0, T]} E(t) + \mathcal{R}, \end{aligned}$$

where we have used the pressure estimate (31) and Lemma 8.5 for the last inequality.

Summing the estimates for \mathcal{I}_1 , \mathcal{I}_2 , \mathcal{I}_3 and integrating (34) from 0 to T , we obtain the inequality,

$$\sup_{t \in [0, T]} (|\tilde{\eta}^3|_4^2 + \|\bar{\partial}^4 \tilde{v}(t)\|_4^2) \leq M_0 + C T P(\sup_{t \in [0, T]} E_{\kappa}(t)).$$

According to Proposition 3,

$$\sup_{t \in [0, T]} \|\operatorname{div} \tilde{v}(t)\|_3^2 \leq M_0 + C T P(\sup_{t \in [0, T]} E_{\kappa}(t)),$$

from which it follows that

$$\sup_{t \in [0, T]} \|\bar{\partial}^4 \operatorname{div} \tilde{v}(t)\|_{H^1(\Omega)'}^2 \leq M_0 + C T P(\sup_{t \in [0, T]} E_{\kappa}(t)).$$

Hence, the normal trace estimate (10) shows that

$$\sup_{t \in [0, T]} (|\tilde{\eta}^3(t)|_4^2 + |\tilde{v}^3(t)|_{3.5}^2) \leq M_0 + C T P(\sup_{t \in [0, T]} E_{\kappa}(t)). \quad (35)$$

□

Combining Proposition 4 with the curl estimates in Proposition 2 and the divergence estimates in Proposition 3 for $\eta(t)$ and $v(t)$ and using (11) provides the following

Proposition 5. For $t \in [0, T_{\kappa}]$,

$$\|\Lambda_{\kappa} \tilde{\eta}(t)\|_{4.5}^2 + \|\tilde{v}(t)\|_4^2 \leq M_0 + C T P(\sup_{t \in [0, T]} E_{\kappa}(t)). \quad (36)$$

Since $\tilde{v}_t = -\tilde{A}_{\kappa}^T D \tilde{q}$, we also obtain

Proposition 6. For $t \in [0, T_{\kappa}]$,

$$\|\tilde{v}_t(t)\|_{3.5}^2 \leq M_0 + C T P(\sup_{t \in [0, T]} E_{\kappa}(t)). \quad (37)$$

Proposition 7. For $t \in [0, T_{\kappa}]$ and $\alpha = 1, 2$,

$$|\sqrt{\kappa} \tilde{v}_t^{\alpha}(t)|_{3.5}^2 + |\kappa \tilde{v}^{\alpha}(t)|_4^2 \leq M_0 + C T P(\sup_{t \in [0, T]} E_{\kappa}(t)). \quad (38)$$

Proof. Multiplying (16a) by $\tilde{\eta}_{\kappa}^j$, we find the identity

$$\tilde{v}_t \cdot \tilde{\eta}_{\kappa, k} = -\tilde{q}_{,k} \text{ for } k = 1, 2, 3.$$

It follows that

$$\tilde{v}_t \cdot \bar{\partial} \eta_{\kappa} = -\bar{\partial} \tilde{q} = 0 \text{ on } \Gamma,$$

so that the tangential component (with respect to the moving boundary $\tilde{\eta}_{\kappa}(t, \Gamma)$) of \tilde{v}_t vanishes on Γ . Hence,

$$\bar{\partial}^3 \tilde{v}_t \cdot \bar{\partial} \tilde{\eta}_{\kappa} = -\tilde{v}_t \cdot \bar{\partial}^4 \tilde{\eta}_{\kappa} - 3 \bar{\partial}^2 \tilde{v}_t \cdot \bar{\partial}^2 \tilde{\eta}_{\kappa} - 3 \bar{\partial} \tilde{v}_t \cdot \bar{\partial}^3 \tilde{\eta}_{\kappa}.$$

Thus,

$$|\bar{\partial}^3 \tilde{v}_t \cdot \bar{\partial} \tilde{\eta}_\kappa|_0 \leq C |\tilde{v}_t|_{2.5} |\tilde{\eta}_\kappa|_4 \leq C |\tilde{\eta}_\kappa|_4,$$

so that using the fundamental theorem of calculus, together with the fact that $\eta_{\kappa,\alpha}(0)$ is proportional to T_α for $\alpha = 1, 2$, we find that

$$|\tilde{v}_t^\alpha|_3 \leq C |\tilde{\eta}_\kappa|_4 + C T |\tilde{v}_t|_3, \quad (39)$$

Similarly,

$$|\bar{\partial}^4 \tilde{v}_t \cdot \bar{\partial} \tilde{\eta}_\kappa|_0 \leq C |\tilde{v}_t|_3 |\tilde{\eta}_\kappa|_5 \leq C |\tilde{\eta}_\kappa|_5,$$

from which it follows that

$$|\tilde{v}_t^\alpha|_4 \leq C |\tilde{\eta}|_5 + C T |\tilde{v}_t|_4. \quad (40)$$

Interpolation between (39) and (40) yields

$$|\tilde{v}_t^\alpha|_{3.5} \leq C |\tilde{\eta}_\kappa|_{4.5} + C T |\tilde{v}_t|_{3.5}.$$

Interpolation between the inequalities $|\tilde{\eta}_\kappa|_4 \leq C |\Lambda_\kappa \tilde{\eta}|_4$ and $|\tilde{\eta}_\kappa|_5 \leq \frac{C}{\kappa} |\Lambda_\kappa \tilde{\eta}|_4$ shows that $\sqrt{\kappa} |\tilde{\eta}_\kappa|_{4.5} \leq C |\Lambda_\kappa \tilde{\eta}|_4$. It thus follows from (36) and (37) that

$$\sqrt{\kappa} |\tilde{v}_t^\alpha(t)|_{3.5} \leq M_0 + C T P\left(\sup_{t \in [0, T]} E_\kappa(t)\right).$$

A similar argument shows that

$$\kappa |\tilde{v}^\alpha(t)|_4 \leq M_0 + C T P\left(\sup_{t \in [0, T]} E_\kappa(t)\right).$$

□

Combining the estimate (38) together with the curl estimates in Proposition 2 and divergence estimates in Proposition 3 for $\kappa \tilde{v}(t)$ and $\sqrt{\kappa} \tilde{v}_t(t)$ proves the following

Proposition 8. *For $t \in [0, T_\kappa]$ and $\alpha = 1, 2$,*

$$\|\sqrt{\kappa} \tilde{v}_t(t)\|_4^2 + \|\kappa \tilde{v}(t)\|_{4.5}^2 \leq M_0 + C T P\left(\sup_{t \in [0, T]} E_\kappa(t)\right). \quad (41)$$

9. Proof of the Main Theorem.

9.1. Time of existence and bounds independent of κ and existence of solutions to (3). Summing the inequalities provided by the above Propositions, we find that

$$\sup_{t \in [0, T]} E_\kappa(t) \leq M_0 + C T P\left(\sup_{t \in [0, T]} E_\kappa(t)\right).$$

Just as in Section 9 of [2], this provides us with a time of existence T_1 independent of κ and an estimate on $(0, T_1)$ independent of κ of the type:

$$\sup_{t \in [0, T_1]} E_\kappa(t) \leq 2M_0. \quad (42)$$

In particular, our sequence of solutions $(\tilde{\eta}, \tilde{v})$ satisfy the κ -independent bound (42) on the κ -independent time-interval $(0, T_1)$.

9.2. The limit as $\kappa \rightarrow 0$. By the κ -independent estimate (42), there exists a subsequence of $\{\tilde{v}_t, \tilde{A}_\kappa, D\tilde{q}\}$ which converges uniformly to (v_t, A, Dq) where $A = [D\eta]^{-1}$, and $\eta = e + \int_0^t v dt'$. Standard arguments show that (η, v) solve (3), and that

$$\sup_{t \in [0, T_1]} E(t) \leq P(E(0)).$$

9.3. Uniqueness. Suppose that on $[0, T_1]$, (η^1, v^1, q^1) and (η^2, v^2, q^2) are both solutions of (3) with initial data $u_0 \in H_{\text{div}}^5(\Omega)$ and $\text{curl } u_0 \in H^{4.5}(\Omega)$, and with both q_1 and q_2 satisfying the stability condition $-\partial q_i / \partial N > 0$ on Γ for $i = 1, 2$.

Setting

$$\mathcal{E}_\eta(t) = 1 + \|\eta(t)\|_{5.5}^2 + \|v(t)\|_5^2,$$

by the method of the previous section (with $\kappa = 0$), we infer that both $\mathcal{E}_{\eta^1}(t)$ and $\mathcal{E}_{\eta^2}(t)$ are bounded by a constant \mathcal{M}_0 depending on the data u_0 and Γ on a time interval $0 \leq t \leq T_1$ for T_1 small enough.

Let

$$w := v^1 - v^2, \quad r := q^1 - q^2, \quad \text{and } \xi := \eta^1 - \eta^2.$$

Then (ξ, w, r) satisfies

$$\xi = \int_0^t w \quad \text{in } \Omega \times (0, T], \quad (43a)$$

$$\partial_t w^i + (a^1)_i^k r_{,k} = (a^2 - a^1)_i^k q_{,k}^2 \quad \text{in } \Omega \times (0, T], \quad (43b)$$

$$(a^1)_i^j w^{i,j} = (a^2 - a^1)_i^j v^{2^i}_{,j} \quad \text{in } \Omega \times (0, T], \quad (43c)$$

$$r = 0 \quad \text{on } \Gamma \times (0, T], \quad (43d)$$

$$(\xi, w) = (0, 0) \quad \text{on } \Omega \times \{t = 0\}. \quad (43e)$$

We set

$$E(t) = 1 + \|\xi(t)\|_{4.5}^2 + \|w(t)\|_4^2.$$

We will show that $E(t) = 0$, which shows that $w = 0$. We follow the identical analysis as in the previous section, and estimate the new error terms, arising from the difference of two solutions, using the additional space regularity in our assumptions. With (43e), we find that $\sup_{t \in [0, T]} E(t) \leq C T P(\sup_{t \in [0, T]} E(t))$.

9.4. Optimal regularity for initial data. We smoothed our initial data u_0 in order to construct solutions to the κ -approximation (16). Having obtained solutions which depend only on $E(0)$, a standard density argument shows that the initial data needs only to satisfy $E(0) < \infty$. All assumptions on from Section 8.3 can now be verified by the fundamental theorem of calculus, and taking T_1 even small if necessary.

Acknowledgments. SS was supported by the National Science Foundation under grant DMS-0701056. We are grateful to the referees for there extremely careful reading of the manuscript. Their numerous suggestions have benefited the presentation immensely.

REFERENCES

- [1] D. Ambrose and N. Masmoudi, *The zero surface tension limit of three-dimensional water waves*, Indiana Univ. Math. J., **58** (2009), 479–521.
- [2] D. Coutand and S. Shkoller, *The interaction between quasilinear elastodynamics and the Navier-Stokes equations*, Arch. Rational Mech. Anal., **179** (2006), 303–352.
- [3] D. Coutand and S. Shkoller, *Well-posedness of the free-surface incompressible Euler equations with or without surface tension*, J. Amer. Math. Soc., **20** (2007), 829–930.
- [4] D. Ebin and J. Marsden, *Groups of diffeomorphisms and the motion of an incompressible fluid*, Ann. of Math., **92** (1970), 102–163.
- [5] D. Lannes, *Well-posedness of the water-waves equations*, J. Amer. Math. Soc., **18** (2005), 605–654.
- [6] H. Lindblad, *Well-posedness for the motion of an incompressible liquid with free surface boundary*, Annals of Math., **162** (2005), 109–194.

- [7] V. I. Nalimov, *The Cauchy-Poisson problem* (in Russian), *Dynamika Splosh. Sredy*, **18** (1974), 104–210.
- [8] J. Shatah and C. Zeng, *Geometry and a priori estimates for free boundary problems of the Euler equation*, *Comm. Pure Appl. Math.*, **61** (2008), 698–744.
- [9] M. Taylor, “Partial Differential Equations, Vol. **I–III**,” Springer, 1996.
- [10] R. Temam, “Navier-Stokes Equations. Theory and Numerical Analysis,” Third edition. *Studies in Mathematics and its Applications*, 2. North-Holland Publishing Co., Amsterdam, 1984.
- [11] S. Wu, *Well-posedness in Sobolev spaces of the full water wave problem in 2-D*, *Invent. Math.*, **130** (1997), 39–72.
- [12] S. Wu, *Well-posedness in Sobolev spaces of the full water wave problem in 3-D*, *J. Amer. Math. Soc.*, **12** (1999), 445–495.
- [13] H. Yosihara, *Gravity waves on the free surface of an incompressible perfect fluid*, *Publ. Res. Inst. Math. Sci.*, **18** (1982), 49–96.
- [14] P. Zhang and Z. Zhang, *On the free boundary problem of three-dimensional incompressible Euler equations*, *Comm. Pure Appl. Math.*, **61**, (2008), 877–940.

Received November 2009; revised January 2010.

E-mail address: D.Coutand@ma.hw.ac.uk

E-mail address: shkoller@math.ucdavis.edu