

Math 210 § 15.2

①

Mass/Density


Center of Mass

Moments of Inertia

Start with simpler
case of $(x, y) \in \mathbb{R}^2$
density = $\delta(x, y) \in \mathbb{R}$

- Calculus extends the physics of point masses to the continuum
- Basic Set up - Fluid Dynamics

Air Foil



$$\delta(x, y) = \text{density} = \frac{\text{mass}}{\text{area}}$$


$$\vec{v} = (v_1, v_2) = \text{velocity}$$

- Start with the density

2

$$\delta(x, y) = \frac{\text{Mass}}{\text{Vol}} \equiv \frac{\text{Mass}}{\text{Area}}$$

2-D plate



$$\delta(x, y) = \lim_{\substack{\Delta x \rightarrow 0 \\ \Delta y \rightarrow 0}} \frac{\Delta M}{\Delta x \Delta y}$$

Eg: Density can be given by a formula:

$$\delta(x, y) = xy^2 = \frac{\text{mass}}{\text{area}}$$

Q: Given the density, how do you recover the total mass in Region R ?

- The Idea: Give a precise definition of $\iint_R f(x,y) dA$ in terms of a Riemann Sum... (3)


- Identify the total mass as the limit of a Riemann Sum...

- Evaluate the total mass by evaluating the integral, which is $\iint_R \delta(x,y) dA$

- We start by giving a precise definition of the Riemann Integral

$$\iint_R \underbrace{f(x,y)}_{\text{general function } f} dA$$

Defn of $\iint_R f(x,y) dA$ (4)



Mesh

$$a = x_0 < x_1 < \dots < x_N = b$$

$$\Delta x = \frac{b-a}{N}$$

$$c = y_0 < y_1 < \dots < y_N = d$$

$$\Delta y = \frac{d-c}{N}$$

Defn: $\iint_R f(x,y) dA = \lim_{N \rightarrow \infty} \sum_{(x_i, y_i) \in R} f(x_i, y_i) \Delta x \Delta y$

Riemann Sum

Approximate Volume
above ΔR_{ij}

5

$$\iint_R f(x,y) dA \equiv \lim_{N \rightarrow \infty} \sum_{(x_i, y_i) \in R} f(x_i, y_i) \Delta x \Delta y$$

Riemann Sum

Idea: You only sum over rectangles w. mesh points in R


$\Rightarrow$ error tends to zero as
 $N \rightarrow \infty$

Defn of total mass

Ex Let R be a metal plate with density $\delta(x, y) = \frac{\text{mass}}{\text{area}}$

Q: How do you define the Total Mass of R ?

Ans: Define it by a Riemann Sum



"Mass in Rectangle ΔR_{ij} " = ΔM_{ij}

$$\Delta M_{ij} = \underbrace{\delta(x_i, y_i)}_{\substack{\text{mass} \\ \text{area}}} \cdot \underbrace{\Delta x \Delta y}_{\text{Area}}$$

Defn: Total Mass $M = \lim_{N \rightarrow \infty} \sum_{i,j \in R} \underbrace{\delta(x_i, y_i) \Delta x \Delta y}_{\Delta M_{ij}}$

$$= \iint_R \delta(x, y) dA$$

8
a **Big Idea**: How to go from point masses to the **Continuum**

- Identify a continuous physical quantity as the limit of a Riemann Sum
- Define it to be the integral

Simplest example:


$\delta(x,y)$ = density (continuous distributed mass)

$$\text{Total Mass} = \iint_R \delta(x,y) dA$$

Similarly: Center of Mass Moments of Inertia

- Center of Mass

Q: Given $\delta(x, y)$ what are the coordinates $(\bar{x}, \bar{y})$ of the center of Mass?



Claim:


$$\bar{x} = \frac{\int_A x \delta(x, y) dA}{M}$$

$$\bar{y} = \frac{\int_A y \delta(x, y) dA}{M}$$

Center of Mass

10

Ex 1-d Find the point of Balance



• At $\bar{x}$: $m_1 d_1 = m_2 d_2$

OR: $(\bar{x} - x_1) m_1 = (x_2 - \bar{x}) m_2$

$$(\bar{x} - x_1) m_1 + \underbrace{(\bar{x} - x_2) m_2}_{\text{neg}} = 0$$

• Solve for $\bar{x}$ to get CM


$$\bar{x} (m_1 + m_2) = x_1 m_1 + x_2 m_2$$

$$\bar{x} = \frac{x_1 m_1 + x_2 m_2}{m_1 + m_2}$$

(It didn't matter where I put $x=0$)

• If there were N -masses

$m_1, \dots, m_N$



Condition for balance

$$(\bar{x} - x_1)m_1 + (\bar{x} - x_2)m_2 + \dots + (\bar{x} - x_N)m_N = 0$$

$$\sum_{k=1}^N (\bar{x} - x_k)m_k = 0$$

neg to right of $\bar{x}$
pos to left


Solve for $\bar{x}$!

$$\bar{x} \sum_{k=1}^N m_k = \sum_{k=1}^N x_k m_k$$

$$\bar{x} = \frac{\sum_{k=1}^N x_k m_k}{\sum_{k=1}^N m_k}$$

Q Consider now a density $\delta(x)$ (1-dimension $x \in \mathbb{R}$)

(12)




$$CM = \lim_{N \rightarrow \infty} \frac{\sum_{k=1}^N x_k \delta(x_k) \Delta x}{\sum_{k=1}^N \delta(x_k) \Delta x}$$

$$= \frac{\int_a^b x \delta(x) dx}{\int_a^b \delta(x) dx} = \frac{M_y}{M}$$

Similarly: in 2-dimensions $\mathbb{R}^2$

(13)



Balance point in x is $\bar{x}$

$$\bar{x} = \lim_{N \rightarrow \infty} \frac{\sum x_i \Delta m_i}{\sum \Delta m_i}$$

$$= \lim_{N \rightarrow \infty} \frac{\sum_{i,j \in R} x_i \delta(x_i, y_i) \Delta x \Delta y}{\sum_{i,j \in R} \delta(x_i, y_i) \Delta x \Delta y}$$

$$= \frac{\left(\iint_R x \delta(x, y) dA \right)}{\left(\iint_R \delta(x, y) dA \right)} = \frac{M_y}{M}$$

Conclude:

$$M_y = \iint_R x \delta(x, y) dA$$

distance to
y-axis is x

$$M = \iint_R \delta(x, y) dA$$

$$\bar{x} = \frac{M_y}{M}, \quad \bar{y} = \frac{M_x}{M}$$