

Continuation of Vector/Matrix Review

Note Title

* Range & Nullspace (or Kernel)

Def. $A \in \mathbb{R}^{m \times n}$.

$\text{range}(A) := \{y \in \mathbb{R}^m \mid y = Ax, x \in \mathbb{R}^n\}$


often written as $\text{Ran}(A)$ or $\text{Im}(A)$.

This is also called the column space of A .

$\text{null}(A) := \{x \in \mathbb{R}^n \mid Ax = 0\}$

is called the nullspace (or kernel) of A

$\text{Ker}(A)$



Ihm $\text{range}(A) = \text{span}\{\alpha_1, \dots, \alpha_n\}$
= a set of all possible
linear combi. of $\{\alpha_1, \dots, \alpha_n\}$

(Proof) Need to show two things

(1) $\text{range}(A) \subset \text{span}\{\alpha_1, \dots, \alpha_n\}$

(2) $\text{span}\{\alpha_1, \dots, \alpha_n\} \subset \text{range}(A)$

Now, (1) is easy since any $y \in \text{range}(A)$ by definition, $\exists x \in \mathbb{R}^n$ s.t. $y = Ax$.

This is a lin. combi. of col vectors of A .
So, $y \in \text{span}\{\alpha_1, \dots, \alpha_n\}$.

(2) Take any $y \in \text{span}\{\alpha_1, \dots, \alpha_n\}$.

By definition, $\exists \{x_1, \dots, x_n\}$ s.t.

$$y = x_1 \alpha_1 + \dots + x_n \alpha_n = A \mathbf{x} \in \text{range}(A)$$

by setting $\mathbf{x} = (x_1, \dots, x_n)^T$ //

* Linear Independence, Bases

Def. The vectors $\{\alpha_1, \dots, \alpha_n\}, \alpha_j \in \mathbb{R}^m$ are called **linearly independent** if

$$\sum_{j=1}^n x_j \alpha_j = \mathbf{0} \iff x_j = 0, 1 \leq j \leq n$$


A set of m linearly independent vectors in \mathbb{R}^m is called a **basis** in \mathbb{R}^m . \Rightarrow a matrix representation of a basis in \mathbb{R}^m is an $m \times m$ matrix.
Note that any vector in \mathbb{R}^m can be written as a lin. combi. of the m basis vectors in \mathbb{R}^m

Def. The **dimension** of $\text{span}\{\alpha_1, \dots, \alpha_n\}$ is the maximal number of linearly independent vectors among $\{\alpha_1, \dots, \alpha_n\}$

i.e., if $\exists j, \alpha_j = x_1 \alpha_1 + \dots + x_{j-1} \alpha_{j-1} + x_{j+1} \alpha_{j+1} + \dots + x_n \alpha_n$

then such α_j is useless in some sense (or more precisely, it is redundant).

Ex.



In \mathbb{R}^3 , these are linearly dependent.

$$\alpha_1 = \alpha_2 + \alpha_3$$

$$\text{So } \dim \text{span}\{\alpha_1, \alpha_2, \alpha_3\} = 2.$$

We cannot write any vector in \mathbb{R}^3 by a lin. combi. of $\{\alpha_2, \alpha_3\}$.

Only a certain subset of vectors in \mathbb{R}^3 can be written as a lin. combi. of $\{\alpha_2, \alpha_3\}$ (no control on the first and second entries of a vector in \mathbb{R}^3 .)

* Rank

Def. The **column rank** of A

$$:= \dim(\text{range}(A))$$

= # of linearly indep. col.vec's of A.

The **row rank** of A

$$:= \dim(\text{range}(A^T))$$

= # of linearly indep. row vec's of A.

$$\text{rank}(A) := \dim(\text{range}(A))$$

$A \in \mathbb{R}^{m \times n}$ is said to be of **full rank** if

$$\underline{\text{rank}(A) = \min(m, n)}.$$

Thm. $A \in \mathbb{R}^{m \times n}$, $m \geq n$ is of full rank
 $\Leftrightarrow \forall \mathbf{x}, \mathbf{y} \in \mathbb{R}^n$, $\mathbf{x} \neq \mathbf{y}$,
 $A\mathbf{x} \neq A\mathbf{y}$.

(Proof) \Rightarrow If $\text{rank}(A) = n$, i.e., full rank,
then $\{\mathbf{a}_1, \dots, \mathbf{a}_n\}$ are lin. indep.

So, they form a basis of range(A).

This means that $\forall \mathbf{b} \in \text{range}(A)$,
 $\exists!$ $\mathbf{x} \in \mathbb{R}^n$ s.t. $\mathbf{b} = \sum_{j=1}^n x_j \mathbf{a}_j$.
There exists a unique

\Leftarrow Suppose A is not of full rank.

Then $\{\mathbf{a}_1, \dots, \mathbf{a}_n\}$ are linearly dep.

i.e., $\exists \mathbf{c} \in \mathbb{R}^n$, $\mathbf{c} \neq \mathbf{0}$ s.t.

$$\sum_{j=1}^n c_j \mathbf{a}_j = \mathbf{0}, \text{ i.e., } A\mathbf{c} = \mathbf{0}.$$

Then set $\mathbf{y} = \mathbf{x} + \mathbf{c} \neq \mathbf{x}$.

$$\begin{aligned} \text{But } A\mathbf{y} &= A(\mathbf{x} + \mathbf{c}) = A\mathbf{x} + \underline{A\mathbf{c}} = \mathbf{0} \\ &= A\mathbf{x} \text{ contradiction!} \end{aligned}$$

#

* Inverse

Def. A is said to be **nonsingular**
or **invertible** \Leftrightarrow A is **square** and
of **full rank**

Hence $A \in \mathbb{R}^{m \times m}$: nonsingular
 $\Rightarrow \{\mathbf{a}_1, \dots, \mathbf{a}_m\}$ form a basis of \mathbb{R}^m

That means : the canonical basis vector $e_j \in \mathbb{R}^m$ can also be written as a lin. combi. of $\{q_1, \dots, q_m\}$

$$e_j = \sum_{i=1}^m z_{ij} q_i, \exists z_{ij}, 1 \leq i \leq m$$

$$\Rightarrow e_j = A z_j, z_j = (z_{1j}, \dots, z_{mj})^T$$

$$\text{So, } [e_1 | e_2 | \dots | e_m] = A [z_1 | z_2 | \dots | z_m]$$

$$\Leftrightarrow \underline{\underline{I}} = A \underline{\underline{Z}}$$

$m \times m$ identity matrix

Such matrix $Z \in \mathbb{R}^{m \times m}$ is called the **inverse** of A and written as $\underline{\underline{A}}^{-1}$.

Any nonsingular matrix has a unique inverse, and $A \underline{\underline{A}}^{-1} = \underline{\underline{A}}^{-1} A = \underline{\underline{I}}$.

Thm (Equivalences of a nonsingular matrix)

For $A \in \mathbb{R}^{m \times m}$, the following statements are equivalent :


- (a) A has an inverse A^{-1}
- (b) $\text{rank}(A) = m$
- (c) $\text{range}(A) = \mathbb{R}^m$
- (d) $\text{null}(A) = \{ \underline{0} \}$
- (e) 0 is not an eigenvalue of A
- (f) 0 is not a singular value of A
- (g) $\det(A) \neq 0$.

* Matrix A^{-1} \times vector

$$\begin{aligned} \mathbf{y} &= A \mathbf{x}, \quad A: \text{nonsingular} \\ \Rightarrow \mathbf{x} &= A^{-1} \mathbf{y}. \end{aligned}$$

This means that $A^{-1} \mathbf{y}$ represents an expansion coefficients of \mathbf{y} in the basis of col's of A .

So, Multiplication by A^{-1} is a change of basis operation!



Note : $\mathbf{y} = \mathbf{I}^{-1} \mathbf{y}$.