

# MAT 271: Applied & Computational Harmonic Analysis Supplementary Notes I by Naoki Saito

## The Generalized Functions

- The generalized functions have more singular behavior than functions (thus the name “generalized functions”), and are always defined as *linear functionals* on the *dual space*. Thus, before we discuss the generalized functions, we need to know the following.

**Definition:** Let  $\mathcal{X}$  be a vector space over, say,  $\mathbb{C}$ . A linear map from  $\mathcal{X}$  to  $\mathbb{C}$  is called a *linear functional* on  $\mathcal{X}$ . If  $\mathcal{X}$  is a normed vector space, then the space  $\mathcal{L}(\mathcal{X}, \mathbb{C})$  of *bounded linear functionals* on  $\mathcal{X}$  is called the *dual space*, and denoted by  $\mathcal{X}^*$  (or  $\mathcal{X}'$ ).

Examples: The dual of  $L^p(\mathbb{R})$ ,  $1 < p < \infty$ , is  $L^q(\mathbb{R})$ , where  $(1/p) + (1/q) = 1$ . These numbers are called *conjugate exponents*. In particular,  $L^2$  is self dual. Similarly, the dual of the sequence space  $\ell^p(\mathbb{Z})$  is  $\ell^q(\mathbb{Z})$ .

**Hölder’s Inequality:** Let  $p$  and  $q$  are conjugate exponents. Then for any  $f \in L^p$ ,  $g \in L^q$ , we have

$$\|fg\|_1 \leq \|f\|_p \|g\|_q.$$

(As you can see, the Cauchy-Schwarz inequality is a special version of this with  $p = q = 2$ . The proof is a great exercise.)

**The Riesz Representation Theorem:** Suppose  $p$  and  $q$  are conjugate exponents with  $1 < p < \infty$ . Then for each linear functional  $\varphi \in (L^p)^*$ , there exists  $g \in L^q$  such that  $\varphi(f) = \int f(x)g(x) dx$  for all  $f \in L^p$ . In other words,  $(L^p)^*$  is isometrically isomorphic to  $L^q$ .

- The more singular the class of the generalized functions, the more regular its dual.
- We now define the *Schwartz class*  $\mathcal{S} := \{f \in C^\infty(\mathbb{R}) \mid \sup_{x \in \mathbb{R}} |x^k \partial^\ell f| < \infty, \text{ for any } k, \ell \in \mathbb{N}\}$ , which are very smooth and decay faster than any polynomial at infinity, i.e., a very nice class of functions. An example: The Gaussian  $g(x) = e^{-x^2}$ .
- Then, we consider the dual  $\mathcal{S}'$ . You can imagine that members of this class can be very singular or “spiky.” This dual space is called the *tempered distributions*. Being as a linear functional, each member of  $\mathcal{S}'$  acts on the Schwartz functions. More precisely, if  $F \in \mathcal{S}'$  and  $\phi \in \mathcal{S}$ , then the value of  $F$  at  $\phi$  ( $F$  is a *linear map* from  $\mathcal{S}$  to  $\mathbb{C}$ !!) is denoted as  $\langle F, \phi \rangle = F(\phi) = \int F(x)\phi(x) dx$ .
- An example: *the Dirac delta function*  $\delta(x) \in \mathcal{S}'$  is defined as  $\langle \delta, \phi \rangle = \phi(0)$ . In other words,  $\int_{-\infty}^{\infty} \delta(x)\phi(x) dx = \phi(0)$ .
- For any  $F \in \mathcal{S}'$  and any  $\phi \in \mathcal{S}$ , we can define the following operations:

**Differentiation:**  $\langle \partial_x^k F, \phi \rangle = (-1)^k \langle F, \partial_x^k \phi \rangle$ .

This can be shown by integration by parts. An example:  $\langle \delta', \phi \rangle = -\langle \delta, \phi' \rangle = -\phi'(0)$ .

**Convolution:**  $F * \phi(x) = \langle F, \tau_x \tilde{\phi} \rangle$ , where  $\tilde{\phi}(y) = \phi(-y)$ .

An example:  $(\delta * \phi)(x) = \phi(x)$ .

**Fourier transform:**  $\langle \hat{F}, \phi \rangle = \langle F, \hat{\phi} \rangle$ .

An example:  $F = \delta$ , then  $\langle \hat{\delta}, \phi \rangle = \langle \delta, \hat{\phi} \rangle = \hat{\phi}(0)$ . This essentially shows that  $\hat{\delta}(\xi) \equiv 1$ . Using the translation operator, we can also have  $\mathcal{F}\{\delta(x-a)\} = e^{-2\pi i \xi a}$ , and  $\mathcal{F}\{e^{-2\pi i x a}\} = \delta(\xi+a)$ .

- **Definition:** A tempered distribution  $F$  on  $\mathbb{R}$  is called *periodic* with period  $A$  if  $\langle F, \tau_A \phi \rangle = \langle F, \phi \rangle$  for all  $\phi \in \mathcal{S}$ . A sequence of tempered distributions  $\{F_n\}$  is said to *converge temperately* to a tempered distribution  $F$  if  $\langle F_n, \phi \rangle \rightarrow \langle F, \phi \rangle$  as  $n \rightarrow \infty$  for all  $\phi \in \mathcal{S}$ . (See that all these operations and definitions are now moved to the *nice spouses* of  $F$ !)

- **[Theorem]** If  $F$  is a periodic tempered distribution, then  $F$  can be expanded in a temperately convergent Fourier series,  $F(x) = \frac{1}{\sqrt{A}} \sum_{-\infty}^{\infty} \alpha_k e^{2\pi i k x / A}$ , i.e.,  $\langle F, \phi \rangle = \sum_{-\infty}^{\infty} \alpha_k \left\langle \frac{1}{\sqrt{A}} e^{2\pi i k \cdot / A}, \phi \right\rangle$  for all  $\phi \in \mathcal{S}$ . Moreover, the coefficients  $\alpha_k$  satisfy  $\alpha_k \leq C(1 + |k|)^N$  for some  $C, N \geq 0$ . Conversely, if  $\{\alpha_k\}$  is any sequence satisfying this estimate, the series  $\frac{1}{\sqrt{A}} \sum_{-\infty}^{\infty} \alpha_k e^{2\pi i k x / A}$  converges temperately to a periodic tempered distribution.

- Define the *Shah function* (or *comb function*),  $\text{III}_A(x) := \sum_{k=-\infty}^{\infty} \delta(x - kA)$ . The facts about this function:


1. Since this is a periodic tempered distribution, we can expand it into the temperately convergent Fourier series;  $\text{III}_A(x) \sim \frac{1}{A} \sum_{-\infty}^{\infty} e^{2\pi i k x / A}$ . Note that  $\alpha_k \equiv 1/\sqrt{A}$  for all  $k \in \mathbb{Z}$ .

2.  $\mathcal{F}\{\text{III}_A\}(\xi) = \frac{1}{A} \text{III}_{1/A}(\xi) = \frac{1}{A} \sum_{-\infty}^{\infty} \delta(\xi - \frac{k}{A})$ .

- Using the Shah function and its Fourier transform, we can see that the Fourier transform of the Fourier series of a periodic function on  $[-A/2, A/2]$  as follows:

$$f(x) \sim \frac{1}{\sqrt{A}} \sum_{-\infty}^{\infty} \alpha_k e^{2\pi i k x / A} \xrightarrow{\mathcal{F}} \frac{1}{\sqrt{A}} \sum_{-\infty}^{\infty} \alpha_k \delta(\xi - \frac{k}{A}) \quad \text{i.e., line spectrum (discrete)}$$

As you can see, as  $A$  gets large, we are doing the finer sampling in the frequency domain, i.e.,



- **Periodization of a function with compact support  $\iff$  Discretization in frequency domain (with amplitude rescaling)**

For the details of the facts in these notes, see [1, Chap. 9], [2, Chap. 9], [3, Chap. 1].

## References

[1] G. B. FOLLAND, *Fourier Analysis and Its Applications*, Amer. Math. Soc., Providence, RI, 1992. Republished by AMS, 2009.

[2] ———, *Real Analysis: Modern Techniques and Their Applications*, John Wiley & Sons, Inc., 2nd ed., 1999.

[3] E. M. STEIN AND G. WEISS, *Introduction to Fourier Analysis on Euclidean Spaces*, Princeton Univ. Press, 1971.