

Lecture 16 : Wavelet Bases II

Note Title

★ Conjugate Mirror Filters

A whole MRA is entirely characterized by the scaling fn ϕ since it generates V_0 and consequently all V_j 's. $j \in \mathbb{Z}$.

An interesting thing is that any scaling fn is specified by a discrete filter called **conjugate mirror filter** (CMF).

- The scaling (or two-scale difference) eqn:

Recall $V_1 \subset V_0$, and $\frac{1}{\sqrt{2}} \phi(\frac{x}{2}) \in V_1$.

Hence $\frac{1}{\sqrt{2}} \phi(\frac{x}{2}) = \sum_{k \in \mathbb{Z}} h_k \phi(x-k)$

This is an expansion of $\frac{1}{\sqrt{2}} \phi(\frac{x}{2}) \in V_1 \subset V_0$ w.r.t. $\{\phi(x-k)\}_{k \in \mathbb{Z}}$ (an ONB of V_0).

$$h_k = \langle \frac{1}{\sqrt{2}} \phi(\frac{\cdot}{2}), \phi(\cdot - k) \rangle$$

$$\sqrt{2} \hat{\phi}(2\xi) = \hat{h}(\xi) \hat{\phi}(\xi), \quad \hat{h}(\xi) := \sum_{k \in \mathbb{Z}} h_k e^{-2\pi i k \xi}$$

$$\Rightarrow \hat{\phi}(2\xi) = \frac{1}{\sqrt{2}} \hat{h}(\xi) \hat{\phi}(\xi)$$

$$\text{i.e., } \hat{\phi}(\xi) = \frac{1}{\sqrt{2}} \hat{h}(\frac{\xi}{2}) \hat{\phi}(\frac{\xi}{2})$$

$$= \frac{1}{\sqrt{2}} \hat{h}(\frac{\xi}{2^2}) \hat{\phi}(\frac{\xi}{2^2})$$

$$\Rightarrow \hat{\phi}(\xi) = \prod_{p=1}^{\infty} \left(\frac{\hat{h}(2^{-p}\xi)}{\sqrt{2}} \right) \hat{\phi}(2^{-P}\xi) = \dots$$

If $\hat{\phi}(\xi)$ is continuous at $\xi = 0$,
 then $\hat{\phi}(2^{-P}\xi) \rightarrow \hat{\phi}(0)$ as $P \rightarrow \infty$
 $\Rightarrow \hat{\phi}(\xi) = \hat{\phi}(0) \prod_{p=1}^{\infty} \left(\frac{\hat{h}(2^{-p}\xi)}{\sqrt{2}} \right)$

Thm (Mallat & Meyer, 1986?)

Let $\phi \in L^2(\mathbb{R})$ be an integrable scaling fcn, i.e., $\int \phi(x) dx < \infty$, $\{\phi(x-k)\}_{k \in \mathbb{Z}}$: an ONB of V_0 .

necessary cond. $\Rightarrow |\hat{h}(\xi)|^2 + |\hat{h}(\xi + \frac{1}{2})|^2 \equiv 2$ a.e. $\xi \in \mathbb{R}$.
 and $\hat{h}(0) = \sqrt{2}$.

Conversely, if $\hat{h}(\xi)$ satisfies:

- sufficient cond. {
- 1) 1-periodic;
 - 2) C^1 in the neighborhood of $\xi = 0$; and
 - 3) $|\hat{h}(\xi)|^2 + |\hat{h}(\xi + \frac{1}{2})|^2 \equiv 2$, $\hat{h}(0) = \sqrt{2}$,
 $\inf_{\xi \in [-\frac{1}{4}, \frac{1}{4}]} |\hat{h}(\xi)| > 0$,

then $\hat{\phi}(\xi) = \prod_{p=1}^{\infty} \frac{\hat{h}(2^{-p}\xi)}{\sqrt{2}}$ is

the Fourier transform of a scaling fcn $\phi \in L^2(\mathbb{R})$.

(Proof) Here, we only prove the necessary cond. for the whole proof, see, e.g., Mallat's book.

$\{\phi(x-k)\}_{k \in \mathbb{Z}}$: an ONB for $V_0 \subset L^2(\mathbb{R})$.

The F.T. of the orthonormality gives as

(*) $\sum_{k \in \mathbb{Z}} |\hat{\phi}(\xi+k)|^2 \equiv 1$ a.e. $\xi \in \mathbb{R}$ as we did before.

By the two-scale diff. eqn. in the Fourier dom.

$$\hat{\phi}(\xi) = \frac{1}{\sqrt{2}} \hat{h}\left(\frac{\xi}{2}\right) \hat{\phi}\left(\frac{\xi}{2}\right), (*) \text{ becomes}$$

$$\sum_{k \in \mathbb{Z}} |\hat{h}\left(\frac{\xi}{2} + \frac{k}{2}\right)|^2 |\hat{\phi}\left(\frac{\xi}{2} + \frac{k}{2}\right)|^2 \equiv 2$$

$$\Leftrightarrow \sum_{l \in \mathbb{Z}} |\hat{h}\left(\frac{\xi}{2} + l\right)|^2 |\hat{\phi}\left(\frac{\xi}{2} + l\right)|^2$$

$$+ |\hat{h}\left(\frac{\xi}{2} + \frac{1}{2} + l\right)|^2 |\hat{\phi}\left(\frac{\xi}{2} + \frac{1}{2} + l\right)|^2 \equiv 2$$

$$\Leftrightarrow |\hat{h}\left(\frac{\xi}{2}\right)|^2 \sum_{l \in \mathbb{Z}} |\hat{\phi}\left(\frac{\xi}{2} + l\right)|^2 \xrightarrow{\text{via } (*)} \frac{1}{2}$$

$$+ |\hat{h}\left(\frac{\xi}{2} + \frac{1}{2}\right)|^2 \sum_{l \in \mathbb{Z}} |\hat{\phi}\left(\frac{\xi}{2} + \frac{1}{2} + l\right)|^2 \equiv 2$$

$$\Leftrightarrow |\hat{h}\left(\frac{\xi}{2}\right)|^2 + |\hat{h}\left(\frac{\xi}{2} + \frac{1}{2}\right)|^2 \equiv 2, \text{ a.e. } \xi \in \mathbb{R}$$

$$\Leftrightarrow |\hat{h}(\xi)|^2 + |\hat{h}\left(\xi + \frac{1}{2}\right)|^2 \equiv 2, \text{ a.e. } \xi \in \mathbb{R} \checkmark$$

Now, $\hat{\phi}(2\xi) = \frac{1}{\sqrt{2}} \hat{h}(\xi) \hat{\phi}(\xi)$ and set $\xi = 0$

$$\Rightarrow \hat{\phi}(0) = \frac{1}{\sqrt{2}} \hat{h}(0) \hat{\phi}(0) \Leftrightarrow \hat{h}(0) = \sqrt{2} \text{ since } \hat{\phi}(0) \neq 0.$$


Example 1 : Piecewise Const. MRA

$$\phi(x) = \chi_{[0,1)}(x)$$

$$h_k = \left\langle \frac{1}{\sqrt{2}} \chi_{[0,2)}, \chi_{[0,1)}(\cdot - k) \right\rangle$$

$$= \begin{cases} \frac{1}{\sqrt{2}} & k=0,1 \\ 0 & \text{o.w.} \end{cases}$$

(no overlap)




Example 2: Shannon MRA

$$\phi(x) = \text{sinc}(x), \quad \hat{\phi}(\xi) = \chi_{[-\frac{1}{2}, \frac{1}{2})}(\xi)$$

From the two-scale diff. eqn. in Fourier,

$$\hat{h}(\xi) = \frac{\sqrt{2} \hat{\phi}(2\xi)}{\hat{\phi}(\xi)} = \sqrt{2} \chi_{[-\frac{1}{4}, \frac{1}{4})}(\xi) \quad \text{for } \forall \xi \in [-\frac{1}{2}, \frac{1}{2})$$



$$\hat{h}(\xi) = \sum_{k \in \mathbb{Z}} h_k e^{-2\pi i k \xi} = \sqrt{2} \chi_{[-\frac{1}{4}, \frac{1}{4})}(\xi)$$

$$\begin{aligned} h_k &= \int_{-\frac{1}{2}}^{\frac{1}{2}} \sqrt{2} \chi_{[-\frac{1}{4}, \frac{1}{4})}(\xi) e^{+2\pi i k \xi} d\xi \\ &= \sqrt{2} \int_{-\frac{1}{4}}^{\frac{1}{4}} e^{2\pi i k \xi} d\xi = \frac{1}{\sqrt{2}} \frac{\sin \frac{\pi k}{2}}{\frac{\pi k}{2}} \\ &= \frac{1}{\sqrt{2}} \text{sinc}\left(\frac{k}{2}\right), \quad k \in \mathbb{Z}. \end{aligned}$$

$\Rightarrow \{h_k\}$: not a finite sequence.

Example 3: Spline MRA

Recall

$$\hat{\phi}(\xi) = \frac{e^{-i\varepsilon\pi\xi}}{\xi^{m+1} \sqrt{S_{2m+2}(\xi)}} \quad S_n(\xi) := \sum_{k \in \mathbb{Z}} (\xi+k)^{-n}$$

$$\hat{h}(\xi) = \frac{\sqrt{2} \hat{\phi}(2\xi)}{\hat{\phi}(\xi)} = e^{-i\varepsilon\pi\xi} \sqrt{\frac{S_{2m+2}(\xi)}{2^{2m+1} S_{2m+2}(2\xi)}}$$

$m=1$: linear case \Rightarrow



$$\text{Recall } S_4(\xi) = \frac{\pi^4}{3} \frac{1 + 2\cos^2 \pi \xi}{\sin^4 \pi \xi}.$$

$$\Rightarrow \hat{h}(\xi) = \sqrt{\frac{1}{2^3} \cdot \frac{1 + 2 \cos^2 \pi \xi}{1 + 2 \cos^2 2\pi \xi} \cdot \frac{\sin^4 2\pi \xi}{\sin^4 \pi \xi}}$$

$$= \sqrt{2} \sqrt{\frac{1 + 2 \cos^2 \pi \xi}{1 + 2 \cos^2 2\pi \xi} \cdot \cos^2 \pi \xi}$$

$$\stackrel{\text{green}}{=} \left(\frac{2 \sin \pi \xi \cdot \cos \pi \xi}{\sin \pi \xi} \right)^4$$

$\Rightarrow \{h_k\}$: numerical table
 spline scaling fun may be relatively localized in x but not compactly supported while $\theta(x)$ is compactly supported.




$H(\xi)$
 $= \hat{h}(\xi)$

★ Mother Wavelet; Wavelet ONB

Recall an MRA of $L^2(\mathbb{R})$

$$\dots \subset V_{j+1} \subset \underbrace{V_j}_{\text{wavy}} \subset V_{j-1} \subset \dots$$

Consider the **orthogonal complement** of V_j in V_{j-1} , i.e., the information contained in V_{j-1} but **not** in V_j .




$$V_j \oplus W_j = V_{j-1}$$

In terms of the orthogonal proj.'s, we can write
 $\forall f \in L^2(\mathbb{R})$,

$$\underbrace{P_{V_{j-1}} f}_{\substack{\text{approx.} \\ \text{at resol.} \\ 2^{-j+1}}} = \underbrace{P_{V_j} f}_{\substack{\text{approx.} \\ \text{at resol.} \\ 2^{-j}}} + \underbrace{P_{W_j} f}_{\substack{\text{detailed} \\ \text{info} \\ \text{necessary} \\ \text{to recover} \\ P_{V_{j-1}} f}}$$

The Concept of Multiresolution Analysis



Multiresolution Analysis by Sinc Wavelets

Multiresolution Decomposition with Haar Basis

In fact,
in this case $\rightarrow f$


$$f = P_{V_0} f + P_{W_1} f$$

$$P_{W_2} f$$

\vdots
 \vdots
 \vdots

$$P_{W_6} f$$

$$P_{V_6} f$$



Multiresolution Approximation with Haar Basis

f


$$P_{V_1} f$$

$$P_{V_2} f$$

\vdots
 \vdots

$$P_{V_6} f \Rightarrow P_{V_5} f$$

$$+ P_{W_6} f \quad P_{V_6} f$$



Father $\phi \rightarrow \phi_{j,k}$, $V_j = \overline{\text{span} \{ \phi_{j,k} \}_{k \in \mathbb{Z}}}$ ONB
 Mother $\psi \rightarrow \psi_{j,k}$, $W_j = \overline{\text{span} \{ \psi_{j,k} \}_{k \in \mathbb{Z}}}$ ONB

Thm (Mallat, Meyer 1986)

Let ϕ be a scaling fcn (father wavelet) and $\{h_k\}_{k \in \mathbb{Z}}$ be the corresponding CMF. Let us define $\psi \in L^2(\mathbb{R})$ whose Fourier transf. has

$$\hat{\psi}(\xi) = \frac{1}{\sqrt{2}} \hat{g}\left(\frac{\xi}{2}\right) \hat{\phi}\left(\frac{\xi}{2}\right)$$

with $\hat{g}(\xi) = e^{-2\pi i \xi} \overline{\hat{h}\left(\xi + \frac{1}{2}\right)}$.

Let $\psi_{j,k}(x) = 2^{-j/2} \psi(2^{-j}x - k)$.

Then, $\{\psi_{j,k}\}_{k \in \mathbb{Z}}$ form an ONB of W_j for each $j \in \mathbb{Z}$, and $\{\psi_{j,k}\}_{(j,k) \in \mathbb{Z}^2}$ form an ONB of $L^2(\mathbb{R})$.

(Proof) We look for a fcn $\psi \in L^2(\mathbb{R})$ s.t.

$$\frac{1}{\sqrt{2}} \psi\left(\frac{x}{2}\right) = \psi_{1,0}(x) \in W_1 \subset V_0$$

and $\{\psi_{1,k}\}_{k \in \mathbb{Z}}$ form an ONB of W_1 .

Suppose $\frac{1}{\sqrt{2}} \psi\left(\frac{x}{2}\right) \in W_1$. Since $W_1 \subset V_0$ and $\{\phi(x-k)\}_{k \in \mathbb{Z}}$: an ONB of V_0 ,

$$\frac{1}{\sqrt{2}} \psi\left(\frac{x}{2}\right) = \sum_{k \in \mathbb{Z}} \underbrace{\left\langle \frac{1}{\sqrt{2}} \psi\left(\frac{\cdot}{2}\right), \phi(\cdot - k) \right\rangle}_{=: g_k} \phi(x-k)$$

$$\sqrt{2} \hat{\psi}(2\xi) = \hat{g}(\xi) \hat{\phi}(\xi), \quad \hat{g}(\xi) = \sum_{k \in \mathbb{Z}} g_k e^{-2\pi i k \xi}$$

Lemma The family $\{\psi_{j,k}\}_{k \in \mathbb{Z}}$ is an ONB of W_j

$$\iff \begin{cases} |\hat{g}(\xi)|^2 + |\hat{g}(\xi + \frac{1}{2})|^2 \equiv 2 & \text{a.e. } \xi \in \mathbb{R} \\ \hat{g}(\xi) \overline{\hat{h}(\xi)} + \hat{g}(\xi + \frac{1}{2}) \overline{\hat{h}(\xi + \frac{1}{2})} \equiv 0 \end{cases}$$

(Proof of Lemma) We'll prove only $j=0$ case since the other cases are easy via S_{2^j} op. once we prove the $j=0$ case.

Using the same argument in the proof of $\{\phi(x-k)\}_{k \in \mathbb{Z}}$ forming an ONB of V_0 , we can show that

$\{\psi(x-k)\}_{k \in \mathbb{Z}}$ are orthonormal

$$\iff I(\xi) := \sum_{k \in \mathbb{Z}} |\hat{\psi}(\xi+k)|^2 \equiv 1 \quad \text{a.e. } \xi \in \mathbb{R}$$

Now, the two-scale diff. eqn. $\hat{\psi}(\xi) = \frac{1}{\sqrt{2}} \hat{g}(\frac{\xi}{2}) \hat{\phi}(\frac{\xi}{2})$

$$I(\xi) = \frac{1}{2} \sum_k |\hat{g}(\frac{\xi}{2} + \frac{k}{2})|^2 |\hat{\phi}(\frac{\xi}{2} + \frac{k}{2})|^2 \quad \hat{g}: 1\text{-periodic}$$

$$= \frac{1}{2} \sum_l (|\hat{g}(\frac{\xi}{2} + l)|^2 |\hat{\phi}(\frac{\xi}{2} + l)|^2$$

$$+ |\hat{g}(\frac{\xi}{2} + l + \frac{1}{2})|^2 |\hat{\phi}(\frac{\xi}{2} + l + \frac{1}{2})|^2)$$

$\hat{g}: 1\text{-periodic}$

$$\stackrel{\downarrow}{=} \frac{1}{2} \left\{ |\hat{g}(\frac{\xi}{2})|^2 \sum_l |\hat{\phi}(\frac{\xi}{2} + l)|^2 + |\hat{g}(\frac{\xi}{2} + \frac{1}{2})|^2 \sum_l |\hat{\phi}(\frac{\xi}{2} + l + \frac{1}{2})|^2 \right\}$$

$$= \frac{1}{2} (|\hat{g}(\frac{\xi}{2})|^2 + |\hat{g}(\frac{\xi}{2} + \frac{1}{2})|^2) \equiv 1 \quad \text{a.e. } \xi \in \mathbb{R}$$

$$\iff |\hat{g}(\xi)|^2 + |\hat{g}(\xi + \frac{1}{2})|^2 \equiv 2 \quad \text{a.e. } \xi \in \mathbb{R}.$$

Finally, we need to show $V_{-1} = V_0 \oplus W_0$.
 We know $\{\sqrt{2} \phi(2x-k)\}_{k \in \mathbb{Z}}$ form an ONB of V_{-1} .

So, $V_{-1} = V_0 \oplus W_0$

$\Leftrightarrow \forall \{a_k\} \in \ell^2(\mathbb{Z}), \exists \{b_k\}, \{c_k\} \in \ell^2(\mathbb{Z})$ s.t.

$$\sum a_k \sqrt{2} \phi(2(x - \frac{k}{2})) = \sum b_k \phi(x-k) + \sum c_k \psi(x-k)$$

$\downarrow \mathcal{F}$

$$\frac{1}{\sqrt{2}} \hat{a}(\frac{\xi}{2}) \hat{\phi}(\frac{\xi}{2}) = \hat{b}(\xi) \hat{\phi}(\xi) + \hat{c}(\xi) \hat{\psi}(\xi)$$

$$\Leftrightarrow \hat{a}(\frac{\xi}{2}) = \hat{b}(\xi) \hat{h}(\frac{\xi}{2}) + \hat{c}(\xi) \hat{g}(\frac{\xi}{2}) \quad (*)$$

\uparrow via $\hat{\phi}(\xi) = \frac{1}{\sqrt{2}} \hat{h}(\frac{\xi}{2}) \hat{\phi}(\frac{\xi}{2}), \hat{\psi}(\xi) = \frac{1}{\sqrt{2}} \hat{g}(\frac{\xi}{2}) \hat{\phi}(\frac{\xi}{2})$

Question: Do such $\{b_k\}, \{c_k\}$ exist?

\Rightarrow Yes!

Define $\hat{b}(2\xi) := \frac{1}{2} [\hat{a}(\xi) \overline{\hat{h}(\xi)} + \hat{a}(\xi + \frac{1}{2}) \overline{\hat{h}(\xi + \frac{1}{2})}]$

$$\hat{c}(2\xi) := \frac{1}{2} [\hat{a}(\xi) \overline{\hat{g}(\xi)} + \hat{a}(\xi + \frac{1}{2}) \overline{\hat{g}(\xi + \frac{1}{2})}]$$

Then these satisfy (*).

In fact,

$$\begin{cases} \hat{b}(\xi) \hat{h}(\frac{\xi}{2}) = \frac{1}{2} [\hat{a}(\frac{\xi}{2}) |\hat{h}(\frac{\xi}{2})|^2 + \hat{a}(\xi + \frac{1}{2}) \hat{h}(\frac{\xi}{2}) \overline{\hat{h}(\xi + \frac{1}{2})}] \\ \hat{c}(\xi) \hat{g}(\frac{\xi}{2}) = \frac{1}{2} [\hat{a}(\frac{\xi}{2}) |\hat{g}(\frac{\xi}{2})|^2 + \hat{a}(\xi + \frac{1}{2}) \hat{g}(\frac{\xi}{2}) \overline{\hat{g}(\xi + \frac{1}{2})}] \end{cases}$$

We can show that $|\hat{h}(\frac{\xi}{2})|^2 + |\hat{g}(\frac{\xi}{2})|^2 \equiv \frac{2}{a.e.} \xi \in \mathbb{R}$
 and $\hat{h}(\frac{\xi}{2}) \overline{\hat{h}(\xi + \frac{1}{2})} + \hat{g}(\frac{\xi}{2}) \overline{\hat{g}(\xi + \frac{1}{2})} \equiv 0$

These can be derived from

$$\begin{cases} |\hat{h}(\frac{\xi}{2})|^2 + |\hat{h}(\xi + \frac{1}{2})|^2 \equiv 2 \\ |\hat{g}(\frac{\xi}{2})|^2 + |\hat{g}(\xi + \frac{1}{2})|^2 \equiv 2 \\ \hat{h}(\frac{\xi}{2})\overline{\hat{g}(\frac{\xi}{2})} + \hat{h}(\xi + \frac{1}{2})\overline{\hat{g}(\xi + \frac{1}{2})} \equiv 0 \end{cases} \quad \text{a.e. } \xi \in \mathbb{R}$$

Hence such $\hat{b}(\xi), \hat{c}(\xi)$ exist.

They are 1-periodic because of their forms
and $\hat{a}, \hat{h}, \hat{g}$ are also 1-periodic

Thus $\exists \{b_k\}, \{c_k\} \in \ell^2(\mathbb{Z})$

i.e., $V_{-1} = V_0 \oplus W_0$!

$$\Leftrightarrow W_0 = V_0^\perp \text{ in } V_{-1}$$

$$\Rightarrow W_j = V_j^\perp \text{ in } V_{j-1}, \quad V_{j-1} = V_j \oplus W_j$$

Lemma done. // //

Now,

$$W_j \perp V_j, \quad W_l \subset V_{l-1} \subset V_j \quad \forall l > j$$

$$\Rightarrow W_j \perp W_l. \quad \text{Hence } L^2(\mathbb{R}) = \bigoplus_{j=-\infty}^{\infty} W_j$$

and for any $L > J$,

$$\begin{aligned} V_J &= V_L \oplus W_L \oplus W_{L-1} \oplus \dots \oplus W_{J-1} \\ &= V_L \oplus \bigoplus_{j=L}^{J-1} W_j \end{aligned}$$

Thm done. // //