

## Homework 3

due February 17

**Problem 1.** Let  $C(x, y) = \prod_{i,j \geq 1} (1 - x_i y_j)^{-1}$ . Show that then for all  $f \in \Lambda$

$$\langle C(x, y), f(x) \rangle = f(y)$$

where the scalar product is taken in the  $x$  variables. In other words,  $C(x, y)$  is a *reproducing kernel* for the scalar product.

**Problem 2.** Show that

$$\prod_{i=1}^m \prod_{j=1}^n (x_i + y_j) = \sum_{\lambda} s_{\lambda}(x) s_{\tilde{\lambda}}(y)$$

where the sum is over all partitions  $\lambda$  with  $\ell(\lambda) \leq m$  and  $\lambda_1 \leq n$  and  $\tilde{\lambda}$  is defined as

$$\tilde{\lambda} = (n - \lambda_m, n - \lambda_{m-1}, \dots, n - \lambda_1).$$

**Problem 3.** Let  $D_{\mu} : \Lambda \rightarrow \Lambda$  be the linear transformation given by

$$D_{\mu}(s_{\lambda}) = s_{\lambda/\mu}.$$

Show that  $D_{\mu} D_{\lambda} = D_{\lambda} D_{\mu}$ .

**Problem 4.** If  $R$  is a ring, then an additive group homomorphism  $D : R \rightarrow R$  is called a derivation if  $D(fg) = (Df)g + f(Dg)$  for all  $f, g \in R$ .

- (1) Show that the linear transformation  $\Lambda \rightarrow \Lambda$  defined by  $D(s_{\lambda}) = s_{\lambda/1}$  is a derivation.
- (2) Show that the bilinear operation  $[f, g]$  on  $\Lambda$  given by

$$[s_{\lambda}, s_{\mu}] = s_{\lambda/1} s_{\mu} - s_{\lambda} s_{\mu/1}$$

defines a Lie algebra structure on  $\Lambda$ . That is, the Jacobi identity holds

$$[f, [g, h]] + [g, [h, f]] + [h, [f, g]] = 0.$$

**Problem 5.** If  $A$  is an algebra over a field  $k$ , then  $A \otimes_k A$  is an algebra with multiplication characterized uniquely by  $(a \otimes b)(c \otimes d) = ac \otimes bd$ . A coproduct is an algebra homomorphism  $\Delta : A \rightarrow A \otimes A$  which is coassociative in the sense that the two maps

$$(1 \otimes \Delta) \circ \Delta \quad \text{and} \quad (\Delta \otimes 1) \circ \Delta$$

from  $A$  to  $A \otimes A \otimes A$  are equal. (This axiom is dual to the associative law for multiplication  $\mu : A \otimes A \rightarrow A$ , which can be formulated as  $\mu \circ (1 \otimes \mu) = \mu \circ (\mu \otimes 1)$ .) Here  $1 : A \rightarrow A$  is the identity map.

Taking  $\Lambda$  to be the algebra of symmetric functions with coefficients in  $k = \mathbb{Q}$  and identifying  $\Lambda \otimes \Lambda$  with  $\Lambda(X)\Lambda(Y)$ , show that  $\Delta(f) = f(X, Y)$  defines a coproduct on  $\Lambda$ .

**Remark:** An algebra equipped with a coproduct is called a bialgebra. If we also define the counit  $\epsilon : \Lambda \rightarrow \mathbb{Q}$  by  $\epsilon(f) = \langle f, 1 \rangle = f(0, 0, \dots)$  and the antipode  $S : \Lambda \rightarrow \Lambda$  by  $Sf = f(-X)$ , these together with  $\Delta$  can be shown to satisfy the axioms of a Hopf algebra.

**Problem 6.** The relevance of this problem will become clear when we talk about differential posets in class.

- (1) Let  $U : \Lambda \rightarrow \Lambda$  and  $D : \Lambda \rightarrow \Lambda$  be linear transformations defined by

$$U(f) = p_1 f$$

$$D(f) = \frac{\partial}{\partial p_1} f$$

where  $\partial/\partial p_1$  is applied to  $f$  written as a polynomial in the  $p_i$ 's. Show that

$$DU - UD = I$$

$$DU^k = kU^{k-1} + U^k D$$

where  $I$  is the identity operator.

- (2) Now let us work in Young's lattice  $Y$  and define

$$D(\lambda) = \sum_{\lambda^- \triangleleft \lambda} \lambda^-$$

$$U(\lambda) = \sum_{\lambda^+ \triangleright \lambda} \lambda^+$$

where the sum over  $\lambda^-$  is over all partitions being covered by  $\lambda$  in  $Y$  and the sum over  $\lambda^+$  is over all partitions covering  $\lambda$  in  $Y$ . Show that

$$DU - UD = I$$

$$DU^k = kU^{k-1} + U^k D.$$

- (3) Derive from (2) that

$$\sum_{\lambda \vdash n} (f^\lambda)^2 = n!$$