

1.) Use what you know about converging geometric series to write each power series as an ordinary function.

a.)  $\sum_{n=2}^{\infty} \frac{x^n}{3^n}$       b.)  $\sum_{n=0}^{\infty} \frac{2^{n-1}(x+3)^{n+1}}{5^n}$   
 c.)  $x^2 - x^{5/2} + x^3 - x^{7/2} + x^4 - x^{9/2} + \dots$       d.)  $\sum_{n=0}^{\infty} (n+1)x^n$

2.) Recall that if  $y = f(x)$  is a function and

$$a_0 + a_1(x-a) + a_2(x-a)^2 + a_3(x-a)^3 + a_4(x-a)^4 + \dots = \sum_{n=0}^{\infty} a_n(x-a)^n$$

is the Taylor Series (or Maclaurin series if  $a = 0$ ) centered at  $x = a$  for  $y = f(x)$ , then  $a_n = \frac{f^{(n)}(a)}{n!}$ . Use this formula to compute the first four nonzero terms and the general formula for the Taylor series expansion for each function about the given value of  $a$ .

a.)  $f(x) = e^x$  centered at  $x = 0$       b.)  $f(x) = e^x$  centered at  $x = \ln 2$   
 c.)  $f(x) = \frac{1}{1-x}$  centered at  $x = 0$       d.)  $f(x) = \sin x$  centered at  $x = 0$   
 e.)  $f(x) = \frac{1}{x}$  centered at  $x = 1$       f.)  $f(x) = \sqrt{x+5}$  centered at  $x = -1$

3.) Use the suggested method to find the first four nonzero terms of the Maclaurin series for each function.

a.)  $f(x) = \frac{1}{1+x^2}$  (Substitute  $-x^2$  into the Maclaurin series for  $\frac{1}{1-x}$ .)  
 b.)  $f(x) = x^3 e^{-3x}$  (Substitute  $-3x$  into the Maclaurin series for  $e^x$  and then multiply by  $x^3$ .)  
 c.)  $f(x) = \frac{e^x}{1-x} = e^x \frac{1}{1-x}$  (Multiply the Maclaurin series for  $e^x$  and  $\frac{1}{1-x}$  term by term and then group like powers of  $x$ .)  
 d.)  $f(x) = \frac{e^x}{1-x}$  (Use polynomial division. Divide the Maclaurin series for  $e^x$  by  $1-x$ .)  
 e.)  $f(x) = 3x^2 \cos(x^3)$  (Substitute  $x^3$  into the Maclaurin series for  $\sin x$  then differentiate term by term.)  
 f.)  $f(x) = \arctan x$  (Integrate the Maclaurin series for  $\frac{1}{1+t^2}$  from 0 to  $x$ .)

4.) The Maclaurin series for  $f(x) = \frac{1}{1+x}$  is  $1 - x + x^2 - x^3 + x^4 - x^5 + \dots$

a.) Show that  $f(x) = \frac{1}{1+x}$  and  $1 - x + x^2 - x^3 + x^4 - x^5 + \dots$  have the same value at  $x = 0$ .

b.) Show that  $f(x) = \frac{1}{1+x}$  and  $1 - x + x^2 - x^3 + x^4 - x^5 + \dots$  have the same value at  $x = 1/2$ .

c.) Show that  $f(x) = \frac{1}{1+x}$  and  $1 - x + x^2 - x^3 + x^4 - x^5 + \dots$  do not have the same value at  $x = 1$ .

d.) For what x-values is  $f(x) = \frac{1}{1+x}$  defined?

e.) For what x-values is the Maclaurin series  $1 - x + x^2 - x^3 + x^4 - x^5 + \dots$  defined?

NOTE: It can be shown that  $f(x) = \frac{1}{1+x}$  and its Maclaurin series  $1 - x + x^2 - x^3 + x^4 - x^5 + \dots$  are equal on the interval  $(-1, 1)$ .

5.) Determine (Use shortcuts.) the third-degree Taylor polynomial,  $P_3(x; 0)$ , for the function  $f(x) = \frac{x}{1+x}$ . Use  $\int_0^1 P_3(x; 0) dx$  to estimate the value of  $\int_0^1 \frac{x}{1+x} dx$ . Now evaluate  $\int_0^1 \frac{x}{1+x} dx$  directly to see how good the estimate is.

6.) The following definite integral cannot be evaluated using the Fundamental Theorem of Calculus. Use the Maclaurin series for  $\cos x$  and the absolute error  $|R_n|$  for an alternating series to estimate the value of this integral with error at most 0.0001:  $\int_0^1 \cos(x^2) dx$

7.) Write each Maclaurin series as an ordinary function.

a.)  $(3x) - \frac{(3x)^3}{3!} + \frac{(3x)^5}{5!} - \frac{(3x)^7}{7!} + \frac{(3x)^9}{9!} - \dots$  (HINT: Use  $\sin x$ .)

b.)  $x^2 - x^3 + x^4 - x^5 + x^6 - \dots$  (HINT: Factor.)

c.)  $\frac{1}{2!} + \frac{x}{3!} + \frac{x^2}{4!} + \frac{x^3}{5!} + \frac{x^4}{6!} + \dots$  (HINT: Use  $e^x$ .)

d.)  $x + 2x^2 + 3x^3 + 4x^4 + 5x^5 + \dots$  (Challenging)

8.) Use any method to find the given Taylor polynomial for each function. Then estimate the Absolute Taylor Error on the indicated interval.

a.)  $f(x) = e^{-2x}$ ,  $P_3(x; 0)$ , for  $[-1/2, 1/3]$

b.)  $f(x) = \sin 2x$ ,  $P_5(x; 0)$ , for  $[0, 3/4]$

c.)  $f(x) = \frac{x}{1-x}$ ,  $P_4(x; 0)$ , for  $[-1/3, 0]$

9.) What should  $n$  be so that the  $n$ th-degree Taylor Polynomial  $P_n(x; a)$  estimates the

value of the given function on the indicated interval with Absolute Taylor Error at most 0.00001 ?

a.)  $f(x) = e^{-x}$  for  $a = 1$  and  $[0, 1]$

b.)  $f(x) = \frac{x+3}{x+1}$  for  $a = 0$  and  $[0, 1/2]$

10.) (Challenging) Use shortcuts to find the first three nonzero terms in the Taylor Series centered at  $x = -1$  for  $f(x) = \frac{x}{3-x}$ .

“In mathematics you don’t understand things. You just get used to them.” – Johann von Neumann