

# 15 FACE NUMBERS OF POLYTOPES AND COMPLEXES

Louis J. Billera and Anders Björner

57805  
=

---

## INTRODUCTION

Geometric objects are often put together from simple pieces according to certain combinatorial rules. As such, they can be described as *complexes* with their constituent *cells*, which are usually polytopes and often simplices. Many constraints of a combinatorial and topological nature govern the incidence structure of cell complexes and are therefore relevant in the analysis of geometric objects. Since these incidence structures are in most cases too complicated to be well understood, it is worthwhile to focus on simpler invariants that still say something nontrivial about their combinatorial structure. The invariants to be discussed in this chapter are the *f-vectors*  $f = (f_0, f_1, \dots)$ , where  $f_i$  is the number of  $i$ -dimensional cells in the complex.

The theory of *f-vectors* can be discussed at two levels: (1) the numerical relations satisfied by the  $f_i$  numbers, and (2) the algebraic, combinatorial, and topological facts and constructions that give rise to and explain these relations. This chapter will summarize the main facts in the numerology of *f-vectors* (i.e., at level 1), with emphasis on cases of geometric interest.

The chapter is organized as follows. To begin with, we treat simplicial complexes, first the general case (Section 15.1), then complexes with various Betti number constraints (Section 15.2), and finally triangulations of spheres, polytope boundaries, and manifolds (Section 15.3). Then we move on to nonsimplicial complexes, discussing first the general case (Section 15.4) and then polytopes and spheres (Section 15.5).

---

## 15.1 SIMPLICIAL COMPLEXES

---

### GLOSSARY

The convex hull of any set of  $j + 1$  affinely independent points in  $\mathbb{R}^n$  is called a *j-simplex*. See Chapter 13 for more about this definition, and for the notions of *faces* and *vertices* of a simplex.

A *geometric simplicial complex*  $\Gamma$  is a finite nonempty family of simplices in  $\mathbb{R}^n$  such that (i)  $\sigma \in \Gamma$  implies that  $\tau \in \Gamma$  for every face  $\tau$  of  $\sigma$ , and (ii) if  $\sigma, \tau \in \Gamma$  and  $\sigma \cap \tau \neq \emptyset$  then  $\sigma \cap \tau$  is a face of both  $\sigma$  and  $\tau$ .

An *abstract simplicial complex*  $\Delta$  is a finite nonempty family of subsets of

some ground set  $V$  (the *vertex set*) such that if  $A \in \Delta$  and  $B \subseteq A$  then  $B \in \Delta$ . (Note that always  $\emptyset \in \Delta$ .) The elements  $A \in \Delta$  are called *faces*. Define the *dimension* of a face  $A$  and of  $\Delta$  itself by  $\dim A = |A| - 1$ ;  $\dim \Delta = \max_{A \in \Delta} \dim A$ . By a *d-complex* we mean a  $d$ -dimensional complex.

With every geometric simplicial complex  $\Gamma$  we associate an abstract simplicial complex by taking the family of vertex sets of its simplices. Conversely, every  $d$ -dimensional abstract simplicial complex  $\Delta$  can be realized in  $\mathbb{R}^n$  for  $n \geq 2d + 1$  (and sometimes less) by some geometric simplicial complex. The latter is unique up to homeomorphism, so it is correct to think of the realization map as a one-to-one correspondence between abstract and geometric simplicial complexes. We will therefore drop the adjectives “abstract” and “geometric” and speak only of a *simplicial complex*.

For a simplicial complex  $\Delta$ , let  $\Delta^i = \{i\text{-dimensional faces}\}$  and let  $f_i = |\Delta^i|$ . The integer sequence  $f(\Delta) = (f_0, f_1, \dots)$  is called the *f-vector* of  $\Delta$ . (The entry  $f_{-1} = 1$  is usually suppressed.) The subcomplex  $\Delta^{\leq i} = \bigcup_{j \leq i} \Delta^j$  is called the *i-skeleton* of  $\Delta$ .

A simplicial complex  $\Delta$  is called *pure* if all maximal faces are of equal dimension. It is called *r-colorable* if there exists a partition of the vertex set  $V = V_1 \cup \dots \cup V_r$  such that  $|A \cap V_i| \leq 1$  for all  $A \in \Delta$  and  $1 \leq i \leq r$ . Equivalently,  $\Delta$  is *r-colorable* if and only if its 1-skeleton  $\Delta^{\leq 1}$  is *r-colorable* in the standard sense of graph theory. An  $(r-1)$ -complex that is both pure and *r-colorable* is sometimes called *balanced*.

For integers  $k, n \geq 1$  there is a unique way of writing

$$n = \binom{a_k}{k} + \binom{a_{k-1}}{k-1} + \dots + \binom{a_i}{i}$$

so that  $a_k > a_{k-1} > \dots > a_i \geq i \geq 1$ . Then define

$$\partial_k(n) = \binom{a_k}{k-1} + \binom{a_{k-1}}{k-2} + \dots + \binom{a_i}{i-1},$$

and

$$\partial^k(n) = \binom{a_k-1}{k-1} + \binom{a_{k-1}-1}{k-2} + \dots + \binom{a_i-1}{i-1}.$$

Also let  $\partial_k(0) = \partial^k(0) = 0$ .

Let  $\mathbb{N}^\infty$  denote the set of sequences  $(n_0, n_1, \dots)$  of nonnegative integers, and  $\mathbb{N}^{(\infty)}$  the subset of sequences such that  $n_k = 0$  for sufficiently large  $k$ . We call  $n \in \mathbb{N}^{(\infty)}$  a *K-sequence* if

$$\partial_{k+1}(n_k) \leq n_{k-1} \quad \text{for all } k \geq 1.$$

We call  $n \in \mathbb{N}^\infty$  an *M-sequence* if

$$n_0 = 1 \text{ and } \partial^k(n_k) \leq n_{k-1} \quad \text{for all } k \geq 2.$$

---

## THE KRUSKAL-KATONA THEOREM AND SOME RELATIVES

The following basic result characterizes the *f*-vectors of simplicial complexes.

**THEOREM 15.1.1** *Kruskal-Katona Theorem*

For  $f = (f_0, f_1, \dots) \in \mathbb{N}^{(\infty)}$  the following are equivalent:

- (i)  $f$  is the  $f$ -vector of a simplicial complex;
- (ii)  $f$  is a  $K$ -sequence.

A simplicial complex is **connected** if its 1-skeleton is connected in the sense of graph theory.

**THEOREM 15.1.2**

For  $f \in \mathbb{N}^{(\infty)}$  the following are equivalent:

- (i)  $f$  is the  $f$ -vector of a connected simplicial complex;
- (ii)  $f$  is a  $K$ -sequence and  $\partial^3(f_2) \leq f_1 - f_0 + 1$ .

Theorem 15.1.1 has a generalization to colored complexes, whose statement will require some additional definitions. Fix an integer  $r > 0$ . Then define  $\binom{n}{k}_r$  as follows: partition  $\{1, \dots, n\}$  into  $r$  subsets  $V_1, \dots, V_r$  as evenly as possible (so every subset  $V_i$  will have  $\lfloor \frac{n}{r} \rfloor$  or  $\lfloor \frac{n}{r} \rfloor + 1$  elements), and let  $\binom{n}{k}_r$  be the number of  $k$ -subsets  $F \subseteq \{1, \dots, n\}$  such that  $|F \cap V_i| \leq 1$  for  $1 \leq i \leq r$ . For  $k \leq r$  every positive integer  $n$  can be uniquely written

$$n = \binom{a_k}{k}_r + \binom{a_{k-1}}{k-1}_{r-1} + \dots + \binom{a_i}{i}_{r-k+i},$$

where  $a_k > a_{k-1} > \dots > a_i \geq i \geq 1$ . Then define

$$\partial_k^{(r)}(n) = \binom{a_k}{k-1}_r + \binom{a_{k-1}}{k-2}_{r-1} + \dots + \binom{a_i}{i-1}_{r-k+i},$$

and let  $\partial_k^{(r)}(0) = 0$ .

**THEOREM 15.1.3**

For  $f = (f_0, \dots, f_{d-1})$ ,  $d \leq r$ , the following are equivalent:

- (i)  $f$  is the  $f$ -vector of an  $r$ -colorable simplicial complex;
- (ii)  $\partial_{k+1}^{(r)}(f_k) \leq f_{k-1}$ , for all  $1 \leq k \leq d-1$ .

Note that for  $r$  sufficiently large Theorem 15.1.3 specializes to Theorem 15.1.1.

---

**MULTICOMPLEXES AND MACAULAY'S THEOREM**

A **multicomplex**  $\mathcal{M}$  is a nonempty collection of monomials in finitely many variables such that if  $m$  is in  $\mathcal{M}$  then so is every divisor of  $m$ . Let  $f_i(\mathcal{M})$  be the number of degree  $i$  monomials in  $\mathcal{M}$ ;  $f(\mathcal{M}) = (f_0, f_1, \dots)$  is called the  **$f$ -vector** of  $\mathcal{M}$ .

**THEOREM 15.1.4** *Macaulay's Theorem*

For  $f \in \mathbb{N}^\infty$  the following are equivalent:

- (i)  $f$  is the  $f$ -vector of a multicomplex;

- (ii)  $f$  is an  $M$ -sequence;
- (iii)  $f_i = \dim_{\mathbf{k}} R_i$ ,  $i \geq 0$ , for some finitely generated commutative graded  $\mathbf{k}$ -algebra  $R = \bigoplus_{i \geq 0} R_i$  such that  $R_0 \cong \mathbf{k}$  (a field) and  $R_1$  generates  $R$ .

A simplicial complex can be identified with a multicomplex of squarefree monomials. Hence, a  $K$ -sequence is (except for a shift in the indexing) an  $M$ -sequence: If  $(f_0, \dots, f_{d-1})$  is a  $K$ -sequence then  $(1, f_0, \dots, f_{d-1})$  is an  $M$ -sequence. For this reason (and others, see, e.g., Theorem 15.2.2), properties of  $M$ -sequences are relevant also even if one is interested only in the special case of simplicial complexes.

A multicomplex is **pure** if all its maximal (under divisibility) monomials have the same degree.

### THEOREM 15.1.5

Let  $(f_0, \dots, f_r)$  be the  $f$ -vector of a pure multicomplex,  $f_r \neq 0$ . Then  $f_i \leq f_j$  for all  $i < j \leq r - i$ .

---

## COMMENTS

Simplicial complexes (abstract and geometric) are treated in most books on algebraic topology; see, e.g., [Mun84, Spa66]. The Kruskal-Katona theorem (independently discovered by M.-P. Schützenberger, J.B. Kruskal, G.O.H. Katona, L.H. Harper, and B. Lindström during the years 1959-1966) is discussed in many places and several proofs have appeared; see, e.g., [And87, Zie95]. Theorems 15.1.2 and 15.1.3 are from [Bjö96] and [FFK88] respectively. A Kruskal-Katona type theorem for simplicial complexes with vertex-transitive symmetry group appears in [FK96].

For Macaulay's theorem we refer to [And87, Sta96]. There is a common generalization of Macaulay's theorem and the Kruskal-Katona theorem due to Clements and Lindström; see [And87]. Theorem 15.1.5 is from [Hib89].

---

## 15.2 BETTI NUMBER CONSTRAINTS

---

### GLOSSARY

The **Euler characteristic**  $\chi(\Delta)$  of a simplicial complex  $\Delta$  with  $f$ -vector  $(f_0, \dots, f_{d-1})$  is  $\chi(\Delta) = \sum_{i=0}^{d-1} (-1)^i f_i$ .

The  **$h$ -vector**  $(h_0, \dots, h_d)$  of a  $(d-1)$ -dimensional simplicial complex is defined by

$$\sum_{i=0}^d h_i x^{d-i} = \sum_{i=0}^d f_{i-1} (x-1)^{d-i}.$$

The corresponding  **$g$ -vector**  $(g_0, \dots, g_{\lfloor d/2 \rfloor})$  is defined by  $g_0 = 1$  and  $g_i = h_i - h_{i-1}$ , for  $i \geq 1$ .

The **Betti number**  $\beta_i(\Delta)$  is the dimension (as a  $\mathbb{Q}$ -vector space) of the  $i$ -th reduced simplicial homology group  $\tilde{H}_i(\Delta, \mathbb{Q})$ ; see any textbook on algebraic topology

(e.g., [Mun84]) for the definition. We call  $(\beta_0, \dots, \beta_{\dim \Delta})$  the **Betti sequence** of  $\Delta$ .

The **link**  $\ell k_{\Delta}(F)$  of a face  $F$  is the subcomplex of  $\Delta$  defined by  $\ell k_{\Delta}(F) = \{A \in \Delta \mid A \cap F = \emptyset, A \cup F \in \Delta\}$ . Note that  $\ell k_{\Delta}(\emptyset) = \Delta$ .

A simplicial complex  $\Delta$  is **acyclic** if  $\beta_i(\Delta) = 0$  for all  $i$ .

A simplicial complex  $\Delta$  is **Cohen-Macaulay** if  $\beta_i(\ell k_{\Delta}(F)) = 0$  for all  $F \in \Delta$  and all  $i < \dim \ell k_{\Delta}(F)$ .

A simplicial complex  $\Delta$  is  **$m$ -Leray** if  $\beta_i(\ell k_{\Delta}(F)) = 0$  for all  $F \in \Delta$  and all  $i \geq m$ .

## FIXED BETTI NUMBERS

The most basic relationship between  $f$ -vectors and Betti numbers is the **Euler-Poincaré formula**:

$$\chi(\Delta) = f_0 - f_1 + f_2 - \dots = 1 + \beta_0 - \beta_1 + \beta_2 - \dots$$

This is in fact the only linear one in the following complete set of relations.

### THEOREM 15.2.1

For  $f = (f_0, f_1, \dots) \in \mathbb{N}^{(\infty)}$  and  $\beta = (\beta_0, \beta_1, \dots) \in \mathbb{N}^{(\infty)}$  the following are equivalent:

- (i)  $f$  is the  $f$ -vector of some simplicial complex with Betti sequence  $\beta$ ;
- (ii) if  $\chi_{k-1} = \sum_{j \geq k} (-1)^{j-k} (f_j - \beta_j)$ ,  $k \geq 0$ , then  $\chi_{-1} = 1$  and  $\partial_{k+1}(\chi_k + \beta_k) \leq \chi_{k-1}$  for all  $k \geq 1$ .

By putting  $\beta_i = 0$  for all  $i$  one gets as a special case a characterization of the  $f$ -vectors of acyclic simplicial complexes, viz.,  $\sum_{i \geq 0} f_{i-1} x^i = (1+x) \sum_{i \geq 0} f'_i x^i$ , where  $(f'_0, f'_1, \dots)$  is a  $K$ -sequence.

## COHEN-MACAULAY COMPLEXES

Examples of Cohen-Macaulay complexes are triangulations of manifolds whose Betti numbers vanish below the top dimension, in particular triangulations of spheres and balls. Other examples are matroid complexes (the independent sets of a matroid), Tits buildings, and the order complexes (simplicial complex of totally ordered subsets) of several classes of posets, e.g., semimodular lattices (including distributive and geometric lattices). Shellable complexes (see Chapters 14 and 17) are Cohen-Macaulay. Cohen-Macaulay complexes are always pure.

The definition of  $h$ -vector given in the glossary shows that the  $h$ -vector and the  $f$ -vector of a complex mutually determine each other via the formulas:

$$h_i = \sum_{j=0}^i (-1)^{i-j} \binom{d-j}{i-j} f_{j-1}, \quad f_{i-1} = \sum_{j=0}^i \binom{d-j}{i-j} h_j,$$

for  $0 \leq i \leq d$ . Hence, we may state  $f$ -vector results in terms of  $h$ -vectors whenever convenient.

**THEOREM 15.2.2**

For  $h = (h_0, \dots, h_d) \in \mathbb{Z}^{d+1}$  the following are equivalent:

- (i)  $h$  is the  $h$ -vector of a  $(d-1)$ -dimensional Cohen-Macaulay complex;
- (ii)  $h$  is the  $h$ -vector of a  $(d-1)$ -dimensional shellable complex;
- (iii)  $h$  is an  $M$ -sequence.

Since there are a total of  $\binom{n+k-1}{k}$  monomials of degree  $k$  in  $n$  variables, and by Theorems 15.1.4 and 15.2.2 the  $h$ -vector of a  $(d-1)$ -dimensional Cohen-Macaulay complex counts certain monomials in  $h_1 = f_0 - d$  variables, we derive the inequalities

$$0 \leq h_i \leq \binom{f_0 - d + i - 1}{i}$$

for the  $h$ -vectors of Cohen-Macaulay complexes. The lower bound can be improved for complexes with fixed-point-free involutive symmetry.

**THEOREM 15.2.3**

Let  $h = (h_0, \dots, h_d)$  be the  $h$ -vector of a Cohen-Macaulay complex admitting an automorphism  $\alpha$  of order 2, such that  $\alpha(F) \neq F$  for all  $F \in \Delta \setminus \{\emptyset\}$ . Then

$$h_i \geq \binom{d}{i} \quad \text{for } 0 \leq i \leq d.$$

Consequently,  $f_{d-1} = h_0 + \dots + h_d \geq 2^d$ .

Another condition on a Cohen-Macaulay complex that forces stricter conditions on its  $h$ -vector is being  $r$ -colorable.

**THEOREM 15.2.4**

For  $h = (h_0, \dots, h_d) \in \mathbb{Z}^{d+1}$  the following are equivalent:

- (i)  $h$  is the  $h$ -vector of a  $(d-1)$ -dimensional and  $d$ -colorable Cohen-Macaulay complex;
- (ii)  $(h_1, \dots, h_d)$  is the  $f$ -vector of a  $d$ -colorable simplicial complex.

Hence in this case the  $h$ -vector is not only an  $M$ -sequence, but the special kind of  $K$ -sequence characterized in Theorem 15.1.3.

**LERAY COMPLEXES**

Examples of Leray complexes arise as follows. Let  $\mathcal{K} = \{K_1, \dots, K_t\}$  be a family of convex sets in  $\mathbb{R}^m$ , and let  $\Delta(\mathcal{K}) = \{A \subseteq \{1, \dots, t\} \mid \bigcap_{i \in A} K_i \neq \emptyset\}$ . Then the simplicial complex  $\Delta(\mathcal{K})$  is  $m$ -Leray.

Fix  $m \geq 0$ , and let  $f = (f_0, \dots, f_{d-1})$  be the  $f$ -vector of a simplicial complex  $\Delta$ . Define

$$h_k^* = \begin{cases} f_k & \text{for } 0 \leq k \leq m-1 \\ \sum_{j \geq 0} (-1)^j \binom{k+j-m}{j} f_{k+j} & \text{for } k \geq m. \end{cases}$$

The sequence  $h^* = (h_0^*, \dots, h_{d-1}^*)$  is the  $h^*$ -vector of  $\Delta$ . The two vectors  $f$  and  $h^*$  mutually determine each other.

**THEOREM 15.2.5**

For  $h^* = (h_0^*, h_1^*, \dots) \in \mathbb{Z}^{(\infty)}$  the following are equivalent:

- (i)  $h^*$  is the  $h^*$ -vector of an  $m$ -Leray complex;
- (ii)  $h^*$  is the  $h^*$ -vector of  $\Delta(\mathcal{K})$  for some family  $\mathcal{K}$  of convex sets in  $\mathbb{R}^m$ ;
- (iii)

$$\begin{cases} h_k^* \geq 0 & \text{for } k \geq 0 \\ \partial_{k+1}(h_k^*) \leq h_{k-1}^* & \text{for } 1 \leq k \leq m-1 \\ \partial_m(h_k^*) \leq h_{k-1}^* - h_k^* & \text{for } k \geq m. \end{cases}$$

**COMMENTS**

The Euler-Poincaré formula (due to Poincaré 1899) is proved in most books on algebraic topology. Theorem 15.2.1 is from [BK88]. A good general source on Cohen-Macaulay complexes is [Sta96]; Theorems 15.2.2, 15.2.3, and 15.2.4, as well as references to the original sources, can be found there. A generalization of Theorem 15.2.2 to complexes whose  $k$ -skeleton is Cohen-Macaulay appears in [Bjö96]. There are several additional results about  $h$ -vectors of Cohen-Macaulay complexes. For instance, for complexes with nontrivial automorphism groups, see [Sta96, Section III.8]; for matroid complexes, see [Sta96, Section III.3]; and for Cohen-Macaulay complexes that are  $r$ -colorable for  $r < d$ , see the references mentioned in [Sta96, Section III.4].

Cohen-Macaulay complexes are closely related to certain commutative rings [Sta96], and via this connection such complexes have also been of use in the theory of splines; see [Sta96, Section III.5] and also Chapter 45.

Theorem 15.2.5 was conjectured by Eckhoff and proved by Kalai [Kal84, Kal86].

---

## 15.3 SIMPLICIAL POLYTOPES, SPHERES, AND MANIFOLDS

---

**GLOSSARY**

A **triangulated  $d$ -ball** is a simplicial complex  $\Delta$  whose realization  $\|\Delta\|$  is homeomorphic to the ball  $\{x \in \mathbb{R}^d \mid x_1^2 + \dots + x_d^2 \leq 1\}$ . A **triangulated  $(d-1)$ -sphere** is a simplicial complex whose realization is homeomorphic to the sphere  $\{x \in \mathbb{R}^d \mid x_1^2 + \dots + x_d^2 = 1\}$ . Equivalently, it is the boundary of a triangulated  $d$ -ball. Examples of triangulated  $(d-1)$ -spheres are given by the boundary complexes of simplicial  $d$ -polytopes.

A **pseudomanifold** is a pure simplicial complex  $\Delta$  such that

- (i) each face of codimension 1 is contained in precisely two maximal faces; and
- (ii) the dual graph (whose vertices are the maximal faces of  $\Delta$  and whose edges are the faces of codimension 1) is connected.

An **Eulerian pseudomanifold** is a pseudomanifold  $\Delta$  such that  $\Delta$  and the link of each face have the Euler characteristic of a sphere of the corresponding dimension.

A pure  $(d-1)$ -dimensional simplicial complex  $\Delta$  is a **homology manifold** if it is connected and the link of each nonempty face has the Betti numbers of a sphere of the same dimension. It is a **homology sphere** if, in addition,  $\Delta$  itself has the Betti numbers of a  $(d-1)$ -sphere. Examples of homology manifolds are given by triangulations of compact connected topological manifolds, i.e., spaces that are locally Euclidean.

The **cyclic  $d$ -polytope with  $n$  vertices**  $C_d(n)$  is the convex hull of any  $n$  points on the moment curve in  $\mathbb{R}^d$ . (See Section 13.1.4.)

The following implications hold among these various classes, all of them strict:

$$\begin{aligned} \text{polytope boundary} &\Rightarrow \text{sphere} \Rightarrow \text{homology sphere} \Rightarrow \\ &\quad \text{Eulerian pseudomanifold} \Rightarrow \text{pseudomanifold} \\ \text{homology sphere} &\Rightarrow \text{homology manifold} \Rightarrow \text{pseudomanifold} \\ \text{homology sphere} &\Rightarrow \text{Cohen-Macaulay complex} \end{aligned}$$

---

## PSEUDOMANIFOLDS

The following results give the basic lower and upper bounds on  $f$ -vectors of pseudomanifolds.

### THEOREM 15.3.1 Lower Bound Theorem

For a  $(d-1)$ -dimensional pseudomanifold  $\Delta$  with  $n$  vertices,

$$f_k(\Delta) \geq \begin{cases} \binom{d}{k}n - \binom{d+1}{k+1}k & \text{for } 1 \leq k \leq d-2 \\ (d-1)n - (d-2)(d+1) & \text{for } k = d-1. \end{cases}$$

### THEOREM 15.3.2 Upper Bound Theorem

Let  $\Delta$  be a  $(d-1)$ -dimensional homology manifold with  $n$  vertices, such that either

- (i)  $d$  is even, or
- (ii)  $d = 2k + 1$  is odd, and either  $\chi(\Delta) = 2$  or  $\beta_k \leq 2\beta_{k-1} + 2 \sum_{i=0}^{k-3} \beta_i$ .

Then  $f_k(\Delta) \leq f_k(C_d(n))$  for  $1 \leq k \leq d-1$ .

This upper bound theorem applies when the homology manifold is Eulerian (irrespective of dimension); in particular, it applies to all simplicial polytopes and spheres. By the geometric operation of “pulling vertices” (Section 14.2), one can extend this to all convex polytopes.

### THEOREM 15.3.3

If  $P$  is any convex  $d$ -polytope with  $n$  vertices, then  $f(P) \leq f(C_d(n))$ .

The given lower and upper bounds are best possible within the class of simplicial polytope boundaries. The lower bound is attained by the class of stacked polytopes



(Sections 14.4.2 and 17.2). To make the upper bound numerically explicit, we give the formula for the  $f$ -vector of a cyclic polytope.

### THEOREM 15.3.4

For  $d \geq 2$  and  $0 \leq k \leq d-1$ , the number of  $k$ -faces of the cyclic polytope  $C_d(n)$  with  $n$  vertices is

$$f_k(C_d(n)) = \frac{n - \delta(n - k - 2)}{n - k - 1} \sum_{j=0}^{\lfloor d/2 \rfloor} \binom{n-1-j}{k+1-j} \binom{n-k-1}{2j-k-1+\delta},$$

where  $\delta = d - 2\lfloor d/2 \rfloor$ . In particular,

$$f_{d-1}(C_d(n)) = \binom{n - \lfloor \frac{d+1}{2} \rfloor}{n-d} + \binom{n - \lfloor \frac{d+2}{2} \rfloor}{n-d}.$$

---

## POLYTOPES AND SPHERES

For boundaries of simplicial  $d$ -polytopes and, more generally, for Eulerian pseudomanifolds, we have the following basic relations.

### THEOREM 15.3.5 Dehn-Sommerville Equations

For  $d$ -dimensional Eulerian pseudomanifolds,

$$h_i = h_{d-i} \quad \text{for all } 0 \leq i \leq d.$$

These equations give a complete description of the linear span of all  $f$ -vectors of  $d$ -polytopes (equivalently,  $(d-1)$ -spheres). (The affine span is defined by including the relation  $h_0 = 1$ .)

One consequence of the Dehn-Sommerville equations is the following relation between the  $h$ -vector of a triangulated ball  $K$  and the  $g$ -vector of its boundary  $\partial K$ .

### THEOREM 15.3.6

For a triangulated  $d$ -ball  $K$  and its boundary  $(d-1)$ -sphere  $\partial K$ ,

$$g_i(\partial K) = h_i(K) - h_{d+1-i}(K) \quad \text{for } i \geq 1.$$

A complete characterization of the  $f$ -vectors of simplicial (and, by duality, simple) convex polytopes is given in terms of the  $h$ -vector and  $g$ -vector.

### THEOREM 15.3.7 $g$ -Theorem

A nonnegative integer vector  $h = (h_0, \dots, h_d)$  is the  $h$ -vector of a simplicial convex polytope if and only if

- (i)  $h_i = h_{d-i}$ , and
- (ii)  $(g_0, \dots, g_{\lfloor d/2 \rfloor})$  is an  $M$ -sequence.

One consequence of (ii) is that  $g_i \geq 0$ . For centrally symmetric polytopes, we get a better lower bound.

**THEOREM 15.3.8**

For centrally symmetric simplicial  $d$ -polytopes,

$$g_i = h_i - h_{i-1} \geq \binom{d}{i} - \binom{d}{i-1} \quad \text{for } i \leq \lfloor d/2 \rfloor.$$

**COMMENTS**

The Lower Bound Theorem 15.3.1 is due to Kalai and Gromov in the generality given here; see [Kal87] including the note added in proof. The  $k = d - 1$  case had earlier been done by Klee and the case of polytope boundaries by Barnette. See [Kal87] for a discussion of the history of this result.

The Upper Bound Theorem 15.3.2 is due to Novik [Nov]. The case of polytopes (Theorem 15.3.3) was first proved by McMullen (see [MS71]), and extended to spheres by Stanley (see [Sta96]). The computation of the  $f$ -vector of the cyclic polytope can be found in [Grü67, Sections 4.7.3 and 9.6.1] or [MS71].

The Dehn-Sommerville equations for polytopes are classical; proofs can be found in [Grü67, Sta86, Zie95]. The extension to Eulerian pseudomanifolds is due to Klee [Kle64]; an equivariant version appears in [Bar92]. The  $g$ -theorem was conjectured by McMullen and proved by Billera, Lee, and Stanley [BL81, Sta80]. More recently, another proof of the necessity of these conditions was given by McMullen [McM93]. It is not known whether the second condition of Theorem 15.3.7 holds for general triangulated spheres. The  $g$ -theorem has a convenient reformulation as a one-to-one correspondence (via matrix multiplication) between  $f$ -vectors of simplicial polytopes and  $M$ -sequences, see [Bjö87, Zie95]. Theorem 15.3.8 was proved by Stanley [Sta87a].

The study of  $f$ -vectors of unbounded polyhedra can be approached by studying the  $f$ -vectors of **polytope pairs**  $(P, F)$ , where  $P$  is a polytope and  $F$  is a maximal face of  $P$ . See [BL93] for a summary of such results.

**15.4 CELL COMPLEXES****GLOSSARY**

*Convex polytopes* and *faces* of such are defined in Chapter 13.

A **polyhedral complex**  $\Gamma$  is a finite collection of convex polytopes in  $\mathbb{R}^n$  such that (i) if  $\pi \in \Gamma$  and  $\sigma$  is a face of  $\pi$ , then  $\sigma \in \Gamma$ ; and (ii) if  $\pi, \sigma \in \Gamma$  and  $\pi \cap \sigma \neq \emptyset$ , then  $\pi \cap \sigma$  is a face of both. The **space** of  $\Gamma$  is  $\|\Gamma\| = \bigcup_{\pi \in \Gamma} \pi$ , a subspace of  $\mathbb{R}^n$ . Examples of polyhedral complexes are given by **boundary complexes**  $\partial P$  of convex polytopes  $P$  (i.e., the collection of all proper faces). A geometric simplicial complex (defined in Section 15.1) is a polyhedral complex all of whose cells are simplices. A **cubical complex** is a polyhedral complex all of whose cells are (combinatorially isomorphic to) cubes.

A **regular cell complex**  $\Gamma$  is a family of closed balls (homeomorphs of  $\{x \in \mathbb{R}^j \mid |x| \leq 1\}$ ) in a Hausdorff space  $\|\Gamma\|$  such that (i) the interiors of the balls

partition  $\|\Gamma\|$  and (ii) the boundary of each ball in  $\Gamma$  is a union of other balls in  $\Gamma$ . The members of  $\Gamma$  are called (closed) **cells** or **faces**. The **dimension** of a cell is its topological dimension and  $\dim \Gamma = \max_{\sigma \in \Gamma} \dim \sigma$ .

A regular cell complex has the **intersection property** if, whenever the intersection of two cells is nonempty, then this intersection is also a cell in the complex. Polyhedral complexes are examples of regular cell complexes with the intersection property. Regular cell complexes with the intersection property can be reconstructed up to homeomorphism from the corresponding “abstract” complex consisting of the family of vertex sets of its cells.

For a regular cell complex  $\Gamma$ , let  $f_i$  be the number of  $i$ -dimensional cells, and let  $\beta_i = \dim_{\mathbb{Q}} \tilde{H}_i(\|\Gamma\|, \mathbb{Q})$ . The latter denotes  $i$ -dimensional reduced singular homology with rational coefficients of the space  $\|\Gamma\|$ ; see [Mun84, Spa66] for explanations of this concept. Then we have the  **$f$ -vector**  $f = (f_0, f_1, \dots)$  and the **Betti sequence**  $\beta = (\beta_0, \beta_1, \dots)$  of  $\Gamma$ . These definitions generalize those previously given in the simplicial case.

---

## BASIC $f$ -VECTOR RELATIONS

Among the classes of complexes

- simplicial complexes
- polyhedral complexes
- regular cell complexes with the intersection property
- regular cell complexes

each is a subclass of its successor. Thus one may wonder how many of the relations for  $f$ -vectors of simplicial complexes given in Sections 15.1–15.3 can be extended to these broader classes of complexes. Also, what new phenomena (not visible in the simplicial case) arise? Some answers will be given in this section and the following one, but current knowledge is quite fragmentary. We begin here with the most general relations.

### THEOREM 15.4.1

$(f_0, \dots, f_d)$  is the  $f$ -vector of a  $d$ -dimensional regular cell complex if and only if  $f_d \geq 1$  and  $f_i \geq 2$  for all  $0 \leq i < d$ .

### THEOREM 15.4.2

$f$  is the  $f$ -vector of a regular cell complex with the intersection property if and only if  $f$  is a  $K$ -sequence.

Let  $\beta = (\beta_0, \beta_1, \dots) \in \mathbb{N}^{(\infty)}$  be fixed, and for every sequence  $f = (f_0, f_1, \dots)$  let

$$\chi_{k-1} = \sum_{j \geq k} (-1)^{j-k} (f_j - \beta_j) \quad \text{for } k \geq 0.$$

### THEOREM 15.4.3

$(f_0, \dots, f_d)$  is the  $f$ -vector of a  $d$ -dimensional regular cell complex with Betti sequence  $\beta$  if and only if  $\chi_{-1} = 1$  and  $\chi_k \geq 1$  for  $0 \leq k < d$ .

**THEOREM 15.4.4**

For  $f \in \mathbb{N}^{(\infty)}$  the following are equivalent:

- (i)  $f$  is the  $f$ -vector of a regular cell complex with the intersection property and with Betti sequence  $\beta$ ;
- (ii)  $\chi_{-1} = 1$  and  $\partial_{k+1}(\chi_k + \beta_k) \leq \chi_{k-1}$  for all  $k \geq 1$ .

These results show that the  $f$ -vectors of regular cell complexes (with or without Betti number constraints) are considerably more general than the  $f$ -vectors of simplicial complexes, but that the two classes of  $f$ -vectors agree in the presence of the intersection property.

**COMMENTS**

Regular cell complexes are known as **regular CW complexes** in the topological literature [LW69]. The nonregular CW complexes offer an even more general class of cell complexes [LW69, Mun84, Spa66], but there is very little one can say about  $f$ -vectors in that generality. See [BLS<sup>+</sup>93, Section 4.7] for a detailed discussion of regular cell complexes from a combinatorial point of view.

For the results of this section see [BK88, BK91, BK89]. A characterization of  $f$ -vectors of (cubical) subcomplexes of a cube can be found in [Lin71], and of regular cell decompositions of spheres in [Bay88].

**15.5 GENERAL POLYTOPES AND SPHERES****GLOSSARY**

A **flag** of faces in a (polyhedral)  $(d-1)$ -complex  $\Delta$  is a chain  $F_1 \subsetneq F_2 \subsetneq \cdots \subsetneq F_k$  in  $\Delta$ . It is an  **$S$ -flag** if

$$S = \{\dim F_1, \dots, \dim F_k\} \subseteq \{0, 1, \dots, d-1\}.$$

If  $f_S = f_S(\Delta)$  denotes the number of  $S$ -flags in  $\Delta$ , then the function  $S \mapsto f_S$ ,  $S \subseteq \{0, 1, \dots, d-1\}$ , is called the **flag  $f$ -vector** of  $\Delta$ .

If 
$$h_S = \sum_{T \subseteq S} (-1)^{|S|-|T|} f_T,$$

then the function  $S \mapsto h_S$ ,  $S \subseteq \{0, 1, \dots, d-1\}$ , is called the **flag  $h$ -vector**.

For  $S \subseteq \{0, \dots, d-1\}$  and noncommuting symbols  $\mathbf{a}$  and  $\mathbf{b}$ , let  $u_S = u_0 u_1 \cdots u_{d-1}$  be the  **$\mathbf{ab}$ -word** defined by  $u_i = \mathbf{a}$  if  $i \notin S$  and  $u_i = \mathbf{b}$  otherwise. When  $\Delta$  is spherical (or, more generally, Eulerian), then the  **$\mathbf{ab}$ -polynomial**  $\sum h_S u_S$  is also a polynomial in  $\mathbf{c} = \mathbf{a} + \mathbf{b}$  and  $\mathbf{d} = \mathbf{ab} + \mathbf{ba}$ . (Note that the degree of  $\mathbf{c}$  is 1 and the degree of  $\mathbf{d}$  is 2.) The resulting  **$\mathbf{cd}$ -polynomial**

$$\sum h_S u_S = \sum \phi_w w,$$

where the right-hand sum is over all  **$\mathbf{cd}$ -words**  $w$  of degree  $d$ , is called the  **$\mathbf{cd}$ -index** of  $\Delta$ . For 2- and 3-polytopes, the  **$\mathbf{cd}$ -index** is  $\mathbf{c}^2 + (f_0 - 2)\mathbf{d}$  and  $\mathbf{c}^3 + (f_0 - 2)\mathbf{dc} + (f_2 - 2)\mathbf{cd}$ , respectively.

For any convex  $d$ -polytope  $P$ , we define the **toric  $h$ -vector** and **toric  $g$ -vector** recursively by  $h(P, x) = \sum_{i=0}^d h_i x^{d-i}$  and  $g(P, x) = \sum_{i=0}^{\lfloor d/2 \rfloor} g_i x^i$ , where  $g_i = h_i - h_{i-1}$  and the following relations hold:

- (i)  $g(\emptyset, x) = h(\emptyset, x) = 1$ ; and
- (ii)  $h(P, x) = \sum_{G \text{ face of } P, G \neq P} g(G, x)(x-1)^{d-1-\dim G}$ .

(Compare to Chapter 14.4.2, where this toric  $h$ -vector is defined for any polyhedral complex. In the notation given there, we have defined  $h$  and  $g$  for the complex  $\partial P$ .) When  $P$  is simplicial, this definition coincides with that of the usual  $h$ -vector, as defined in Section 15.2.

A **rational polytope** is one whose vertices all have rational coordinates.

A **cubical polytope** is one that has a cubical boundary complex. For any cubical  $(d-1)$ -complex with  $f$ -vector  $(f_0, \dots, f_{d-1})$ , define the **cubical  $h$ -vector**  $h^c = (h_0^c, \dots, h_d^c)$  by

$$h_i^c = (-1)^i 2^{d-1} + \sum_{j=1}^i (-1)^{i-j} 2^{j-1} f_{j-1} \sum_{k=0}^{i-j} \binom{d-j}{k} \quad \text{for } i = 0, \dots, d.$$

The **cubical  $g$ -vector**  $g^c = (g_0^c, \dots, g_{\lfloor d/2 \rfloor}^c)$  is defined by  $g_0^c = h_0^c = 2^{d-1}$  and  $g_i^c = h_i^c - h_{i-1}^c$  for  $i \geq 1$ .

An **Eulerian polyhedral complex** is one whose first barycentric subdivision is an Eulerian pseudomanifold. Examples are boundary complexes of polytopes and **spherical** polyhedral complexes, i.e., those whose underlying space is homeomorphic to a sphere.

A (central) **hyperplane arrangement** is a collection  $\mathcal{H}$  of  $n$  linear hyperplanes in  $\mathbb{R}^d$ , given by normal vectors  $x_1, \dots, x_n$  (see Section 6.1.3). The arrangement is **essential** if the normals  $x_i$  span  $\mathbb{R}^d$ . The associated **zonotope** is the Minkowski sum of the  $n$  line segments  $[-x_i, x_i]$ , i.e.,  $Z = \{\sum \lambda_i x_i \mid -1 \leq \lambda_i \leq 1\}$  (see Section 13.1.4).

## LINEAR RELATIONS

We give the linear relations on the invariants defined above that are known to hold for all boundary complexes of polytopes and, more generally, for all Eulerian polyhedral complexes.

### THEOREM 15.5.1

For  $(d-1)$ -dimensional Eulerian polyhedral complexes, the following relations always hold for the flag  $h$ , the toric  $h$ , and the flag  $f$ :

- (i)  $h_S = h_{\{0, \dots, d-1\} \setminus S}$  for all  $S \subseteq \{0, \dots, d-1\}$ ;
- (ii)  $h_i = h_{d-i}$  for  $0 \leq i \leq d$ ; and
- (iii)  $\sum_{j=i+1}^{k-1} (-1)^{j-i-1} f_{S \cup \{j\}} = (1 - (-1)^{k-i-1}) f_S$  whenever  $i, k \in S \cup \{-1, d\}$  with  $i \leq k-2$  and  $S \cap \{i+1, \dots, k-1\} = \emptyset$ .

It is known that the relations in Theorem 15.5.1(iii), the **generalized Dehn-Sommerville equations**, completely describe the linear span of all flag  $f$ -vectors of Eulerian complexes, and so they imply those in (i). Since the toric  $h$  is known to be a linear function of the flag  $f$ , they imply those in (ii) as well. The linear span of flag  $f$ -vectors has dimension  $e_d$ , where  $e_d$  is the  $d^{\text{th}}$  Fibonacci number (defined by

the recurrence  $e_d = e_{d-1} + e_{d-2}$ ,  $e_0 = e_1 = 1$ ). There are  $e_d$   $cd$ -words of degree  $d$ . Furthermore, the coefficients  $\phi_w$  of the  $cd$ -index, considered as linear expressions in the  $f_S$ , form a linear basis for the span of flag  $f$ -vectors of  $d$ -polytopes. The affine span of all flag  $f$ -vectors is defined by including the relation  $f_\emptyset = 1$ .

For cubical polytopes and spheres, the cubical  $h$ -vector satisfies the analogue of the Dehn-Sommerville equations.

### THEOREM 15.5.2

For cubical  $d$ -polytopes and cubical  $(d-1)$ -spheres,

$$h_i^c = h_{d-i}^c \quad \text{for all } 0 \leq i \leq d.$$

These give all linear relations satisfied by  $f$ -vectors of cubical polytopes and spheres. The cubical  $h$ -vector satisfies, as well, the equations of Theorem 15.3.6, linking the  $h$  of a cubical ball to the  $g$  of its boundary sphere.

---

## LINEAR INEQUALITIES

Some linear inequalities that hold for flag  $f$ -vectors of all polytope boundaries are given in this section. The list is not thought to be complete, although there are no conjectures for what the complete set might be.

For a Cohen-Macaulay polyhedral complex, i.e., one whose first barycentric subdivision is a Cohen-Macaulay simplicial complex, the flag  $h$  is always nonnegative.

### THEOREM 15.5.3

For a Cohen-Macaulay polyhedral  $(d-1)$ -complex  $\Gamma$ ,

$$h_S(\Gamma) \geq 0 \quad \text{for all } S \subseteq \{0, \dots, d-1\}.$$

For general convex polytopes, we also have nonnegativity of the  $cd$ -index.

### THEOREM 15.5.4

For a convex  $d$ -polytope  $P$ ,  $\phi_w \geq 0$  for all  $cd$ -words  $w$  of degree  $d$ .

For rational convex polytopes, it is known, further, that the toric  $h$  is unimodal.

### THEOREM 15.5.5

For a rational convex  $d$ -polytope,  $g_i \geq 0$  for  $i \leq \lfloor d/2 \rfloor$ .

Related to this is the following *nonlinear* inequality holding between the  $g$ -vectors of a polytope  $P$  and any of its faces  $F$ . We denote by  $P/F$  the *link* of  $F$  in  $P$ , i.e., the polytope whose lattice of faces is (isomorphic to) the interval  $[F, P]$  in the face lattice of  $P$ .

### THEOREM 15.5.6

For a rational polytope  $P$  and any face  $F$ , we have the polynomial inequality

$$g(P, t) - g(F, t)g(P/F, t) \geq 0,$$

i.e., all coefficients of this polynomial are nonnegative.

Finally, we have the following lower bound for the number of vertices of polytopes with no triangular faces (this includes the class of cubical polytopes).

### THEOREM 15.5.7

*A  $d$ -polytope with no triangular 2-face has at least  $2^d$  vertices.*

---

## HYPERPLANE ARRANGEMENTS AND ZONOTOPES

An essential hyperplane arrangement  $\mathcal{H}$  defines a decomposition of  $\mathbb{R}^d$  into polyhedral cones (as in Section 6.1.3). This decomposition  $\Gamma_{\mathcal{H}}$ , a regular cell complex if intersected with the unit sphere, has a flag  $f$ -vector dual to that of its associated zonotope  $Z$ , in the sense that  $f_S(\Gamma_{\mathcal{H}}) = f_{d-S}(Z)$ , where  $S = \{i_1, \dots, i_k\} \subseteq \{1, \dots, d\}$  and  $d - S = \{d - i_k, \dots, d - i_1\}$ .

### THEOREM 15.5.8

*The flag  $f$ -vector of an arrangement (or zonotope) depends only on the matroid (linear dependency structure) of the underlying point configuration  $\{x_1, \dots, x_n\}$ .*

Although a fairly special subclass of polytopes, the zonotopes nonetheless are varied enough to carry all the linear information carried by flag numbers of general polytopes.

### THEOREM 15.5.9

*The flag  $f$ -vectors of zonotopes (and thus of hyperplane arrangements) satisfy the generalized Dehn-Sommerville equations, and no other linear relations not implied by these.*

When it comes to linear *inequalities*, however, the difference between zonotopes and general polytopes emerges. The following result has the most direct interpretation when it is stated for arrangements, where it bounds the average number of  $S = \{i_1, \dots, i_k\}$ -flags in an  $i_k$ -face by the number of  $S$ -flags in an  $i_k$ -cube.

### THEOREM 15.5.10

*For a hyperplane arrangement  $\mathcal{H}$  in  $\mathbb{R}^d$  and  $S = \{i_1, \dots, i_k\} \subseteq \{1, \dots, d\}$  with  $k \geq 2$ ,*

$$\frac{f_S(\Gamma_{\mathcal{H}})}{f_{i_k}(\Gamma_{\mathcal{H}})} < \binom{i_k}{i_1, i_2 - i_1, \dots, i_k - i_{k-1}} 2^{i_k - i_1}.$$

There is a straightforward reformulation of Theorem 15.5.10 for zonotopes that is easily seen not to be valid for all polytopes.

---

## GENERAL 3- AND 4-POLYTOPES

We describe here the situation for flag  $f$ -vectors of 3- and 4-polytopes. The equations in Theorem 15.5.1(iii) reduce consideration to  $(f_0, f_2)$  when  $d = 3$  and to  $(f_0, f_1, f_2, f_{02})$  when  $d = 4$ .

**THEOREM 15.5.11**

For 3-polytopes, the following is known about the vector  $(f_0, f_2)$ .

- (i) An integer vector  $(f_0, f_2)$  is the  $f$ -vector of a 3-polytope if and only if  $f_0 \leq 2f_2 - 4$  and  $f_2 \leq 2f_0 - 4$ .
- (ii) An integer vector  $(f_0, f_2)$  is the  $f$ -vector of a cubical 3-polytope if and only if  $f_2 = f_0 - 2$ ,  $f_0 \geq 8$ , and  $f_0 \neq 9$ .
- (iii) If  $(f_0, f_2) = (f_0(Z), f_2(Z))$  for a 3-zonotope  $Z$ , then  $f_0$  and  $f_1$  are both even integers,  $f_0 \leq 2f_2 - 4$ , and  $f_2 \leq f_0 - 2$ .

For 4-polytopes, much less is known.

**THEOREM 15.5.12**

Flag  $f$ -vectors  $(f_0, f_1, f_2, f_{02})$  of 4-polytopes satisfy the following inequalities.

- (i)  $f_{02} \geq 3f_2$
- (ii)  $f_{02} \geq 3f_1$
- (iii)  $f_{02} + f_1 + 10 \geq 3f_2 + 4f_0$
- (iv)  $6f_1 \geq 6f_0 + f_{02}$
- (v)  $f_0 \geq 5$
- (vi)  $f_0 + f_2 \geq f_1 + 5$
- (vii)  $2(f_{02} - 3f_2) \leq \binom{f_0}{2}$
- (viii)  $2(f_{02} - 3f_1) \leq \binom{f_2 - f_1 + f_0}{2}$
- (ix)  $f_{02} - 4f_2 + 3f_1 - 2f_0 \leq \binom{f_0}{2}$
- (x)  $f_{02} + f_2 - 2f_1 - 2f_0 \leq \binom{f_2 - f_1 + f_0}{2}$ .

It is not known, for example, whether (i)–(vi) give all linear inequalities holding for flag  $f$ -vectors of 4-polytopes.

**COMMENTS**

It is thought that the best route to an eventual characterization of  $f$ -vectors of general polytopes lies in an understanding of their flag  $f$ -vectors. The latter inherit many of the algebraic properties of  $f$ -vectors of simplicial polytopes that led to their characterization.

The relations in Theorem 15.5.1 hold more generally for the case of enumeration of chains in Eulerian posets; see the article by Stanley in [BMSW94]. The relations in Theorem 15.5.1(iii) are proved in [BB85]. An expression for the toric  $h$  in terms of the flag  $f$ , as well as a discussion of the convolution product (originally due to Kalai), can be found in the article by Bayer in [BMSW94]. The article by Kalai in the same volume contains an extensive discussion of  $g$ -vectors for both simplicial and general polytopes. The form of the cubical Dehn-Sommerville equations given in Theorem 15.5.2 appeared in [Adi96].

Theorem 15.5.4 holds as well for certain shellable spheres (see [Sta96, Section III.4]). Theorem 15.5.3 can also be found in [Sta96, Theorem III.4.4] (where  $h_S$  is denoted  $\beta(S)$ ). Theorem 15.5.5 appears in [Sta87b]. Theorem 15.5.7 is due to Blind and Blind [BB90]. There is a notion of convolution product of flag  $f$



numbers that can be used to produce new linear inequalities from given ones; see [BL93, Section 3.10].

Note that Theorem 15.5.6, due to Braden and MacPherson [BM97], gives a connection between the  $g$ -vector of a polytope  $P$  and that of one of its faces. This is an example of a “monotonicity theorem” related to face numbers. For similar theorems relating  $h$ -vectors of subcomplexes and subdivisions of a simplicial complex  $\Delta$ , see Sections III.9–10 of [Sta96] and the references given there.

For the fact that the flag  $f$ -vector of a zonotope or arrangement (or, more generally, of an oriented matroid) depends only on the underlying matroid, see [BLS<sup>+</sup>93, Cor. 4.6.3]. That the only linear relations satisfied by zonotopes are the generalized Dehn-Sommerville equations of Theorem 15.5.1(iii) is proved in [BER]. Theorem 15.5.10 is due to Varchenko for the case  $k = 2$  (see [BLS<sup>+</sup>93, Proposition 4.6.9]) and to Liu [Liu95, Theorem 4.8.2] in the form given here.

Theorem 15.5.11(i) can be found in [Grü67, Section 10.3]; 15.5.11(ii) appears in dual form (for 4-valent 3-polytopes) in [Bar83]; 15.5.11(iii) can be derived using the methods of [Grü67, Section 18.2] (see also [BER]). Theorem 15.5.12 can be found in [Bay87].

## 15.6 OPEN PROBLEMS

### PROBLEM 15.6.1

*Characterize the  $f$ -vectors of triangulations of the  $(d-1)$ -sphere. It has been conjectured that the conditions of the  $g$ -theorem provide the answer.*

### PROBLEM 15.6.2

*Characterize the  $f$ -vectors of triangulations of the  $d$ -ball.*

### PROBLEM 15.6.3

*Characterize the  $f$ -vectors of triangulations of the  $d$ -torus. It is known that  $f(2\text{-torus}) = \{(n, 3n, 2n) \mid n \geq 7\}$ , but the question is open for  $d \geq 3$ .*

### PROBLEM 15.6.4

*Characterize the  $f$ -vectors of  $d$ -polytopes. The answer is known for  $d \leq 3$  (Theorem 15.5.11(i)), but for  $d \geq 4$  there is not even a conjectured answer.*

### PROBLEM 15.6.5 I. Bárány

*Does there exist a constant  $c_d > 0$  such that  $f_i \geq c_d \cdot \min\{f_0, f_{d-1}\}$  for all  $d$ -polytopes and all  $i$ ? Will  $c_d = 1$  do?*

### PROBLEM 15.6.6

*Characterize the  $f$ -vectors of centrally symmetric  $d$ -polytopes. The question is open in the simplicial as well as in the general case. Even an upper bound conjecture in the simplicial and centrally symmetric case is missing.*

### PROBLEM 15.6.7 Conjecture of G. Kalai

*The total number of faces (counting  $P$  but not  $\emptyset$ ) of a centrally symmetric convex*

$d$ -polytope  $P$  is  $\geq 3^d$ . (Verified in the simplicial case as a consequence of Theorem 15.3.8.)

**PROBLEM 15.6.8**

The **clique complex** of a graph is the collection of vertex sets of all its cliques (complete induced subgraphs). Characterize the  $f$ -vectors of clique complexes.

**PROBLEM 15.6.9** G. Kalai

Is the  $f$ -vector of an  $(r-1)$ -dimensional clique complex the  $f$ -vector of an  $r$ -colorable complex?

**PROBLEM 15.6.10** Conjecture of Charney and Davis [Sta96, p. 100]

Let  $(g_0, \dots, g_k)$  be the  $g$ -vector of a clique complex homeomorphic to the sphere  $S^{2k-1}$ . Then  $g_k - g_{k-1} + \dots + (-1)^k g_0 \geq 0$ .

**PROBLEM 15.6.11** Conjecture of Stanley [Sta96, p. 102]

Every coefficient  $\phi_w$  of the **cd-index** of a sphere is nonnegative.

**PROBLEM 15.6.12** Adin [Adi96] (The case  $i = 1$  is implied by Theorem 15.5.7.)

The generalized lower bound conjecture for cubical  $d$ -polytopes and  $d$ -spheres:  $g_i^c \geq 0$  for  $i \leq \lfloor d/2 \rfloor$ . This has been shown to be the best possible set of linear inequalities for cubical  $d$ -spheres [BBC]. More generally, characterize  $f$ -vectors of cubical polytopes.

**PROBLEM 15.6.13**

Characterize the flag  $f$ -vectors of polytopes and of zonotopes. In particular, determine a complete set of linear inequalities holding for flag  $f$ -vectors of polytopes and of zonotopes.

**PROBLEM 15.6.14**

Characterize (toric)  $h$ -vectors of general polytopes.

---

## 15.7 SOURCES AND RELATED MATERIAL

---

### FURTHER READING

Surveys of  $f$ -vector theory are given in [BL93, Bjö87, BK89, KK95, Sta85]. Books treating  $f$ -vectors (among other things) include [And87, BMSW94, Grü67, MS71, Sta96, Zie95].

---

### RELATED CHAPTERS

Chapter 6: Oriented matroids  
Chapter 13: Basic properties of convex polytopes  
Chapter 14: Subdivisions and triangulations of polytopes

## Chapter 45: Splines and geometric modeling

## REFERENCES

- [Adi96] R.M. Adin. A new cubical  $h$ -vector. *Discrete Math.*, 157:3–14, 1996.
- [And87] I. Anderson. *Combinatorics of Finite Sets*. Clarendon Press, Oxford, 1987.
- [BBC] E.K. Babson, L.J. Billera, and C. Chan. Neighborly cubical spheres and a cubical lower bound conjecture. *Israel J. Math.*, to appear.
- [Bar83] D.W. Barnette. *Map Coloring, Polyhedra, and the Four Color Theorem*. Number 8 of *Dolciani Math. Exp.*, Math. Assoc. America, Washington, 1983.
- [Bar92] A.I. Barvinok. On equivariant generalization of Dehn-Sommerville equation. *Europ. J. Combin.*, 13:419–428, 1992.
- [Bay87] M.M. Bayer. The extended  $f$ -vectors of 4-polytopes. *J. Combin. Theory. Ser. A*, 44:141–151, 1987.
- [Bay88] M.M. Bayer. *Barycentric subdivisions*. *Pacific J. Math.*, 135:1–16, 1988.
- [BB85] M.M. Bayer and L.J. Billera. Generalized Dehn-Sommerville relations for polytopes, spheres and Eulerian partially ordered sets. *Invent. Math.*, 79:143–157, 1985.
- [BL93] M.M. Bayer and C.W. Lee. Combinatorial aspects of convex polytopes. In P.M. Gruber and J.M. Wills, editors, *Handbook of Convex Geometry*, pages 485–534. North-Holland, Amsterdam, 1993.
- [BER] L.J. Billera, R. Ehrenborg, and M. Readdy. The  $cd$ -index of zonotopes and arrangements. In *Mathematical Essays in Honor of Gian-Carlo Rota*, Birkhäuser, Boston, to appear.
- [BL81] L.J. Billera and C.W. Lee. A proof of the sufficiency of McMullen's conditions for  $f$ -vectors of simplicial polytopes. *J. Combin. Theory Ser. A*, 31:237–255, 1981.
- [BMSW94] T. Bisztriczky, P. McMullen, R. Schneider, and A. Ivić Weiss, editors. *Polytopes: Abstract, Convex and Computational*. Volume 440 of *NATO Adv. Sci. Inst. Ser. C: Math. Phys. Sci.* Kluwer, Dordrecht, 1994.
- [Bjö87] A. Björner. Face numbers of complexes and polytopes. In *Proc. Internat. Cong. Math., Berkeley, 1986*, pages 1408–1418. Amer. Math. Soc., Providence, 1987.
- [Bjö96] A. Björner. Nonpure shellability,  $f$ -vectors, subspace arrangements and complexity. In L.J. Billera, C. Greene, R. Simion, and R. Stanley, editors, *Formal Power Series and Algebraic Combinatorics, DIMACS Ser. in Discrete Math. and Theor. Comput. Sci.*, pages 25–53. Amer. Math. Soc., Providence, 1996.
- [BK88] A. Björner and G. Kalai. An extended Euler-Poincaré theorem. *Acta Math.*, 161:279–303, 1988.
- [BK89] A. Björner and G. Kalai. On  $f$ -vectors and homology. In G. Bloom, R. Graham, and J. Malkevitch, editors, *Combinatorial Mathematics: Proc. 3rd Internat. Conf., New York, 1985*, volume 555 of *Ann. New York Acad. Sci.*, pages 63–80. New York Acad. Sci., 1989.
- [BK91] A. Björner and G. Kalai. Extended Euler-Poincaré relations for cell complexes. In P. Gritzmann and B. Sturmfels, editors, *Applied Geometry and Discrete Mathematics—The Victor Klee Festschrift*, pages 81–89, volume 4 of *DIMACS Series in Discrete Math. and Theor. Comput. Sci.* Amer. Math. Soc., Providence, 1991.
- [BLS<sup>+</sup>93] A. Björner, M. LasVergnas, B. Sturmfels, N. White, and G.M. Ziegler. *Oriented Matroids*. Volume 46 of *Encyclopedia Math. Appl.*, Cambridge University Press, 1993.
- [BB90] G. Blind and R. Blind. Convex polytopes without triangular faces. *Israel Jour. Math.*, 71:129–134, 1990.

- [BM97] T.C. Braden and R. MacPherson. Intersection homology of toric varieties and a conjecture of Kalai. Preprint, 1997.
- [FFK88] P. Frankl, Z. Füredi, and G. Kalai. Shadows of colored complexes. *Math. Scand.*, 63:169–178, 1988.
- [FK96] E. Friedgut and G. Kalai. Every monotone graph property has a sharp threshold. *Proc. Amer. Math. Soc.*, 124:2993–3002, 1996.
- [Grü67] B. Grünbaum. *Convex Polytopes*. Interscience, London, 1967; revised edition (V. Klee and P. Kleinschmidt, editors), *Grad. Texts in Math.*, Springer-Verlag, in preparation.
- [Hib89] T. Hibi. What can be said about pure O-sequences? *J. Combin. Theory Ser. A*, 50:319–322, 1989.
- [Kal84] G. Kalai. A characterization of  $f$ -vectors of families of convex sets in  $\mathbb{R}^d$ . Part I: Necessity of Eckhoff's conditions. *Israel J. Math.*, 48:175–195, 1984.
- [Kal86] G. Kalai. A characterization of  $f$ -vectors of families of convex sets in  $\mathbb{R}^d$ . Part II: Sufficiency of Eckhoff's conditions. *J. Combin. Theory Ser. A*, 41:167–188, 1986.
- [Kal87] G. Kalai. Rigidity and the lower bound theorem 1. *Invent. Math.*, 88:125–151, 1987.
- [Kle64] V. Klee. A combinatorial analogue of Poincaré's duality theorem. *Canad. J. Math.*, 16:517–531, 1964.
- [KK95] V. Klee and P. Kleinschmidt. Convex polytopes and related complexes. In R. Graham, M. Grötschel, and L. Lovász, editors, *Handbook of Combinatorics*, pages 875–917. North-Holland, Amsterdam, 1995.
- [Lin71] B. Lindström. The optimal number of faces in cubical complexes. *Ark. Mat.*, 8:245–257, 1971.
- [Liu95] N. Liu. *Algebraic and Combinatorial Methods for Face Enumeration in Polytopes*. Ph.D. Thesis, Cornell Univ., Ithaca, 1995.
- [LW69] A.T. Lundell and S. Weingram. *The Topology of CW Complexes*. Van Nostrand, New York, 1969.
- [McM93] P. McMullen. On simple polytopes. *Invent. Math.*, 113:419–444, 1993.
- [MS71] P. McMullen and G.C. Shephard. *Convex Polytopes and the Upper Bound Conjecture*. Volume 3 of *London Math. Soc. Lecture Note Ser.*, Cambridge University Press, 1971.
- [Mun84] J.R. Munkres. *Elements of Algebraic Topology*. Addison-Wesley, Reading, 1984.
- [Nov] I. Novik. Upper bound theorems for simplicial manifolds. *Israel J. Math.*, to appear.
- [Spa66] E.H. Spanier. *Algebraic Topology*. McGraw-Hill, New York, 1966.
- [Sta80] R.P. Stanley. The number of faces of simplicial convex polytopes. *Adv. Math.*, 35:236–238, 1980.
- [Sta85] R.P. Stanley. The number of faces of simplicial polytopes and spheres. In J.E. Goodman, E. Lutwak, J. Malkevitch, and R. Pollack, editors, *Discrete Geometry and Convexity*, volume 440 of *Ann. New York Acad. Sci.*, pages 212–223. New York Acad. Sci., 1985.
- [Sta86] R.P. Stanley. *Enumerative Combinatorics*, Volume I. Wadsworth, Monterey, 1986.
- [Sta87a] R.P. Stanley. On the number of faces of centrally-symmetric simplicial polytopes. *Graphs Combin.*, 3:55–66, 1987.
- [Sta87b] R.P. Stanley. Generalized  $h$ -vectors, intersection cohomology of toric varieties, and related results. In M. Nagata and H. Matsumura, editors, *Commutative Algebra and Combinatorics*, volume 11 of *Adv. Stud. Pure Math.*, pages 187–213. Kinokuniya, Tokyo and North-Holland, Amsterdam/New York, 1987.
- [Sta96] R.P. Stanley. *Combinatorics and Commutative Algebra*, 2nd Ed. Volume 41 of *Progr. Math.*, Birkhäuser, Boston, 1996.
- [Zie95] G.M. Ziegler. *Lectures on Polytopes*. Volume 152 of *Graduate Texts in Math.*, Springer-Verlag, New York, 1995.