

# The Associahedron and Triangulations of the $n$ -gon

Carl W. Lee<sup>1</sup>

Department of Mathematics

University of Kentucky

Lexington, KY 40506

USA

Revised March 17, 1988

---

<sup>1</sup>Research supported, in part, by NSF grants MCS-8201653 and DMS-8504050, and an Alexander von Humboldt Foundation Fellowship.

**ABSTRACT:** Let  $P_n$  be a convex  $n$ -gon in the plane,  $n \geq 3$ . Consider  $\Sigma_n$ , the collection of all sets of mutually noncrossing diagonals of  $P_n$ . Then  $\Sigma_n$  is a simplicial complex of dimension  $n - 4$ . We prove that  $\Sigma_n$  is isomorphic to the boundary complex of some  $(n - 3)$ -dimensional simplicial convex polytope and that this polytope can be geometrically realized to have the dihedral group  $D_n$  as its group of symmetries. Formulas for the  $f$ -vector and  $h$ -vector of this polytope and some implications for related combinatorial problems are discussed.

# 1 Introduction

Let  $P_n$  be a convex  $n$ -gon in the plane,  $n \geq 3$ . Apart from the  $n$  edges of  $P_n$ , the  $n$ -gon has  $\binom{n}{2} - n = n(n-3)/2$  diagonals. Two different diagonals are said to *cross* if they intersect at a point other than, possibly, a common endpoint. Consider  $\Sigma_n$ , the collection of all sets of mutually noncrossing diagonals. The maximum size of such a set is  $n-3$ . We may therefore regard  $\Sigma_n$  as a simplicial complex of dimension  $n-4$ , having  $n(n-3)/2$  vertices.

Perles [12] asked whether  $\Sigma_n$  is isomorphic to the boundary complex of some  $(n-3)$ -dimensional simplicial polytope. He cited Huguet and Tamari [8] in which a related polytopal object was discussed. Because maximum sets in  $\Sigma_n$  correspond to triangulations of  $P_n$ , we seek an  $(n-3)$ -dimensional polytope  $Q_n$  with one vertex for each diagonal of  $P_n$  and one facet for each triangulation of  $P_n$ . In this paper we show that such a polytope exists. We then consider formulas for the  $f$ -vector and  $h$ -vector of this polytope and discuss some implications for related combinatorial problems, which we list at the end of Section 6.

Haiman [7] independently solved Perles' problem by constructing the dual of the desired  $Q_n$ , obtaining a defining set of inequalities, one for each diagonal of the  $n$ -gon. Because of the correspondence between triangulations of the  $n$ -gon and ways of parenthesizing a sequence of  $n-1$  symbols, we will adopt Haiman's designation and refer to any polytope combinatorially equivalent to  $Q_n$  as the  $(n-3)$ -dimensional *associahedron*. Recall that the

number of triangulations of the  $n$ -gon, and hence the number of facets of  $Q_n$ , is the  $(n-1)^{\text{st}}$  Catalan number  $c_{n-1} = \frac{1}{n-1} \binom{2n-4}{n-2}$ ,  $n \geq 2$ . See Gardner [5] for a pleasant introduction to this often-encountered sequence.

## 2 Simplicial Complexes

For convenience we review some properties of simplicial complexes. A *simplicial complex*  $\Delta$  is a nonempty collection of subsets of a finite set  $V$  with the property that  $F \in \Delta$  whenever  $F \subseteq G$  for some  $G \in \Delta$ . For  $F \in \Delta$  we say  $F$  is a *face* of  $\Delta$  and the *dimension* of  $F$ ,  $\dim F$ , equals  $(\text{card } F) - 1$ . The *dimension* of  $\Delta$ ,  $\dim \Delta$ , is defined to be  $\max\{\dim F : F \in \Delta\}$ . Faces of  $\Delta$  of dimension 0, 1,  $(\dim \Delta) - 1$  and  $\dim \Delta$  are called *vertices*, *edges*, *subfacets* and *facets* of  $\Delta$ , respectively. For any finite set  $F$ , the set of all subsets of  $F$  will be denoted  $\overline{F}$ , and the set of all proper subsets of  $F$  will be denoted  $\partial\overline{F}$ . We will write  $v_1 v_2 \cdots v_k$  as an abbreviation for the set  $\{v_1, v_2, \dots, v_k\}$  and will write  $\overline{v}$  as an abbreviation for  $\overline{\{v\}}$ .

Let  $\Delta$  be a simplicial complex. If  $F \in \Delta$ , the *link* of  $F$  in  $\Delta$  is the simplicial complex  $\text{lk}_\Delta F = \{G \in \Delta : G \cap F = \emptyset, G \cup F \in \Delta\}$ . If  $F \neq \emptyset$ , the *deletion* of  $F$  from  $\Delta$  is the simplicial complex  $\Delta \setminus F = \{G \in \Delta : F \not\subseteq G\}$ .

Let  $\Delta_1$  and  $\Delta_2$  be simplicial complexes with disjoint sets of vertices. The *join* of  $\Delta_1$  and  $\Delta_2$  is the simplicial complex  $\Delta_1 \cdot \Delta_2 = \{F_1 \cup F_2 : F_1 \in \Delta_1, F_2 \in \Delta_2\}$ . Suppose  $F \neq \emptyset$  is a face of a simplicial complex  $\Delta$ . Then the *stellar subdivision* of  $F$  in  $\Delta$  is the simplicial

complex  $\text{st}(v, F)[\Delta] = (\Delta \setminus F) \cup (\bar{v} \cdot \partial \bar{F} \cdot \text{lk}_\Delta F)$ , where  $v$  is a new vertex that is not a vertex of  $\Delta$ . Note that during a stellar subdivision, the only old faces of  $\Delta$  that are lost are those containing  $F$ , and the only new ones that are created are those containing  $v$ . We also observe that if  $F$  itself is a vertex, then  $\text{st}(v, F)[\Delta]$  is isomorphic to  $\Delta$ , the vertex  $F$  simply being relabeled.

If a simplicial complex  $\Delta$  is *polytopal*, i.e., if  $\Delta$  is isomorphic to the boundary complex  $\Sigma(P)$  of some simplicial convex polytope  $P$ , then so is  $\text{st}(v, F)[\Delta]$  for any  $\emptyset \neq F \in \Delta$ . One can, for example, choose a point  $v$  just "above" the centroid of the face of  $P$  corresponding to  $F$ , and form the polytope  $Q = \text{conv}(P \cup \{v\})$ , where *conv* means *convex hull*. Then  $\text{st}(v, F)[\Delta]$  is isomorphic to  $\Sigma(Q)$ .

It is easy to verify the next lemma.

**Lemma 1** *Let  $\Delta_1, \Delta_2, \dots, \Delta_{m+1}$  be a sequence of simplicial complexes,  $F_1, F_2, \dots, F_m$  be a sequence of faces, and  $v_1, v_2, \dots, v_m$  be a sequence of vertices, such that  $\Delta_{i+1} = \text{st}(v_i, F_i)[\Delta_i]$ ,  $1 \leq i \leq m$ . Suppose in addition we assume that for particular numbers  $j$  and  $k$ ,  $1 \leq j < k \leq m$ , we have  $F_k \in \Delta_j$  and  $F_j \cup F_k \notin \Delta_j$ . Then  $v_j v_k \notin \Delta_{m+1}$ .*

### 3 Constructing the Associahedron

Assume  $n \geq 4$  and number the vertices of  $P_n$  from 0 to  $n - 1$  consecutively around the perimeter. Let  $S$  be the collection of all sets of consecutive integers of the form  $\{i, i+1, \dots, j\}$ ,

where  $1 \leq i \leq j \leq n - 2$ , excluding the set  $\{1, 2, \dots, n - 2\}$ . If we associate each such set with the diagonal of  $P_n$  joining vertices  $i - 1$  and  $j + 1$ , we establish a bijection between the members of  $S$  and the diagonals of the  $n$ -gon.

Let  $\Delta_1$  be the boundary complex of any  $(n - 3)$ -dimensional geometric simplex in  $\mathbb{R}^{n-3}$  and number the vertices of  $\Delta_1$  from 1 to  $n - 2$ . The members of  $S$  now correspond to certain faces of  $\Delta_1$ . Order the members of  $S$ ,  $F_1, F_2, \dots, F_m$ , so that  $i < j$  whenever  $F_i \supset F_j$ . Set  $\Delta_{i+1} = \text{st}(v_i, F_i)[\Delta_i]$ ,  $1 \leq i \leq m$ , where  $v_i$  is not a vertex of  $\Delta_i$ . Note that when  $F_j$  is subdivided, only faces containing it are lost, so that  $F_{j+1}, F_{j+2}, \dots, F_m$  are not lost, and hence the  $\Delta_i$  are well-defined. We remark also that the singleton sets in  $S$  correspond precisely to the original vertices of  $\Delta_1$ , which need not, therefore, be subdivided.

In this manner we obtain  $\Delta_{m+1}$ , which we call  $\Delta^*$  for short, whose vertices are in one-to-one correspondence with the diagonals of  $P_n$ . The fact that  $\Delta^*$  is polytopal is clear since it is obtained from the boundary complex of an  $(n - 3)$ -dimensional polytope (namely, a simplex) by a sequence of stellar subdivisions. So  $\Delta^*$  is isomorphic to  $\Sigma(Q_n)$  for some simplicial polytope  $Q_n$ . We will show that  $\Delta^*$  is isomorphic to  $\Sigma_n$ , and hence that  $Q_n$  is the desired associahedron. Low values of  $n$ , say,  $4 \leq n \leq 6$ , can be checked directly; the procedure even works formally for  $n = 3$ , yielding a 0-dimensional polytope  $Q_3$  with  $\Sigma(Q_3) = \{\emptyset\} = \Sigma_3$ . See Figure 1.

INSERT FIGURE 1 NEAR HERE

The first step in showing that  $\Delta^*$  is isomorphic to  $\Sigma_n$  will be to prove that if  $u$  and  $v$  are vertices of  $\Delta^*$  corresponding to crossing diagonals of  $P_n$ , then  $uv$  is not an edge of  $\Delta^*$ . For suppose  $u$  and  $v$  correspond to the sets  $F = \{p, p+1, \dots, q\}$  and  $G = \{r, r+1, \dots, s\}$  in  $S$ , respectively. If the associated diagonals cross, it is easy to see that we may assume  $p < r$ ,  $q < s$  and  $r \leq q+1$ . Hence  $H = \{p, p+1, \dots, s\}$  is a set of consecutive integers containing  $F$  and  $G$  strictly. If  $H = \{1, 2, \dots, n-2\}$  then  $H$  is not a face of  $\Delta_1$ , and so  $uv \notin \Delta^*$  by Lemma 1. If  $H \neq \{1, 2, \dots, n-2\}$  then  $H \in S$  and  $H$  is subdivided before both  $F$  and  $G$ . After its subdivision  $H = F \cup G$  is no longer a face, and Lemma 1 again implies that  $uv \notin \Delta^*$ .

We now know that every face of  $\Delta^*$  corresponds to a set of noncrossing diagonals of  $P_n$ . In particular, each facet of  $\Delta^*$  represents a triangulation of the  $n$ -gon and so corresponds to a facet of  $\Sigma_n$ . To show the converse, it is sufficient to note that the following two properties hold for both  $\Delta^*$  and  $\Sigma_n$ : (1) Every subfacet is contained in exactly two facets. (2) Between every pair of facets  $F$  and  $G$  there is a path  $F = F_1, F_2, \dots, F_k = G$  of facets such that  $F_i$  and  $F_{i+1}$  share a common subfacet,  $i = 1, \dots, k-1$ . From this we can conclude that every facet of  $\Sigma_n$  corresponds to one in  $\Delta^*$ . Therefore there is one facet of  $\Delta^*$  for every triangulation of  $P_n$ ,  $\Delta^*$  is isomorphic to  $\Sigma_n$ , and  $Q_n$  is the  $(n-3)$ -dimensional associahedron, establishing

the following theorem<sup>2</sup>.

**Theorem 1** *Let  $\Sigma_n$  be the simplicial complex consisting of the collection of all sets of mutually noncrossing diagonals of the  $n$ -gon. Then  $\Sigma_n$  is realizable as the boundary complex of an  $(n - 3)$ -dimensional simplicial polytope  $Q_n$ .*

## 4 The Associahedron and Gale Diagrams

In this section we describe another way to verify that  $\Sigma_n$  is polytopal which will eventually lead to a realization of  $Q_n$  that geometrically reflects the symmetry of the regular  $n$ -gon. Our primary tool will be that of Gale transforms and Gale diagrams. We refer the reader to Grünbaum [6] and McMullen-Shephard [11] for definitions and explanations of any properties of Gale diagrams we may subsequently use.

Assume  $n \geq 5$  and consider any convex  $n$ -gon  $P_n$  (not necessarily regular) with the vertices again numbered from 0 to  $n - 1$ . Let  $X'$  denote this set of vertices and choose a point  $O$  in the interior of  $P_n$  such that  $O$  satisfies at least one of the following two conditions:

1.  $O$  is in the interior of  $\text{conv}(X' \setminus \{x'\})$  for all  $x' \in X'$ .

---

<sup>2</sup>We thank Gil Kalai and Micha Perles for pointing out this argument for the converse. The original argument showed by induction that the facet  $F = \{1, 2, \dots, n - 2\} \setminus \{j\}$  of  $\Delta_1$  was ultimately subdivided into  $c_j c_{n-j-1}$  facets of  $\Delta^*$ ,  $1 \leq j \leq n - 2$ . Then the identity  $\sum_{j=1}^{n-2} c_j c_{n-j-1} = c_{n-1}$  verifies that all of the facets of  $\Sigma_n$  are present in  $\Delta^*$ , offering a nice geometric manifestation of the Catalan recurrence relation.



2.  $O$  lies on no diagonal of  $P_n$ .

Establish a Cartesian coordinate system for the plane such that the origin is at  $O$ . Vertex  $i$  of the  $n$ -gon can then be thought of as a vector  $x'_i$  in  $\mathbb{R}^2$ ,  $0 \leq i \leq n-1$ . Because  $O$  is in the interior of  $P_n$ , there exist positive numbers  $\lambda_i$ ,  $0 \leq i \leq n-1$ , such that  $\sum_{i=0}^{n-1} \lambda_i x'_i = 0$ . This says that  $O$  is the centroid of the vectors  $\lambda_i x'_i$  and implies that the original points  $x'_i$  constitute the Gale diagram of some set of  $n$  points  $X = \{x_0, x_1, \dots, x_{n-1}\}$  in  $\mathbb{R}^{n-3}$  such that  $\text{conv}(X)$  is a (not necessarily simplicial)  $(n-3)$ -dimensional polytope. We remark that some of the points in  $X$  may not be vertices of the polytope. There is a natural correspondence between the element  $x_i$  of  $X$  and the element  $i (= x'_i)$  of  $X'$ ,  $0 \leq i \leq n-1$ , which induces the obvious correspondence between subsets  $Y$  of  $X$  and  $Y'$  of  $X'$ .

Let  $\Psi$  be the boundary complex of this polytope. The Gale diagram has the property that for every  $Y \subseteq X$  we have  $Y \in \Psi$  if and only if  $O$  is in the relative interior of  $\text{conv}(X' \setminus Y')$ , which we write  $O \in \text{relint conv}(X' \setminus Y')$ .

We now consider the facets, i.e., the maximal faces of  $\Psi$ . It is readily seen that  $F \subseteq X$  is a facet of  $\Psi$  if and only if  $X' \setminus F'$  is the set of vertices of a triangle  $T$  or a diagonal  $D$  containing  $O$  in its relative interior. In the first case  $\dim \text{conv}(F) = n-4$  and  $\text{card } F = n-3$ , and so  $\text{conv}(F)$  is a simplex.

In the second case  $\dim \text{conv}(F) = n-4$  but  $\text{card } F = n-2$ , and so  $\text{conv}(F)$  is not

a simplex. Suppose  $D$  has endpoints  $i$  and  $j$ . Let  $G'_1 = \{i+1, i+2, \dots, j-1\}$  and  $G'_2 = \{j+1, j+2, \dots, n-1, 0, 1, \dots, i-1\}$ . It is easy to check that the only proper supersets  $H'$  of  $\{i, j\}$  for which  $O \in \text{relint conv}(H')$  are the sets of the form  $H' = \{i, j\} \cup H'_1 \cup H'_2$ , where  $H'_i$  is a nonempty subset of  $G'_i$ ,  $i = 1, 2$ . This immediately implies that the boundary complex of the facet  $\text{conv}(F)$  is the simplicial complex  $\partial \overline{G}_1 \cdot \partial \overline{G}_2$ , and that with the exception of such nonsimplicial facets  $F$ , every face of every dimension of  $\Psi$  corresponds to a simplex.

To construct the associahedron, we begin by subdividing each nonsimplicial facet  $F$  in a manner analogous to stellar subdivision by removing  $F$  and adding all faces of the form  $\{v\} \cup G$ , where  $G \in \partial \overline{G}_1 \cdot \partial \overline{G}_2$ . When this is done for every such  $F$ ,  $\Psi$  is transformed into a simplicial complex  $\Psi_1$ . The same argument as for stellar subdivisions shows that  $\Psi_1$  is polytopal: we can place a point  $v$  just "above" the centroid of  $\text{conv}(F)$  and take the convex hull. Note that apart from the nonsimplicial facets of  $\Psi$ , no other face of  $\Psi$  is lost.

A proper subset of vertices of  $X'$  will be called *consecutive* if it is a set of consecutive integers, mod  $n$ . Consider any diagonal of  $P_n$  not containing the origin. When extended, the diagonal determines two open half-planes, one of which contains  $O$ . Associate with the diagonal the set  $F'$  of consecutive vertices in the opposite open half-space. Let  $S'$  be the collection of all subsets of  $X'$  derived in this way. We then have a bijection between the members of  $S'$  and the diagonals of  $P_n$  not containing the origin. We observe that if  $F' \in S'$ , then every consecutive subset of  $F'$  is also in  $S'$ . Further, if  $G'_i$  is one of the two

consecutive sets associated with a diagonal containing  $O$  as previously described, then every proper consecutive subset of  $G'_i$  is in  $S'$ .

By the property of Gale diagrams, every member of  $S'$  corresponds to a face of  $\Psi$ , and hence of  $\Psi_1$ . Note in particular that the singleton sets in  $S'$  correspond precisely to the original vertices of  $\Psi$ . Order the faces of  $\Psi_1$  associated with the members of  $S'$ ,  $F_1, F_2, \dots, F_r$ , so that  $i < j$  whenever  $F_i \supset F_j$ , and set  $\Psi_{i+1} = \text{st}(v_i, F_i)[\Psi_i]$ ,  $1 \leq i \leq r$ . Once again we obtain a polytopal simplicial complex  $\Psi^* = \Psi_{r+1}$  whose vertices are in one-to-one correspondence with the diagonals of the  $n$ -gon. See Figure 2. The argument showing  $\Psi^*$  is isomorphic to  $\Sigma_n$  will parallel the discussion of the previous section.

*INSERT FIGURE 2 NEAR HERE*

Suppose  $u$  and  $v$  are vertices of  $\Psi^*$  associated with crossing diagonals  $D$  and  $E$ , respectively. If  $D$  and  $E$  both contain  $O$ , then  $u$  and  $v$  were introduced to triangulate two distinct nonsimplicial facets of  $\Psi$ . Hence  $uv \notin \Psi_1$  and so  $uv \notin \Psi^*$ . Suppose  $O \in D$  but  $O \notin E$ . The only way we could have  $uv \in \Psi^*$  is if  $F \in \text{lk}_{\Psi_1} u$ , where  $F$  is the face subdivided by  $v$ . But  $\text{lk}_{\Psi_1} u = \partial \overline{G}_1 \cdot \partial \overline{G}_2$ , where  $G'_1$  and  $G'_2$  are the two consecutive sets defined by the two open half-planes associated with  $D$ . Hence  $F' \subseteq G'_1$  or  $F' \subseteq G'_2$ , and in either case  $D$  and  $E$  cannot cross.

Finally, suppose neither  $D$  nor  $E$  contain  $O$ . If  $u$  and  $v$  correspond to consecutive sets

$F'$  and  $G'$ , respectively, then one can verify that  $H' = F' \cup G'$  is a set of consecutive vertices strictly containing both  $F'$  and  $G'$ . If  $H$  is not a face of  $\Psi_1$  then  $uv \notin \Psi^*$  by Lemma 1. If  $H$  is a face of  $\Psi_1$  then  $H$  is a face of  $\Psi$  and it is easy to see that  $H'$  must also be in  $S'$ . Hence  $H$  is subdivided before both  $F$  and  $G$ . When  $H$  is subdivided, then  $F \cup G$  is no longer a face, so once again  $uv \notin \Psi^*$ .

We now know that every facet of  $\Psi^*$  corresponds to a facet of  $\Sigma_n$ . The proof of the converse is identical to the previous argument for  $\Delta^*$ . Hence  $\Sigma_n$  is isomorphic to  $\Psi^*$  and thus to  $\Sigma(Q_n)$  for some simplicial  $(n-3)$ -polytope  $Q_n$ . The above construction includes the construction of the previous section as a special case. One need only choose  $O$  to be suitably near a point in the relative interior of the edge joining 0 and  $n-1$ .

Since the boundary complex of any  $(n-3)$ -polytope with at most  $n$  vertices can be refined to that of a simplicial  $(n-3)$ -polytope with  $n$  vertices, and since every such simplicial polytope has a Gale diagram consisting of a convex  $n$ -gon with origin  $O$  in its interior satisfying condition (2), we have the following result.

**Theorem 2** *For any  $(n-3)$ -polytope  $P$  with at most  $n$  vertices, there exists a refinement of the boundary complex that is isomorphic to  $\Sigma_n$ . Moreover if  $P$  is simplicial, the refinement is achievable by a sequence of stellar subdivisions.*

## 5 Symmetrical Realizations

We will now determine a realization of  $Q_n$  that geometrically reflects the symmetry of the regular  $n$ -gon. Specifically, we will construct  $Q_n$  in such a way that its symmetry group is isomorphic to the dihedral group  $D_n$ . Suppose  $P_n$  is a regular  $n$ -gon with vertex  $j$  having coordinates  $(\cos j\theta, \sin j\theta)$ ,  $0 \leq j \leq n-1$ , where  $\theta = 2\pi/n$ . The dihedral group is generated by elements  $g_1$  and  $g_2$ , where  $g_1(j) = j+1 \pmod{n}$  and  $g_2(j) = n-j \pmod{n}$ ,  $0 \leq j \leq n-1$ .

Because in the above situation the origin  $O$  is the centroid of the vertices of  $P_n$ , we in fact have a Gale diagram that is a Gale transform of some  $(n-3)$ -polytope  $R_n$  if  $n \geq 5$ . Moreover,  $R_n$  has  $n$  vertices  $x_0, x_1, \dots, x_{n-1}$  which are in one-to-one correspondence with the vertices  $0, 1, \dots, n-1$  of the  $n$ -gon.

To find the coordinates of the vertices of  $R_n$ , we first consider the set of  $n$  nonzero vectors  $\{u^0, u^1, \dots, u^{\lfloor \frac{n}{2} \rfloor}, v^1, v^2, \dots, v^{\lfloor \frac{n-1}{2} \rfloor}\}$ , where  $\lfloor \cdot \rfloor$  denotes the integer rounddown function, defined by

$$\begin{aligned} u^k &= (u_0^k, u_1^k, \dots, u_{n-1}^k), \quad 0 \leq k \leq \lfloor \frac{n}{2} \rfloor, \\ u_j^k &= \cos kj\theta, \quad 0 \leq j \leq n-1, \\ v^k &= (v_0^k, v_1^k, \dots, v_{n-1}^k), \quad 1 \leq k \leq \lfloor \frac{n-1}{2} \rfloor, \\ v_j^k &= \sin kj\theta, \quad 0 \leq j \leq n-1. \end{aligned}$$

Note in particular that

$$u^0 = (1, 1, \dots, 1),$$

$$u^1 = (\cos 0\theta, \cos 1\theta, \dots, \cos(n-1)\theta),$$

$$v^1 = (\sin 0\theta, \sin 1\theta, \dots, \sin(n-1)\theta), \text{ and}$$

$$u^{\lfloor \frac{n}{2} \rfloor} = (1, -1, 1, -1, \dots, -1) \text{ if } n \text{ is even.}$$

Using the fact that  $\sum_{j=0}^{n-1} \omega^{mj} = 0$  if  $n$  does not divide  $m$ , where  $\omega$  is the complex  $n^{\text{th}}$  root of unity  $\cos \theta + i \sin \theta$ , and other elementary trigonometric identities, it is easy to check that we have a set of  $n$  nonzero mutually orthogonal vectors, one of which is the vector  $(1, 1, \dots, 1)$ .

If we list vectors  $u^1$  and  $v^1$  as the rows of a  $2 \times n$  matrix, the columns provide the coordinates of the regular  $n$ -gon. This implies that if we list all of our vectors except  $u^0 = (1, 1, \dots, 1)$ ,  $u^1$  and  $v^1$  as the rows of an  $(n-3) \times n$  matrix, the columns of the matrix provide the coordinates of  $x_0, x_1, \dots, x_{n-1}$ , respectively. Thus we may take

$$x_j = (\cos 2j\theta, \sin 2j\theta, \dots, \cos(\frac{n-1}{2}j\theta, \sin(\frac{n-1}{2}j\theta), \quad 0 \leq j \leq n-1, \text{ if } n \text{ is odd, and}$$

$$x_j = (\cos 2j\theta, \sin 2j\theta, \dots, \cos(\frac{n-2}{2}j\theta, \sin(\frac{n-2}{2}j\theta, (-1)^j), \quad 0 \leq j \leq n-1, \text{ if } n \text{ is even.}$$

In the former case  $R_n$  is a cyclic  $(n-3)$ -polytope, and in the latter case  $R_n$  is the projection of a cyclic  $(n-2)$ -polytope.

Suppose  $n$  is odd. Define  $g'_1$  to be the  $(n-3) \times (n-3)$  matrix  $\text{diag}(B_2, B_3, \dots, B_{(\frac{n-1}{2})})$

where  $B_k$  is the  $2 \times 2$  block

$$\begin{bmatrix} \cos k\theta & -\sin k\theta \\ \sin k\theta & \cos k\theta \end{bmatrix}.$$

If  $n$  is even, define  $g'_1$  to be the  $(n-3) \times (n-3)$  matrix  $\text{diag}(B_2, B_3, \dots, B_{(\frac{n-2}{2})}, -1)$  with the  $2 \times 2$  blocks  $B_k$  defined in the same way. Whatever the parity of  $n$ , define  $g'_2$  to be the  $(n-3) \times (n-3)$  matrix  $\text{diag}(1, -1, \dots, (-1)^{n-2})$ . It is easy to check that  $g'_1$  and  $g'_2$  generate the group of orthogonal symmetries of  $R_n$  isomorphic to the dihedral group, where  $g'_1(x_j) = x_{j+1 \pmod n}$  and  $g'_2(x_j) = x_{n-j \pmod n}$ ,  $0 \leq j \leq n-1$ .

It is also straightforward to verify that every face of  $R_n$  to be subdivided is mapped by any element of the group onto another such face, and that centroids are mapped onto centroids. Therefore all the necessary subdivisions to the boundary complex of  $R_n$  can be carried out geometrically in such a way that the group is also the group of symmetries of the resulting associahedron  $Q_n$ . For example, if a face  $F$  with centroid  $y$  is to be subdivided via a vertex  $z$ , choose  $z = (1 + \epsilon)y$  where  $\epsilon$  is a suitably small positive number taken to be the same for all faces in the orbit of  $F$ . See Figure 3.

*INSERT FIGURE 3 NEAR HERE*

## 6 The $f$ -vector and $h$ -vector of the Associahedron

In this section we investigate the number of  $j$ -dimensional faces  $f_j$ ,  $0 \leq j \leq n-4$ , of the  $(n-3)$ -dimensional polytope  $Q_n$ . Of course, we know that  $f_j$  equals the number of ways of choosing a set of  $j+1$  mutually noncrossing diagonals of the convex  $n$ -gon  $P_n$ . In particular,  $f_{n-4} = c_{n-1}$ . The  $f$ -vector of  $Q_n$  is the vector  $f(Q_n) = (f_{-1}, f_0, f_1, \dots, f_{n-4})$ , where we take  $f_{-1} = 1$  by convention.

The  $h$ -vector of  $Q_n$  is defined by  $h(Q_n) = (h_0, h_1, \dots, h_{n-3})$ , where

$$h_i = \sum_{j=0}^i (-1)^{i-j} \binom{n-j-3}{n-i-3} f_{j-1}, \quad 0 \leq i \leq n-3, \quad (1)$$

and the  $f$ -vector can be recovered from the  $h$ -vector by

$$f_{j-1} = \sum_{i=0}^j \binom{n-i-3}{n-j-3} h_i, \quad 0 \leq j \leq n-3. \quad (2)$$

See, for example, McMullen-Shephard [11] where our  $h_i$  is their  $g_{i-1}^{(d)} = g_{i-1}^{(n-3)}$ . Past experience has shown that the  $h$ -vector is often more tractable than the  $f$ -vector, and this turns out to be the case here too.

**Theorem 3** *For the associahedron  $Q_n$ ,*

$$f_{j-1} = \frac{1}{n-1} \binom{n-3}{j} \binom{n+j-1}{j+1}, \quad 0 \leq j \leq n-3, \text{ and}$$

$$h_i = \frac{1}{n-1} \binom{n-3}{i} \binom{n-1}{i+1}, \quad 0 \leq i \leq n-3.$$



**Proof:** The first formula is that of Kirkman [9] and Cayley [2], and the second follows from (1).  $\square$

The fact that  $h_i = h_{n-3-i}$  is a manifestation of the *Dehn-Sommerville equations* (see [6,11]) which hold for any triangulated sphere.

Our next objective is to describe the components of the  $h$ -vector combinatorially. Fix any triangulation  $T$  of  $P_n$ ,  $n \geq 4$ . We will color each of its diagonals either red or green, according to the following method. Choose a diagonal  $D$  and remove it, leaving a "hole" in the shape of a quadrilateral. There are exactly two diagonals of  $P_n$  that are also diagonals of the quadrilateral. One is  $D$ ; call the other  $D'$ . Notice that  $D$  and  $D'$  are crossing, and in particular share no common endpoint. Labeling the vertices of the  $n$ -gon as before, traverse them in the order  $0, 1, \dots, n-1$ , noting for which of  $D, D'$  you encounter an endpoint first. If  $D$  is met first, color  $D$  green; otherwise color it red.

We now observe that given any set of mutually noncrossing diagonals of  $P_n$  (not necessarily a triangulation) there is exactly one way to complete the set to a triangulation  $T$  such that every newly added diagonal is green in  $T$ . For suppose we have not yet completed the set to a triangulation. Then there is at least one convex  $m$ -gon,  $m \geq 4$ , in this subdivision, bounded by diagonals from the set and sides of  $P_n$ . Let its vertices be  $\{i_1, i_2, \dots, i_m\}$ , where  $i_1 < i_2 < \dots < i_m$ . No new green diagonal in a triangulation extending the given set can have  $i_m$  as an endpoint; hence any such triangulation must contain the diagonal joining  $i_1$

and  $i_{m-1}$ . By repeating this argument, the uniquely determined  $T$  is constructed<sup>3</sup>.

**Theorem 4** *For the associahedron  $Q_n$ ,  $h_i$  equals the number of triangulations of  $P_n$  having exactly  $i$  red diagonals.*

**Proof:** Let  $g_i$  be the number of triangulations with exactly  $i$  red diagonals. Let  $F$  be any set of  $j$  mutually noncrossing diagonals of  $P_n$ . There is exactly one way to complete  $F$  to a triangulation so that all of the  $n - j - 3$  new diagonals are green. This means we can count the number of such  $F$  by counting the number of ways we can choose a triangulation with exactly  $i$  red diagonals,  $i \leq j$ , and then remove  $n - j - 3$  of the  $n - i - 3$  green diagonals.

Thus

$$f_{j-1} = \sum_{i=0}^j \binom{n-i-3}{n-j-3} g_i, \quad 0 \leq j \leq n-3.$$

Formulas (1) and (2) immediately imply  $g_i = h_i$ ,  $0 \leq i \leq n-3$ .  $\square$

The Dehn-Sommerville equations are a consequence of being able to interchange the colors green and red. For a dual version of this type of counting argument, see Brøndsted [1].

The components of  $h(Q_n)$  can be interpreted in terms of some of the many problems isomorphic to that of triangulating an  $n$ -gon [5]:

---

<sup>3</sup>This argument, suggested by a referee, is essentially isomorphic to our original argument but avoids recasting the problem in terms of parenthesizing a sequence of  $n-1$  symbols.

1. Consider all ways of completely parenthesizing a sequence of  $n - 1$  symbols using  $n - 2$  pairs of parentheses. Then  $h_i$  equals the number of parenthesizations containing exactly  $i$  internal groups of left (respectively right) parentheses. Modifying the technique discussed in [4] to obtain the formula for the Catalan numbers, one can exploit this isomorphism to derive the formula for  $h_i$  directly, from which the formula for  $f_{j-1}$  is an easy corollary.
2. Consider all sequences of length  $2n - 4$  composed of  $n - 2$  zeros and  $n - 2$  ones, such that at no position along the sequence have you encountered more zeros than ones. Then  $h_i$  equals the number of sequences with  $i + 1$  blocks of ones.
3. Consider all paths from the point  $(0,0)$  to the point  $(n - 2, n - 2)$  in the Cartesian plane, where only unit steps upward and to the right are allowed, and where you must never pass through a point above the line joining  $(0,0)$  and  $(n - 2, n - 2)$ . Then  $h_i$  equals the number of paths with  $i$  changes of direction from upward to right.
4. Consider all rooted, planar, trivalent trees with one root and  $n - 1$  other nodes of degree 1. Then  $h_i$  equals the number of trees with  $i$  branchings to the left (respectively right).
5. Consider all rooted, planar trees with one root and  $n - 1$  other nodes, whether of degree

one or not. Let us say there are  $k - 2$  branchings at a node of degree  $k \geq 3$ . Then  $h_i$  equals the number of trees with a total of  $i$  branchings.

Notice the appearance of the Dehn-Sommerville equations again in (1) and (4).

## 7 Concluding Remarks

We wish to mention another polytope associated with the triangulations of the  $n$ -gon. Dantzig, Hoffman and Hu [3] have shown how to describe a polytope by linear equations in nonnegative variables whose vertices correspond to the triangulations of the  $n$ -gon and whose facets correspond to the diagonals. The dual of this polytope has therefore one vertex for every diagonal and one facet for every triangulation. But this dual is not isomorphic to  $Q_n$ ; in general it is higher dimensional. It is true, however, that adjacent triangulations correspond to adjacent facets, though the converse does not hold.

Given any  $d$ -dimensional convex polytope  $P$ . One might consider the set  $\Sigma$  of all subdivisions of  $P$ , partially ordered by refinement, and ask whether  $\Sigma$  is realizable as the boundary complex of some simplicial convex polytope  $Q$  of dimension  $n - d - 1$ , with facets of  $Q$  corresponding to triangulations of  $P$ . As we have shown, this is true if  $d = 2$ . It also turns out to be true if  $n \leq d + 3$ , but fails in general (for example, when  $d = 3$  and  $n = 7$ ). Nevertheless, there always exists a nice  $(n - d - 2)$ -dimensional spherical complex of some, but not necessarily all, subdivisions of the polytope [10].

## 8 Acknowledgments

I wish to thank Mark Haiman, Gil Kalai, Micha Perles, and an anonymous referee for their helpful suggestions and comments.

## References

- [1] A. Brøndsted, *An Introduction to Convex Polytopes*, Springer-Verlag, New York, New York, USA, 1983.
- [2] A. Cayley, On the partitions of a polygon, *Proc. London Math. Soc.* (1) **22** (1890–1891), 237–262.
- [3] G.B. Dantzig, A.J. Hoffman and T.C. Hu, Triangulations (tilings) and certain block triangular matrices, *Math. Programming* **31** (1985), 1–14.
- [4] H. Dörrie, Euler's problem of polygon division, *100 Great Problems of Elementary Mathematics*, D. Antin, trans., Dover, New York, New York, USA, 1965, 21–27.
- [5] M. Gardner, Catalan numbers, *Time Travel and Other Mathematical Bewilderments*, W.H. Freeman, New York, New York, USA, 1988, 253–266.
- [6] B. Grünbaum, *Convex Polytopes*, Interscience, London, Great Britain, 1967.
- [7] M. Haiman, Constructing the associahedron, manuscript.

- [8] D. Huguet and D. Tamari, La structure polyédrale des complexes de parenthésages, Jr. *Comb., Inf. & Syst. Sci.* 3 (1978), 69–81.
- [9] T.P. Kirkman, On the  $k$ -partitions of the  $r$ -gon and  $r$ -ace, *Philos. Trans. Roy. Soc. London* 147 (1857), 217–272.
- [10] C.W. Lee, Some notes on triangulating polytopes, *Proceedings, 3. Kolloquium über Diskrete Geometrie*, Institut für Mathematik, Universität Salzburg, Salzburg, Austria, May 1985, 173–181.
- [11] P. McMullen and G.C. Shephard, *Convex Polytopes and the Upper Bound Conjecture*, Cambridge Univ. Press, Cambridge, Great Britain, 1971.
- [12] M.A. Perles, List of problems, Conference on Convexity, Oberwolfach, Federal Republic of Germany, July 1984.




Figure 1




Figure 2






Figure 3