

to $N = K$ can be deduced from (27) to be

$$(29) \quad \begin{array}{cccc} U & L & U & U \\ U & L & L & U \\ U & U & L & U. \end{array}$$

According to § 7, patterns that are in the list (26) but not in (29) correspond to hanging keyblocks—in this case all three do. Likewise, all three patterns in (27), and hence also those in (28), also correspond to hanging keyblocks.

Finally, we observe that if the excavation is below, rather than above, the sloping excavation face, so that (26) is replaced by the list

$$(30) \quad \begin{array}{cccc} U & L & L & U \\ U & L & L & L \\ U & U & L & U, \end{array}$$

then there is no overlap with (27), so that in this case no blocks that intersect the corner are keyblocks. The pattern $U L L L$ is the only one from (30) that is not in (29), and hence it represents the only hanging keyblock in this case.

Appendix—Statement of Tucker's Theorem of the Alternative. For the convenience of readers we give a statement of Tucker's Theorem of the Alternative, as presented in [3, p. 29].

Let B, C and D be given matrices, with B being nonsingular. Then either the system $Bx \geq 0$ but not all zero, $Cx \leq 0$, $Dx = 0$ has a solution x or the system $\lambda'B + \mu'C + \nu'D = 0$, $\lambda > 0$, $\mu \geq 0$ has a solution λ, μ, ν , but never both.

This theorem was first given by Tucker in [7].

Acknowledgments. The authors are grateful to Messrs W. N. Merten, W. L. van Heerden and F. A. Vreede of the National Mechanical Engineering Research Institute of the CSIR, Pretoria for stimulating discussions on the practical importance of the Shi-Goodman work, and for prompting our interest in the keyblock identification problem.

REFERENCES

- [1] S. A. CARTNEY, *The ubiquitous joint method*, in *Cavern Design at Dinorwic Power Station*, Tunnels and Tunneling, 1971, pp. 54–57.
- [2] G. F. HADLEY, *Linear Programming*, Addison-Wesley, Reading, MA, 1962.
- [3] O. L. MANGASARIAN, *Nonlinear Programming*, McGraw-Hill, New York, 1969.
- [4] R. T. ROCKAFELLAR, *Convex Analysis*, Princeton Univ. Press, Princeton, NJ, 1970.
- [5] G. H. SHI AND R. E. GOODMAN, *Underground support design using block theory to determine keyblock holding requirements*, Proc. SANGORM Symposium on Rock Mechanics in the Design of Tunnels, Pretoria, 1965, pp. 81–105.
- [6] ———, *A new concept for support of underground and surface excavations in discontinuous rocks based on a keystone principle*, Proc. 22nd U.S. Symposium on Rock Mechanics, Massachusetts Institute of Technology, Cambridge, MA, 1961, pp. 290–296.
- [7] A. W. TUCKER, *Dual systems of homogeneous linear relations*, in H. W. Kuhn and A. W. Tucker, *Linear Inequalities and Related Systems*, Annals of Mathematics Studies No. 38, Princeton Univ. Press, Princeton, NJ, 1956.

HARD ENUMERATION PROBLEMS IN GEOMETRY AND COMBINATORICS*

NATHAN LINIAL†

Abstract. A number of natural enumeration problems in geometry and combinatorics are shown to be complete in the class #P introduced by Valiant. Among others this is established for the enumeration of vertices and of faces of a polytope, acyclic orientations of a graph and satisfying assignments of implicative boolean formulas.

Key words. #P, polytopes, partial orders, acyclic orientations

AMS(MOS) subject classifications. 05C20, 52A25, 68C25

Introduction. This article contains a contribution to the theory of hard enumeration problems. The foundations of this area were laid by Valiant [Val1], [Val2] who define the class #P of enumeration problems and the subclass of problems complete in #P. The most interesting of his results is the #P-completeness of computing permanent matchings in bipartite graphs. While deciding whether a bipartite graph has a perfect matching can be done in polynomial time [H] the enumeration problem is #P-complete. Valiant's pioneering work was continued by a recent article of Provan and Ball [PB] who prove the #P-completeness of a number of natural enumeration problems.

With every enumeration problem there is an associated decision problem. Instead of asking for the number of objects in question we ask whether this number is zero or not. The decision problem associated with the computation of the permanent function is the question whether a given bipartite graph has a perfect matching. While this decision is solvable in polynomial time, it is by no means trivial. Notice, however, that for many of the problems discussed in [PB] the situation is even more extreme: Consider for example the problem of enumerating independent sets in a bipartite graph, the decision problem associated with this enumeration problem is trivial: Every graph has an independent set of vertices. So an enumeration problem can be #P-complete even if the existence problem is trivial.

This article assumes acquaintance with the theory of #P-completeness as presented in [Val1], [PB] and [GJ]. Our purpose is to present a number of natural enumeration problems which belong to the class of #P-complete problems. The problems are geometric, combinatorial and from propositional calculus.

Here is our main theorem:

THEOREM. *The following enumeration problems are #P-complete.*

- (1) *Vertices in a polytope.*
Input: A system of linear inequalities $Ax \leq b$ defining a polytope $P \subseteq \mathbb{R}^n$.
Output: The number of vertices of P .
- (2) *d-dimensional faces of a polytope (fixed d).*
Input: As in (1).
Output: The number of d-dimensional faces of P .

* Received by the editors December 10, 1984.

† Institute of Mathematics and Computer Science, Hebrew University, Jerusalem 91904, Israel.

- (3) *Facets of a polytope.*
Input: A finite set of points in \mathbb{R}^n .
Output: The number of facets ($(n-1)$ -dimensional faces) of P .
- (4) *Components of slotted space.*
Input: A set $\{H_i | i \in I\}$ of hyperplanes in \mathbb{R}^n .
Output: The number of connected components of $\{\mathbb{R}^n \setminus H_i | i \in I\}$.
- (5) *Acyclic orientations of a graph.*
Input: A graph $G = (V, E)$.
Output: The number of orientations of G with no directed circuit.
- (6) *3-colorings of a bipartite graph.*
Input: A bipartite graph $G = (A, B, E)$.
Output: The number of ways to properly color G with 3 colors.
- (7) *Satisfying assignments of an implicative Boolean formula.*
Input: A Boolean formula B on variables x_1, \dots, x_n of the form $B = \bigwedge_{i=1}^r (x_{i1} \vee x_{i2})$.
Output: Number of truth assignments for x_1, \dots, x_n which makes B true.

Proof. (1) We use the fact from [PB] that enumerating order ideals is #P-complete. Given a poset (P, \cong) with $P = [n]$ we associate with P a polytope $B = B(P)$ in \mathbb{R}^n as follows:

$$B = \{x \in \mathbb{R}^n | 1 \cong x_i \cong 0, x_i > x_j \text{ if } i \cong j \text{ in } P\}.$$

(See [Sz2], [Lj], [KS] where use is made of this polytope.) We claim that the vertices of B are in 1:1 correspondence with the order ideals of (P, \cong) . First we prove that all vertices of B have 0-1 coordinates. Let $x \in B$ have some $0 < x_i < 1$. If $\alpha = \max\{x_j | 0 < x_j < 1\}$, then by replacing all coordinates $x_j = \alpha + \epsilon$ or by $\alpha - \epsilon$ we will get a point of B . This implies that x is not a vertex of B . The correspondence between vertices and ideals is as follows:

$$x \in \text{vert}(B) \leftrightarrow S = \{1 \leq j \leq n | x_j = 0\}.$$

It is easily verified that S is an ideal and that this correspondence is bijective.

(2) Suppose that for some fixed d we can find $f_d(K)$ the number of d -dimensional faces of a polytope K . Consider r -fold pyramids P , with K as basis. In [Grn, p. 55] one finds

$$f_d(P) = \sum_i \binom{r}{i} f_{d-i}(K).$$

If we write (*) for $r = 0, \dots, d$ and have all $f_d(P)$ evaluated, then we obtain a system of equations in unknown

$$f_0(K), \dots, f_d(K).$$

This system of equations has a triangular matrix and so they can be solved successively and $f_0(K)$ can be determined in polynomial time. Since evaluating $f_0(K)$ = the number of vertices of K is #P-complete by (1), our claim follows.

(3) This is just the dual of (1). See [Grn, p. 46] for polytope duality.

(5) The proof here is based on two observations.
PROPOSITION [St1]. Let $G = (V, E)$ be a graph with n vertices and let $P(G, \lambda)$ be its chromatic polynomial. Then $(-1)^{|P(G, -1)|}$ equals the number of acyclic orientations of G .

For the other observation we have to define the operation of join of two graphs $G = (V_1, E_1), H = (V_2, E_2)$ where $V_1 \cap V_2 = \emptyset$. The join $G + H$ has $V_1 \cup V_2$ as its vertex set and

$$E_1 \cup E_2 \cup \{\{x, y\} | x \in V_1, y \in V_2\}$$

as its edge set. The following observation is immediate.

PROPOSITION. $P(G + K_n, \lambda) = \lambda(\lambda - 1) \cdot \dots \cdot (\lambda - t + 1) P(G, \lambda - t)$.
 Now we can combine these two facts as follows. Being able to enumerate acyclic orientations is equivalent to computing $P(G, -1)$ for the graph. But if we have the values of $P(G + K_n, -1)$ for $t = 1, \dots, n$, that means we can calculate the integers

$$P(G, -j) \quad (n+1 \cong j \cong 2).$$

But P is a monic polynomial of degree n so from these numbers we can compute $P(G, \lambda)$, the chromatic polynomial of G . This is a #P-complete problem because the reduction to coloring is parsimonious [GJ, p. 169].

(4) The proof here makes use of (5) that enumerating acyclic orientations is #P-complete and on the following result of Greene. $H_d \subseteq \mathbb{R}^n$ is the hyperplane given by $\{x \in \mathbb{R}^n | x_i = x_j\}$.

PROPOSITION [Gre]. Let $G = (V, E)$ be a graph on n vertices and consider

$$S(G) = \mathbb{R}^n \setminus \bigcup H_d$$

where the union is over all i, j such that $[i, j] \in E$. The number of connected components of $S(G)$ equals the number of acyclic orientations of G .

(6) We base this proof on the #P-completeness of enumerating independent sets in bipartite graphs [PB]. Let $G = (A, B, E)$ be a partite graph for which we want to find $I(G)$ the number of independent sets. Consider a graph H which is obtained by adding two new vertices a, b with a being adjacent to all vertices in $B \cup \{b\}$ and b to all vertices of $A \cup \{a\}$. Now let us compute $\chi(H, 3)$, the number of 3-colorings of H . Suppose w. l.o.g. that a, b are colored 1, 2 respectively. The 3-coloring is now uniquely defined by the set of vertices colored 3. This can be any independent set of G and so

$$\chi(H, 3) = 6I(G).$$

This proves the #P-completeness of computing $\chi(H, 3)$.

(7) This follows from #P-completeness of enumerating ideals in posets [PB]: Let (P, \cong) be a poset with $P = \{p_1, \dots, p_n\}$. Associate with it the Boolean expression

$$B = \bigwedge \{x_i \vee x_j | p_j > p_i \text{ in } P\}.$$

It is fairly easy to verify that the set of x_i which are assigned a true value in an assignment satisfying B is an ideal in (P, \cong) and that all ideals are obtained in this way.

Let us mention in closing a most intriguing problem in this field: For a poset (P, \cong) a linear extension is a 1:1 mapping $f: P \rightarrow \{1, \dots, |P|\}$ such that if $x < y$ in then $f(x) < f(y)$. Consider the problem:

Enumeration of linear extensions.

Input: A poset (P, \cong) .

Output: $L(P)$, the number of linear extensions of (P, \cong) .

Conjecture. The enumeration of linear extensions is a #P-complete problem.

A proof of this conjecture will provide a first explicit statement to the effect that computing the volume of a convex polytope is a hard computational problem. To see this we remind the reader about the polytope $B(P)$ which was used in proving part of our main theorem. We quote without proof of the following fact from [Lj]:

PROPOSITION. For a poset (P, \cong) on $|P| = p$ elements $L(P)$ the number of linear extensions of P satisfies

$$L(P) = p! \text{vol}(B(P)).$$

The connection between the #P-completeness of enumerating linear extensions and the complexity of evaluating the volume of a convex polytope is now clear.

Let us also comment about the relationship between the number of linear extensions of a poset and enumerating order ideals. We use $I(P)$ to denote the number of ideals in the poset P . For posets P, Q we define their product $P \times Q$ to be a partial order on the cartesian product of P and Q with $(x_1, y_1) \cong (x_2, y_2)$ if $x_1 \cong x_2$ in P and $y_1 \cong y_2$ in Q . A mapping $f: P \rightarrow Q$ is order preserving if $x \cong y$ in P implies $f(x) \cong f(y)$ in Q .

PROPOSITION. Let (P, \cong) be a poset and let C_i be the chain on i elements. Then $I(P \times C_i)$ equals the number of order preserving maps $f: P \rightarrow \{0, 1, \dots, i\}$.

Proof. With an ideal $J \subseteq P$ we associate a function $f: P \rightarrow \{0, \dots, i\}$ as follows: For every $x \in P$ there is unique $t \cong j \cong 0$ such that $(x, j) \in J$ and $(x, j-1) \notin J$. Let $f(x) = i-j$ for that value j . Since J is an ideal, f is well defined and easily seen to be order preserving. It is also a routine matter verifying that this correspondence is bijective.

Now we come to the expression for the number of linear extensions of a poset. THEOREM. For a poset (P, \cong) on $|P| = n$ elements, the number of linear extensions $L(P)$ satisfies

$$L(P) = I(P \times C_{n-1}) - nI(P \times C_{n-2}) + \binom{n}{2}I(P \times C_{n-3}) - \dots + \pm \binom{n}{n-2}I(P \times C_1) \mp \binom{n}{n-1}.$$

Proof. This follows from the previous proposition and Inclusion-Exclusion. Classify order preserving maps $f: P \rightarrow \{0, \dots, n-1\}$ according to their range. There are $I(P \times C_{n-1})$ such mappings altogether. Say f has property t ($n-1 \geq t \geq 0$) if t is not in the range of f . $L(P)$ is the number of order preserving maps which are onto, i.e., have no property and there are

$$\binom{n}{j} I(P + C_{n-j-1})$$

maps having a given set of j properties.

REFERENCES

[Gr] C. GREENE, *Acyclic orientations (Notes)*, in Higher Combinatorics, M. Aigner, ed., D. Reidel, Dordrecht, 1977, pp. 65-68.
 [Grn] B. GRÜNBAUM, *Convex Polytopes*, Wiley Interscience, New York, 1967.
 [GJ] M. GAREY AND D. JOHNSON, *Computers and Intractability: A Guide to the Theory of NP-Completeness*, W. H. Freeman, San Francisco, 1979.
 [H] M. HALL, JR., *An algorithm for distinctive representations*, Amer. Math. Monthly, 63 (1956), pp. 716-717.
 [KS] J. KAHN AND M. E. SAKS, *Every poset has a good comparison*, Combinatorica, to appear.
 [PB] S. PROVAN AND M. O. BALL, *On the complexity of counting cuts and of computing the probability that a graph is connected*, SIAM J. Comput., 12 (1983), pp. 777-788.
 [L] N. LINTAL, *The information theoretic bound is good for merging*, SIAM J. Comput., to appear.

[St1] R. P. STANLEY, *Acyclic orientations of graphs*, Discrete Math., 5 (1973), pp. 171-178.
 [St2] ———, *Two combinatorial applications of the Alexandroff-Fenchel inequalities*, J. Combin. Theory, ser. A, 31 (1981), pp. 56-65.
 [Val1] L. G. VALIANT, *The complexity of computing the permanent*, Theor. Comput. Sci., 8 (1979), pp. 189-201.
 [Val2] ———, *The complexity of enumeration and reliability problems*, SIAM J. Comput., 8 (1979), pp. 410-421.

13, 795-801
(1984)