

or as part of a published lecture, review or thesis); that it is not under consideration for publication elsewhere; that its publication has been approved by all coauthors, if any, as well as by the responsible authorities at the institution where the work has been carried out; that, if and when the manuscript is accepted for publication, the authors agree to automatic transfer of the copyright to the publisher; that the manuscript will not be published elsewhere in any language without the consent of the copyright holders; and that written permission of the copyright holder is obtained by the authors for the material used from other copyrighted sources.

All articles published in this journal are protected by copyright, which covers the exclusive rights to reproduce and distribute the article (e.g., as offprints), as well as all translation rights. No material published in this journal may be reproduced photographically or stored on microfilm, in electronic databases, video disks, etc., without first obtaining written permission from the publisher.

The use of general descriptive names, trade names, trademarks, etc., in this publication, even if not specifically identified, does not imply that these names are not protected by the relevant laws and regulations.

While the advice and information in this journal is believed to be true and accurate at the date of going to press, neither the authors, the editors, nor the publisher can accept any legal responsibility for any errors or omissions that may be made. The publisher makes no warranty, express or implied, with respect to the material contained herein. Springer-Verlag publishes advertisements in this journal in reliance upon the responsibility of the advertiser to comply with all legal requirements relating to the marketing and sale of products or services advertised. Springer-Verlag and the editors are not responsible for claims made in the advertisements published in the journal. The appearance of advertisements in Springer-Verlag publications does not constitute endorsement, implied or intended, of the product advertised or the claims made for it by the advertiser. Authorization to photocopy items for internal or personal use, or the internal or personal use of specific clients, is granted by Springer-Verlag New York, Inc., provided that the appropriate fee is paid directly to Copyright Clearance Center, 222 Rosewood Drive, Danvers, MA 01923, USA (Tel. 508-750-8400), stating the ISSN 0179-5376, volume, and first and last page numbers of each article copied. The copyright owner's consent does not include copying for general distribution, promotion, advertising, or resale. In these cases, specific written permission must first be obtained from the publisher.

The Canada Institute for Scientific and Technical Information (CISTI) provides a comprehensive, world-wide document delivery service for all Springer-Verlag journals. For more information, or to place an order for a copyright-cleared Springer-Verlag document, please contact Client Assistant, Document Delivery, Canada Institute for Scientific and Technical Information, Ottawa K1A 0S2, Canada (Tel: 613-993-9251; Fax: 613-952-8243; e-mail: cisti.docdel@nrc.ca).

Mathematics and computer science, covering a broad range of research articles of high quality in discrete geometry and combinatorics. DCG publishes papers on such topics as configurations, geometric complexity, polytopes, point location, geometric topology, probabilistic techniques in computational geometry, papers with a distinct geometric flavor in such areas as robotics, solid modeling, computer graphics, combinatorial VLSI design, and robotics.

Discrete & Computational Geometry, Springer-Verlag New York Inc., Journal Fulfillment Department, 233 Spring Street, New York, NY 10013-1578, USA. Distributed in North America on acid-free paper.

On Conway's Thrackle Conjecture*

L. Lovász,¹ J. Pach,² and M. Szegedy³

¹ Department of Computer Science, Yale University,
New Haven, CT 06517, USA
lovasz-laszlo@cs.elte.hu

² Department of Computer Science, City College, CUNY
and
Courant Institute, NYU,
251 Mercer Street, New York, NY 10012, USA
pach@cims.nyu.edu

³ Mathematical Sciences Research Center, AT&T Laboratories,
600 Mountain Avenue, Murray Hill, NJ 07974, USA
ms@research.att.com

Abstract. A *thrackle* is a graph drawn in the plane so that its edges are represented by Jordan arcs and any two distinct arcs either meet at exactly one common vertex or cross at exactly one point interior to both arcs. About 40 years ago, J. H. Conway conjectured that the number of edges of a thrackle cannot exceed the number of its vertices. We show that a thrackle has at most twice as many edges as vertices. Some related problems and generalizations are also considered.

1. Introduction

Let G be a graph with vertex set $V(G)$ and edge set $E(G)$, and assume that it has no loops or multiple edges. A *drawing* of G is a representation of G in the plane such that every vertex corresponds to a point, and every edge is represented by a Jordan arc connecting the corresponding two points without passing through any other vertex. Two edges (arcs) are said to *cross* each other if they have an interior point p in common. For simplicity, we always assume that no three edges cross at the same point. A crossing p is

* The first author was supported by NSF Grant CCR-94-02916. The second author was supported by NSF Grant CCR-91-22103, PSC-CUNY Research Award 663472, and OTKA-4269.

called *proper* if in a small neighborhood of p one edge passes from one side of the other edge to the other side. Due to its aesthetic appeal and wide range of applications in VLSI layout, computer-aided design, software visualization, etc., the area of graph drawings has received a lot of attention in the past two decades. For a recent bibliography of graph drawing algorithms, see [DETT].

There are many interesting results in topological graph theory characterizing all graphs embeddable on a given surface without crossing (see [WB]). However, we know very little about the possible intersection patterns determined by the edges of a graph drawn on a surface. In particular, even for some very simple graphs we do not know how to find the *crossing number* of G , i.e., the minimum number of crossing pairs of edges in a planar drawing of G . In the case when G is a complete bipartite graph, this is Turán's brick factory problem [T1], [G3]. The determination of the crossing number is known to be NP-complete [G3].

Another well-known open problem that illustrates our ignorance about graph drawings was raised by Conway about 40 years ago. He defined a *thrackle* as a drawing of a graph G with the property that any two distinct edges either:

- (i) share an endpoint, and then they do not have any other point in common, or
- (ii) do not share an endpoint, in which case they meet exactly once and determine a proper crossing.

Thrackle Conjecture. The number of edges of a thrackle cannot exceed the number of its vertices.

A graph that can be drawn as a thrackle is said to be *thrackable*. Assuming that the above conjecture is true, Woodall [W] characterized all thrackable graphs. With this assumption, a finite graph is thrackable if and only if it has at most one odd cycle, it has no cycle of length four, and each of its connected components contains at most one cycle. Note that it is quite straightforward to check the necessity of these conditions (see Lemma 2.1). Using a construction suggested by Conway, the thrackle conjecture can be reduced to the following statement: If a graph G consists of two even cycles meeting in a single vertex, then G is not thrackable ([W1] and [PRS1]). It is worth mentioning that the thrackle conjecture is true for *straight-line thrackles*, i.e., for drawings where every edge is represented by a segment [HP], [FS], [PA]. See [LST1] for a surprising relation between straight-line thrackles and triangulations of certain polytopes, and [G1] for another geometric application.

Any two edges of a thrackle intersect in exactly one point, including the endpoints. For finite set-systems satisfying a similar condition we have the following well-known result [F], [BE].

Fisher Inequality. Let F be a family of subsets of a finite set X such that any two members of F have exactly one element in common. Then F has at most as many members as the number of elements of X .

An interesting modular version of this inequality was discovered by Bertekamp [B]. Suppose that every member of F has an odd number of elements and that the intersection of any two members is even. Then $|F| \leq |X|$. These results and their generalizations

originate in linear algebra and play a central role in finite geometries and in the theory of combinatorial designs (see [BF]).

Since thrackles do not contain cycles of length four, it follows from [KST] that the maximum number of edges a thrackle of n vertices can have is $O(n^{3/2})$. Our next theorem represents a substantial improvement on this bound.

Theorem 1.1. Every thrackle of n vertices has at most $2n - 3$ edges.

The proof is based on the following result.

Theorem 1.2. Every thrackable bipartite graph is planar.

Just like the Fisher inequality, the thrackle conjecture has some modular versions, too. For example, call a graph drawing a *generalized* (or *modulo 2*) *thrackle* if any two edges meet an odd number of times, where "meet" means either "meet at a common vertex" or "meet at a proper crossing."

Theorem 1.3. Every generalized thrackle of n vertices has at most $3n - 4$ edges.

Theorem 1.4. A bipartite graph can be drawn as a generalized thrackle if and only if it is planar.

Woodall [W2] asked whether the thrackle conjecture remains true for generalized thrackles. Our last theorem implies that the answer to this question is in the negative, because a bipartite planar graph of n vertices can have as many as $2n - 4$ edges.

2. Three Lemmas

In what follows a thrackle and its underlying "abstract" graph are both denoted by G . If there is no danger of confusion, we make no notational distinction between a vertex (edge) of the graph and the corresponding point (arc).

Lemma 2.1. Let G be a thrackable graph. Then G contains (i) no cycle of length four, (ii) no two vertex-disjoint odd cycles.

Proof. To show (ii), notice that a pair of vertex-disjoint odd cycles would be represented in a thrackle by two closed curves that properly cross each other an odd number of times. \square

Lemma 2.2. Let C_1 and C_2 be two cycles in a graph G that have precisely one vertex v in common. Suppose that G can be drawn as a thrackle. Then the two closed curves representing C_1 and C_2 cross each other in a small neighborhood of v if and only if both cycles are odd.

Proof. Let k_i denote the length of C_i , $i = 1, 2$. The closed curve representing C_i divides the plane into $k_i(k_i - 3)/2 + 2$ connected cells. Color these cells with black

and while so that no two cells that share a boundary arc have the same color. The curve representing C_2 intersects C_1 exactly $2(k_1 - 2) + (k_2 - 2)k_1 \equiv k_1 k_2 \pmod{2}$, not counting v . Every time C_2 intersects C_1 , it passes from one cell to another whose color is different. Assume that in a small neighborhood of v the initial segment of an edge of C_2 incident to v lies in a white region. Then the initial segment of the other edge of C_2 incident to v lies in a black region if and only if $k_1 k_2$ is odd. \square

A graph consisting of three internally disjoint paths P_i , $i = 1, 2, 3$, between u and v is called a Θ -graph. A drawing of this Θ -graph is said to be a *preserver* if in a small neighborhood of u the initial pieces of the paths P_i follow each other in the same circular order (clockwise or counterclockwise) as the final pieces do around v . Otherwise, the drawing is called a *converter*. Note that, using this terminology, if G is a planar graph drawn in the plane without crossing, then any Θ -subgraph of this drawing is a converter. The proof of the next lemma is very similar to that of the previous one.

Lemma 2.3. *A Θ -subgraph of a thrackle is a converter if and only if at most one of its three paths has odd length.*

Remark. With the exception of Lemma 2.1(i), all statements and proofs in this section remain valid for generalized thrackles.

3. Bipartite Thrackles

Proof of Theorem 1.2. By Kuratowski's theorem, it is sufficient to show that a thrackable bipartite graph G does not contain a subdivision of K_5 or of $K_{3,3}$.

Suppose that G contains a subdivision of K_5 , whose vertices are v_0, \dots, v_4 . Assume without loss of generality that in a thrackle drawing of G the initial pieces of the edges incident to v_0 follow each other in the clockwise order $v_0 v_1, \dots, v_0 v_4$. Then there are two (even) cycles through v_0, v_1, v_3 and v_0, v_2, v_4 that have no vertex in common other than v_0 . The corresponding two curves cross each other in a small neighborhood of v_0 , contradicting Lemma 2.2.

Suppose next that G contains a subdivision of $K_{3,3}$ with vertex classes $\{u_1, u_2, u_3\}$ and $\{v_1, v_2, v_3\}$. Denote this subdivision by K . Assume first that the lengths of all nine paths in K connecting the u_i 's and the v_j 's have the same parity. Deleting from K the point u_3 together with the three paths connecting it to the v_j 's, we obtain a Θ -graph. In view of Lemma 2.3, it is a converter between u_1 and u_2 . Similarly, deleting u_2 (u_1) we obtain a converter between u_1 and u_3 (u_2 and u_3 , respectively). We say that the type of u_i is *clockwise* or *counterclockwise* according to the circular order of the initial segments of the paths $u_i v_1, u_i v_2, u_i v_3$ around u_i . It follows from the definition of a converter that any two u_i 's must have opposite types, which is impossible.

There are two other essentially different cases according to the parities of the nine paths forming K . It turns out that in both cases one can arrive at a contradiction by showing that there is exactly one pair of points among u_1, u_2, u_3 having opposite types. \square

Proof of Theorem 1.4. In view of the remark at the end of the previous section, the above argument also proves that every bipartite graph that can be drawn as a *generalized thrackle* is planar. To establish the theorem, we have to show that the reverse of this statement is also true, i.e., every bipartite planar graph G can be drawn as a generalized thrackle. To see this, consider a crossing-free embedding of G in the plane such that

- (i) $V(G) = V_1 \cup V_2$, where all points of V_1 are mapped into the upper half-plane and all points of V_2 below the line $y = -1$;
- (ii) every edge $e \in E(G)$ connects a vertex of V_1 to a vertex of V_2 , and each piece of e belonging to the strip $-1 \leq y \leq 0$ is a vertical segment.

Now erase the part of the drawing in the strip $-1 \leq y \leq 0$, and replace the part in the upper half-plane by its reflection about the y -axis. Reconnecting the corresponding pairs of points on the lines $y = -1$ and $y = 0$ by straight-line segments, we obtain a drawing of G such that any pair of independent edges meet an odd number of times. This can be turned into a generalized thrackle by slightly modifying the edges in a small neighborhood of their endpoints so as to reverse the circular order of edges around each vertex of G . \square

We could have completed our proof without using Lemma 2.2. The fact that a thrackle contains no subdivision of K_5 can also be deduced from Lemma 2.3 in a slightly more complicated way.

The proof of Theorem 1.2 also yields the following.

Corollary 3.1. *A graph is planar if and only if it has a drawing whose every Θ -subgraph is a converter.*

For a related result, see [T2].

4. Reduction to the Bipartite Case

Every graph can be made bipartite by the removal of fewer than half of its edges. It follows from Euler's polyhedral formula that any bipartite planar graph of n vertices has at most $2n - 4$ edges ($n > 2$). If in addition the graph has no cycles of length four then this bound can be replaced by $\lfloor 3n/2 \rfloor - 3$ ($n > 3$). Thus, Theorem 1.4 and Lemma 2.1(i) immediately imply the following.

Corollary 4.1. *Let $n > 3$. Then*

- (i) *every thrackle of n vertices has at most $3n - 7$ edges;*
- (ii) *every generalized thrackle of n vertices has at most $4n - 9$ edges.*

In the rest of this section we sketch how to reduce the bound in Corollary 4.1(i) roughly by n .

Let G be a thrackle of n vertices, $n > 3$. One can assume that G is not bipartite, otherwise its number of edges cannot exceed $\lfloor 3n/2 \rfloor - 3$. Let C denote a shortest odd

cycle of G with length c . By Lemma 2.1(i) and by the minimality of C , any vertex of G not on C has at most one neighbor belonging to C . Hence, there are at most n edges of G which are not on C incident to vertices of C . It follows from Lemma 2.1(ii) that the graph $G - C$ obtained from G by the removal of all points of C is bipartite. Thus,

$$|E(G)| \leq |E(G - C)| + n \leq \frac{3(n - c)}{2} + n = \frac{5n}{2} - \frac{3c}{2}.$$

One can refine this argument, as follows. The closed curve representing C cuts the plane into a number of cells that can be colored with black and white so that no two cells with a common boundary arc have the same color. Let b and w denote the number of vertices of $G - C$ lying in black and in white cells, respectively. Clearly, $c + b + w = n$, and one can assume without loss of generality that $b \leq w$, so that

$$b \leq \frac{n - c}{2}. \quad (1)$$

Observe that if an edge e connects a point of C to (say) a black vertex, then in a small neighborhood of this point the initial piece of e must be white. There are at most b such edges, and if one removes all of them together with all edges of C , the resulting graph (thackle) becomes bipartite. This with (1) yields the inequality

$$|E(G)| \leq \left\lfloor \frac{3n}{2} \right\rfloor - 3 + b + c \leq 2n + \frac{c}{2} - 3.$$

Comparing the last two inequalities, we obtain that $|E(G)| < (2 + \frac{1}{8})n$.

One can further reduce this bound by utilizing an idea of Conway (see [W1], [G2], [PRS2], and [PRS1]). Now we replace each vertex and edge of C by two nearby vertices and edges, respectively. More precisely, we split each vertex v of C into two vertices, v_b and v_w , and connect all black and white neighbors of v not on C to v_b and v_w , respectively. Furthermore, if v and v' are two consecutive vertices of C , we connect v_b to v'_w and v'_b to v_b . It is not hard to see that this construction can be carried out in such a way that the resulting drawing G' is a thrackle, which becomes bipartite after the removal of all edges between v_b 's and black vertices. Thus,

$$|E(G')| - b = |E(G)| + c - b \leq \left\lfloor \frac{3(n + c)}{2} \right\rfloor - 3,$$

which implies by (1) that

$$|E(G)| \leq 2n - 3,$$

as stated in Theorem 1.1.

5. Small Forbidden Configurations

All of the results in the previous sections were based on parity arguments. Theorem 1.4 shows that if we want to settle Conway's original conjecture, we have to go beyond

these methods. In the proof of Theorem 1.1 we were able to explore a property of thrackles that does not hold for generalized thrackles. Namely, we used the fact that a thrackable graph has no cycle of length four (Lemma 2.1(i)). By excluding some other small configurations that would contradict the thrackle conjecture, one can easily improve the bound in Theorem 1.1. The trouble is that it is quite difficult to find any new nontrivial forbidden subgraph, because even a relatively small graph may have an enormous number of topologically different drawings such that no two edges meet more than once. In this section, we illustrate these difficulties by an example.

Let Θ_3 denote a graph consisting of two vertices connected by three internally disjoint paths of length three.

Theorem 5.1. *A thrackable graph cannot contain Θ_3 as a subgraph.*

For the proof we need some preparation. Let G be a fixed thrackle whose edges are smooth curves. Given two directed edges e and f that do not share an endpoint, we say that e *meets f clockwise* if at their intersection point a tangent vector to e can be carried into a tangent vector of f by a clockwise turn with angle less than π .

Let $P = e_1 e_2 e_3 e_4$ be a directed path in G with length four, directed toward e_4 . Associate P with a 4×4 matrix M such that $M_{ij} = 0$ if $i = j$ or if e_i and e_j do not have an interior point in common. Otherwise, let $M_{ij} = 1$ or -1 depending on whether e_i meets e_j clockwise or counterclockwise. Clearly, M is antisymmetric and it is determined by the triple (M_{13}, M_{14}, M_{24}) . This triple is called the *type* of P . It turns out that there are only six possible types:

$$\begin{array}{lll} a = (1, 1, -1); & b = (1, -1, -1); & c = (1, -1, 1); \\ A = (-1, -1, 1); & B = (-1, 1, 1); & C = (-1, 1, -1). \end{array}$$

Lemma 5.2. *Let e_1, e_2, \dots, e_6 be six directed edges of a thrackle that form a simple directed cycle, and let $P_i = e_i e_{i+1} e_{i+2} e_{i+3}$, where the indices are taken mod 6. Then $\text{type}(P_1) \text{ type}(P_2) \dots \text{type}(P_6)$ must be one of the following sequences: $AaAaAa$, $aAaAaA$, $BbBbBb$, $bBbBbB$.*

Given a directed path $P = e_1 e_2 e_3 e_4$, let the *reverse* of P be defined as $P^{-1} = e_4^{-1} e_3^{-1} e_2^{-1} e_1^{-1}$, where e_i^{-1} denotes the same edge as e_i but with reversed orientation. If $e_1 \dots e_5$ is a simple directed path, we say that $P' = e_2 e_3 e_4 e_5$ can be obtained from $P = e_1 e_2 e_3 e_4$ by a *shift*.

Lemma 5.3. *Let P be a path of length four in a thrackle, and assume that $\text{type}(P) \in \{a, b, A, B\}$.*

- (i) *type(P^{-1}) = b, a, B , or A according to whether $\text{type}(P) = a, b, A$, or B .*
- (ii) *If P' can be obtained from P by a shift and $\text{type}(P') \in \{a, b, A, B\}$, then $\text{type}(P) \text{ type}(P')$ must be one of the following six pairs: aA, aB, bB, Aa, Ab, Bb .*

Proof of Theorem 5.1. Assume that there is a thrackle containing Θ_3 as a subgraph. By Lemma 5.2, the type of every directed path of Θ_3 belongs to the set $\{a, b, A, B\}$. Consider

a path P whose type belongs to $\{b, B\}$. (If P does not satisfy this condition, then its reverse does.) Observe that the topology of Θ_3 allows us to transform P into its reverse by a series of shifts. It follows from Lemma 5.3(ii) that the types of all paths obtained during this process belong to $\{b, B\}$. However, by Lemma 5.3(i), $\text{type}(P^{-1}) \in \{a, A\}$, which is a contradiction. \square

References

- [BF] L. Babai and P. Frankl, Linear Algebra Methods in Combinatorics, Part I, Technical Report, University of Chicago, 1988.
- [B] E. R. Berlekamp, On subsets with intersections of even cardinality, *Canadian Mathematical Bulletin* **12** (1969), 363–366.
- [BE] N. G. de Bruin and P. Erdős, On a combinatorial problem, *Proc. Konink. Nederl. Akad. Wetensch. Ser. A* **51** (1948), 1277–1279 (= *Indagationes Mathematicae* **10** (1948), 421–423).
- [DET] G. Di Battista, P. Eades, R. Tamassia, and I. G. Tollu, Algorithms for drawing graphs: An annotated bibliography, *Computational Geometry: Theory & Applications* **4** (1994), 235–282.
- [FS] W. Fenchel and J. Sutherland, Lösung der Aufgabe 167, *Jahresbericht der Deutschen Mathematiker-Vereinigung* **45** (1935), 33–35.
- [GJ] M. R. Garey and D. S. Johnson, Crossing number is NP -complete, *SIAM Journal of Algebraic Discrete Methods* **4** (1983), 312–316.
- [G1] R. L. Graham, The largest small hexagon, *Journal of Combinatorial Theory* **18** (1975), 165–170.
- [G2] J. Green-Collingsham, Thrackles, Surfaces and Drawings of Graphs, Doctoral Dissertation, Clemson University, 1993.
- [G3] R. K. Guy, Crossing numbers of graphs, In: *Graph Theory and Applications*, Lecture Notes in Mathematics, Vol. 303, Springer-Verlag, Berlin, 1972, pp. 111–124.
- [HH] H. Hopf and E. Panwitz, Aufgabe Nr. 167, *Jahresbericht der Deutschen Mathematiker-Vereinigung* **43** (1934), 114.
- [KST] T. Kovari, V. T. Sós, and P. Turán, On a problem of K. Zarankiewicz, *Colloquium Mathematicum* **3** (1954), 50–57.
- [LST] J. A. de Loera, B. Sturmfels, and R. R. Thomas, Gröbner bases and triangulations of the second hypersimplex, *Combinatorica*, to appear.
- [PA] J. Pach and P. K. Agarwal, *Combinatorial Geometry*, Wiley, New York, 1995.
- [PRS1] B. Piazza, R. Ringelsen, and S. Stueckle, On Conway's reduction of the thrackle conjecture and some associated drawings, to appear.
- [PRS2] B. Piazza, R. Ringelsen, and S. Stueckle, Properties of non-minimum crossings for some classes of graphs, In: *Graph Theory, Combinatorics, and Applications*, Vol. 2 (Y. Alavi et al., eds.), Wiley, New York, 1991, pp. 975–989.
- [RSP] R. Ringelsen, S. Stueckle, and B. Piazza, Subgraphs and bounds on maximum crossings, *Bulletin ICA* **2** (1991), 33–46.
- [T1] P. Turán, A note of welcome, *Journal of Graph Theory* **1** (1977), 7–9.
- [T2] W. T. Tutte, Toward a theory of crossing number, *Journal of Combinatorial Theory* **8** (1970), 45–53.
- [WB] A. T. White and L. W. Beineke, Topological graph theory, In: *Selected Topics in Graph Theory* (L. Beineke and R. Wilson, eds.), Academic Press, London, 1978, pp. 15–49.
- [W1] D. R. Woodall, Thrackles and deadlock, In: *Combinatorial Mathematics and Its Applications* (D. J. A. Welsh, ed.), Academic Press, London, 1969, pp. 335–348.
- [W2] D. R. Woodall, Open problems, In: *Combinatorics, Proceedings of the Conference on Combinatorial Mathematics* (D. J. A. Welsh and D. R. Woodall, eds.), Institute of Mathematics and Its Applications, Southend-on Sea, Essex, 1972, pp. 341–350.

Received July 11, 1995, and in revised form April 16, 1996

An Efficient Algorithm for Euclidean Shortest Paths Among Polygonal Obstacles in the Plane

S. Kapoor,¹ S. N. Maheshwari,¹ and J. S. B. Mitchell²

¹Department of Computer Science and Engineering, Indian Institute of Technology, Hauz Khas, New Delhi, India

{skapoor, snml}@cse.iitdernet.in

²Department of Applied Mathematics and Statistics, State University of New York, Stony Brook, NY 11794–3600, USA

jsbm@ams.sunysb.edu

Abstract. We give an algorithm to compute a (Euclidean) shortest path in a polygon with h holes and a total of n vertices. The algorithm uses $O(n)$ space and requires $O(n + h^2 \log n)$ time.

1. Introduction

Let P denote a (multiply connected, closed) polygon in the plane having h holes ("obstacles") and a total of n vertices. The problem of computing a shortest ("geodesic") path from a point $s \in P$ to a point $t \in P$ has been well studied in computational geometry; see [11], [14], [6], [8], [10], [13], [14], [17], and [21]–[23], as well as the recent survey chapter by Mitchell [15]. In the case that $h = 0$, the shortest path can be computed in $O(n)$ time [6], [12]. In the case that $h > 0$, the complexity of the problem has been worst-case quadratic ($O(n^2)$), until the recent $O(n^{1.5+\epsilon})$ algorithm of Mitchell [14], which develops the "continuous Dijkstra" paradigm, and its improvement by Hershberger and Suri [7], [8], which results in an algorithm whose running time is $O(n \log n)$.

A lower bound of $\Omega(n + h \log h)$ is easy to establish (from sorting or convex hulls), but, to date, there is no matching upper bound. In fact, no upper bounds of the form $O(n + f(h))$, having *linear* dependence on n , have previously been published.

Here, we offer a simple algorithm whose dependence on n is linear, both in time and space. The time dependence on h , however, is slightly worse than quadratic: $O(n + h^2 \log n)$. Thus, while our algorithm is optimal for values of h that are roughly $O(\sqrt{n})$, it is inferior to the best known methods in cases in which h is not relatively small compared