

ON THE BETTI NUMBERS OF REAL VARIETIES

J. MILNOR

The object of this note will be to give an upper bound for the sum of the Betti numbers of a real affine algebraic variety. (*Added in proof.* Similar results have been obtained by R. Thom [10].)

Let V be a variety in the real Cartesian space \mathbb{R}^m , defined by polynomial equations

$$f_1(x_1, \dots, x_m) = 0, \dots, f_p(x_1, \dots, x_m) = 0.$$

The q th Betti number of V will mean the rank of the Čech cohomology group $H^q(V)$, using coefficients in some fixed field F .

THEOREM 2. *If each polynomial f_i has degree $\leq k$, then the sum of the Betti numbers of V is $\leq k(2k-1)^{m-1}$.*

Analogous statements for complex and/or projective varieties will be given at the end.

I wish to thank W. May for suggesting this problem to me.

REMARK A. This is certainly not a best possible estimate. (Compare Remark B.) In the examples which I know, the sum of the Betti numbers of V is always $\leq k^m$. Consider for example the m polynomials

$$f_i(x_1, \dots, x_m) = (x_i - 1)(x_i - 2) \cdots (x_i - k),$$

where $i = 1, 2, \dots, m$. These define a zero-dimensional variety, consisting of precisely k^m points.

The proofs of Theorem 2 and of Theorem 1 (which will be stated later) depend on the following.

LEMMA 1. *Let $V_0 \subset \mathbb{R}^m$ be a zero-dimensional variety defined by polynomial equations $f_1 = 0, \dots, f_m = 0$. Suppose that the gradient vectors df_1, \dots, df_m are linearly independent at each point of V_0 . Then the number of points in V_0 is at most equal to the product $(\deg f_1)(\deg f_2) \cdots (\deg f_m)$.*

PROOF. Approximate f_1, \dots, f_m by real polynomials F_1, \dots, F_m of the same degrees whose coefficients are algebraically independent. Now consider the variety $V_{\mathbb{C}}$ in the complex Cartesian space \mathbb{C}^m defined by the equations $F_1 = 0, \dots, F_m = 0$. It follows from van der Waerden [9, §41] that the number of points in $V_{\mathbb{C}}$ is equal to $(\deg f_1)(\deg f_2) \cdots (\deg f_m)$. Since each point of V_0 lies close to some real point of $V_{\mathbb{C}}$; this proves Lemma 1.

Received by the editors January 17, 1963.

Next let W be a compact, nonsingular hypersurface in \mathbb{R}^m , defined by the equation $f=0$, where f is a polynomial of degree $2k$.

THEOREM 1. *The sum of the Betti numbers of W is $\leq 2k(2k-1)^{m-1}$*

Similar estimates have been given by Olneĭnik [5].

The proof follows. Consider the function $\mathbf{n}: W \rightarrow S^{m-1}$ which assigns to each point of W its unit normal vector $\text{grad } f / |\text{grad } f|$. According to a theorem of Sard, the set of critical values of \mathbf{n} has measure zero in S^{m-1} . (See [7] or [6, p. 10].) Hence, after rotating the coordinate system if necessary, we may assume that the two points $(0, \dots, 0, \pm 1)$ of S^{m-1} are not critical values of \mathbf{n} . In terms of local coordinates u_1, \dots, u_{m-1} on W , this means that the matrix $(\partial \mathbf{n}_i / \partial u_j)_{i,j < m}$ is nonsingular whenever $\mathbf{n} = (0, \dots, 0, \pm 1)$.

Now note that the "height function" $(x_1, \dots, x_m) \rightarrow x_m$ on W has no degenerate critical points. In fact near each critical point of this function we can choose local coordinates u_1, \dots, u_{m-1} so that

$$x_1 = u_1, \dots, u_{m-1} = u_{m-1}, \quad x_m = h(u_1, \dots, u_{m-1}),$$

where h is the height function. Then

$$\mathbf{n}(u_1, \dots, u_{m-1}) = \pm (\partial h / \partial u_1, \dots, \partial h / \partial u_{m-1}, -1) / \sqrt{(\sum (\partial h / \partial u_j)^2 + 1)};$$

hence

$$\partial \mathbf{n}_i / \partial u_j = \pm \partial^2 h / \partial u_i \partial u_j$$

at each critical point. Therefore the matrix $(\partial^2 h / \partial u_i \partial u_j)$ is nonsingular; as was asserted.

Now applying Morse theory to the height function $h: W \rightarrow \mathbb{R}$, it follows that the sum of the Betti numbers of W is less than or equal to the number of critical points of h . (See for example Seifert and Threlfall [8, §§5 and 8].)

The critical points of h can be characterized as the solutions of the m polynomial equations

$$\partial f / \partial x_1 = 0, \dots, \partial f / \partial x_{m-1} = 0, \quad f = 0.$$

Note that at each critical point of h the m gradient vectors

$$d(\partial f / \partial x_1), \dots, d(\partial f / \partial x_{m-1}), df$$

are linearly independent.

PROOF. If we differentiate the identity

$$f(u_1, \dots, u_{m-1}, h(u_1, \dots, u_{m-1})) = 0$$

twice, and then eval

$$\partial^2 f / \partial x_i \partial x_j$$

which shows that this the linear inde

Now Lemma 1 as than or equal to the

$$(\text{deg } \partial f / \partial x_i)$$

Therefore the sum this number. This

REMARK B. UHF Morse theory direct slightly sharper in

Using this, Theore

PROOF OF THEOREM For each r we will

Given numbers ϵ for which

$$f_1^2 +$$

This is a compact If $r \leq \delta / \epsilon$ note that which the equality

holds will be denou is an algebraic hyp $\{\text{deg } f_p\} \geq 1$. This lar value of the fixed, it follows singular for almost and δ are chosen s

Applying The $\leq 2k(2k-1)^{m-1}$. B

1964]

twice, and then evaluate at a critical point, we obtain

$$\partial^2 f / \partial x_i \partial x_j + (\partial f / \partial x_m)(\partial^2 h / \partial u_i \partial u_j) = 0;$$

which shows that the matrix $(\partial^2 f / \partial x_i \partial x_j)_{i,j < m}$ is nonsingular. From this the linear independence of the m gradient vectors follows easily.

Now Lemma 1 asserts that the number of critical points of h is less than or equal to the product

$$(\deg \partial f / \partial x_1) \cdots (\deg \partial f / \partial x_{m-1})(\deg f) = (2k - 1)^{m-1} 2k.$$

Therefore the sum of the Betti numbers of W is less than or equal to this number. This proves Theorem 1.

REMARK B. Using a more complicated argument, which applies Morse theory directly to the function f , one can prove the following slightly sharper inequality:

$$\text{rank } H^*W \leq (2k - 1)^m + 1.$$

Using this, Theorem 2 can be sharpened to the form

$$\text{rank } H^*V \leq \frac{1}{2}((2k - 1)^m + 1).$$

PROOF OF THEOREM 2. Let rD^m denote the disk of radius r in R^m . For each r we will show that $H^*(V \cap rD^m)$ has rank $\leq k(2k - 1)^{m-1}$.

Given numbers $\epsilon, \delta > 0$ let $K(\epsilon, \delta)$ denote the set of points in R^m for which

$$f_1^2 + \cdots + f_p^2 + \epsilon^2(x_1^2 + \cdots + x_m^2) \leq \delta^2.$$

This is a compact set, since it is contained in the disk of radius δ/ϵ . If $r \leq \delta/\epsilon$ note that $V \cap rD^m \subset K(\epsilon, \delta)$. The set of points in R^m for which the equality

$$f_1^2 + \cdots + f_p^2 + \epsilon^2|x|^2 = \delta^2$$

holds will be denoted by $\partial K(\epsilon, \delta)$, and called the boundary of K . This is an algebraic hypersurface of degree $2k$, where $k = \text{Max}\{\deg f_1, \dots, \deg f_p\} \geq 1$. This hypersurface is nonsingular if and only if δ^2 is a regular value of the function $f_1^2 + \cdots + f_p^2 + \epsilon^2|x|^2$. Hence, keeping ϵ fixed, it follows from Sard's theorem that this hypersurface is nonsingular for almost all values of δ . We will assume henceforth that ϵ and δ are chosen so that this is the case.

Applying Theorem 1 to ∂K we see that $H^*(\partial K)$ has rank $\leq 2k(2k - 1)^{m-1}$. But the Alexander duality theorem implies that

$$\text{rank } H^*(K) = \frac{1}{2} \text{rank } H^*(\partial K).$$

(See for example [3, p. 279].) Hence

$$\text{rank } H^*K \leq k(2k - 1)^{m-1}.$$

Now choose sequences $\{\epsilon_i\}$ and $\{\delta_i\}$ so that $\{\epsilon_i\}$ decreases monotonely with limit zero; so that $\{\delta_i/\epsilon_i\}$ decreases monotonely with limit r ; and so that each $\partial K(\epsilon_i, \delta_i)$ is nonsingular. Then

$$K(\epsilon_1, \delta_1) \supset K(\epsilon_2, \delta_2) \supset K(\epsilon_3, \delta_3) \supset \dots$$

with intersection $V \cap rD^m$. (These inclusion relations become clear if we put the defining inequality for $K(\epsilon, \delta)$ in the form

$$\sum f_i^2/\delta^2 + |x|^2 \epsilon^2/\delta^2 \leq 1,$$

and note that $\delta_{i+1} \leq \delta_i \epsilon_{i+1}/\epsilon_i \leq \delta_i$.) Therefore the Čech cohomology group $H^*(V \cap rD^m)$ is equal to the direct limit of the groups $H^*K(\epsilon_i, \delta_i)$. (Compare Eilenberg and Steenrod [1, p. 261].) Hence

$$\text{rank } H^*(V \cap rD^m) \leq \limsup(\text{rank } H^*K(\epsilon_i, \delta_i)) \leq k(2k - 1)^{m-1}.$$

There remains one small but tricky point: the passage to the limit as $r \rightarrow \infty$. One approach is to make use of the theorem that V can be triangulated. (See for example Lefschetz and Whitehead [2].) It follows that the Betti numbers of V based on singular homology theory are the same as those based on Čech cohomology theory. But the singular homology group $H_*(V)$ is equal to the direct limit of the singular homology groups of the compact subsets¹ of V . If $P \subset V \cap rD^m \subset V$ is any compact sub-polyhedron, then the restriction homomorphism $H^*V \rightarrow H^*P$ has rank $\leq k(2k - 1)^{m-1}$; hence the corresponding homology homomorphism $H_*P \rightarrow H_*V$ also has rank $\leq k(2k - 1)^{m-1}$. Therefore the limit H_*V has rank $\leq k(2k - 1)^{m-1}$; which completes the proof.

An alternative approach which does not assume triangulability can be sketched as follows. By an argument similar to that in [4, p. 338] one constructs an exact sequence

$$0 \rightarrow \mathcal{L}'H^{q-1}(V \cap rD^m) \rightarrow H^qV \rightarrow \mathcal{L}H^q(V \cap rD^m) \rightarrow 0,$$

where \mathcal{L} denotes the inverse limit functor, and \mathcal{L}' denotes its first derived functor. Since each $H^{q-1}(V \cap rD^m)$ is a finite-dimensional

¹ In contrast note that the Čech homology group of a space need not be the direct limit of the Čech homology groups of its compact subsets. A counter-example is provided by the union of the curve $y = \sin(1/x)/x$ and the y -axis.

for space, an argument
Steenrod in [1, p. 226] that

$$\text{rank } H^*V = \text{rank}$$

which completes the alter
as corollaries one ob
the, for complex or pro

COROLLARY 1 (THE COM
polynomial equations of

COROLLARY 2 (THE COM
ned by homogeneous
 $\text{rank } H^*V_2 \leq mk(2k - 1)^2$

COROLLARY 3 (THE RE
ned by homogeneous poly
 $\leq mk(2k - 1)^m$. Here the
only in the special case

In order to prove Corol
a real variety in R^{2m} . To p
ogeneous equations can be
The intersection $V_2 \cap S^{2m-1}$
is. Using the Gysin sequ
estimate. Corollary 3 is pr
is a 2-fold covering of V_2

Added in proof. A real
siders a locus which is de
any estimate must depend
but also upon their num
ities

$$(x - 1)^2 \geq \frac{1}{10},$$

on the real line define a

THEOREM 3. If $X \subset R^n$
form

with total degree $d = \deg f$

rank H^*

vector space, an argument similar to that used by Eilenberg and Steenrod in [1, p. 226] shows that the \mathcal{L}' term is zero. Therefore

$$\text{rank } H^*V = \text{rank } (\mathcal{L}H^*(V \cap rD^m)) \leq k(2k - 1)^{m-1},$$

which completes the alternative proof.

As corollaries one obtains similar estimates, presumably rather crude, for complex or projective varieties.

COROLLARY 1 (THE COMPLEX AFFINE CASE). *If $V_1 \subset \mathbf{C}^m$ is defined by polynomial equations of degree $\leq k$, then $\text{rank } H^*V_1 \leq k(2k - 1)^{2m-1}$.*

COROLLARY 2 (THE COMPLEX PROJECTIVE CASE). *If $V_2 \subset \mathbf{P}_m(\mathbf{C})$ is defined by homogeneous polynomial equations of degree $\leq k$, then $\text{rank } H^*V_2 \leq mk(2k - 1)^{2m+1}$.*

COROLLARY 3 (THE REAL PROJECTIVE CASE). *If $V_3 \subset \mathbf{P}_m(\mathbf{R})$ is defined by homogeneous polynomial equations of degree $\leq k$, then $\text{rank } H^*V_3 \leq mk(2k - 1)^m$. Here the initial factor m in this expression is needed only in the special case when the coefficient field F has characteristic 2.*

In order to prove Corollary 1 it is only necessary to think of V_1 as a real variety in \mathbf{R}^{2m} . To prove Corollary 2 note that the same homogeneous equations can be used to define an affine variety $V'_2 \subset \mathbf{C}^{m+1}$. The intersection $V'_2 \cap S^{2m+1}$ is the total space of a circle bundle over V_2 . Using the Gysin sequence of this bundle, one arrives at the given estimate. Corollary 3 is proved in the same way. In this case $V'_3 \cap S^m$ is a 2-fold covering of V_3 ; so that the prime 2 plays a special role.

Added in proof. A rather different problem arises when one considers a locus which is defined by polynomial inequalities. In this case any estimate must depend not only on the degrees of the inequalities but also upon their number. As an example the p quadratic inequalities

$$(x - 1)^2 \geq \frac{1}{10}, \quad (x - 2)^2 \geq \frac{1}{10}, \quad \dots, \quad (x - p)^2 \geq \frac{1}{10}$$

on the real line define a locus with $p + 1$ components.

THEOREM 3. *If $X \subset \mathbf{R}^n$ is defined by polynomial inequalities of the form*

$$f_1 \geq 0, \dots, f_p \geq 0$$

with total degree $d = \text{deg } f_1 + \dots + \text{deg } f_p$, then

$$\text{rank } H^*X \leq \frac{1}{2}(2 + d)(1 + d)^{m-1}.$$

PROOF. Again we first replace X by $X \cap rD^m$. This can be done simply adjoining the additional inequality

$$f_0 = r^2 - |x|^2 \geq 0.$$

Given numbers $\epsilon \geq \epsilon^{p+1} \geq \delta > 0$, let $L(\epsilon, \delta)$ denote the set of points for which

$$f_0 + \epsilon \geq 0, \dots, f_p + \epsilon \geq 0$$

and

$$(f_0 + \epsilon)(f_1 + \epsilon) \cdots (f_p + \epsilon) - \delta \geq 0.$$

This set is compact. Its boundary is obtained by setting only the last expression equal to zero.

Given ϵ we may assume that δ is chosen so that the boundary is nonsingular. Therefore the proof (if not the exact statement) of Theorem 1 shows that

$$\text{rank } H^*(\partial L(\epsilon, \delta)) \leq (2+d)(1+d)^{m-1};$$

and hence that

$$\text{rank } H^*L(\epsilon, \delta) \leq \frac{1}{2}(2+d)(1+d)^{m-1}.$$

Now choose numbers ϵ_i tending monotonely to zero and choose $0 < \delta_i < (\epsilon_i - \epsilon_{i+1})^{p+1}$; so that $L(\epsilon_1, \delta_1) \supset L(\epsilon_2, \delta_2) \supset \dots$ with intersection $X \cap rD^m$. As in the proof of Theorem 2, it follows that

$$\text{rank } H^*X \leq \limsup (\text{rank } H^*(X \cap rD^m)) \leq \frac{1}{2}(2+d)(1+d)^{m-1}.$$

REFERENCES

1. S. Eilenberg and N. Steenrod, *Foundations of algebraic topology*, Princeton Univ. Press, Princeton, N. J., 1952.
2. S. Lefschetz and J. H. C. Whitehead, *On analytical complexes*, Trans. Amer. Math. Soc. **35** (1933), 510-517.
3. J. Milnor, *On the immersion of n -manifolds in $(n+1)$ -space*, Comment. Math. Helv. **30** (1956), 275-284.
4. ———, *On axiomatic homology theory*, Pacific J. Math. **12** (1962), 337-341.
5. O. A. Oleĭnik, *Estimates of the Betti numbers of real algebraic hypersurfaces*, Mat. Sb. (N.S.) **28** (70) (1951), 635-640. (Russian)
6. G. de Rham, *Variétés différentiables*, Hermann, Paris, 1955.
7. A. Sard, *The measure of the critical values of differentiable maps*, Bull. Amer. Math. Soc. **48** (1942), 883-897.
8. H. Seifert and W. Threlfall, *Variationsrechnung im Grossen*, Teubner, Berlin, 1938.
9. B. van der Waerden, *Einführung in die algebraische Geometrie*, Springer, Berlin, 1939.
10. R. Thom, *Sur l'homologie des variétés algébriques réelles*, (to appear).

PRINCETON UNIVERSITY

A TOTALLY NONARCHIVAL SPACE

This paper gives a new example of a connected, Hausdorff space which assure the existence of continuous paths.

Example 1 gives a compact, Hausdorff space in which each subset of cardinality \aleph_1 will be used in describing the space.

EXAMPLE 1. Let I be the interval $[0, 1]$. Partition I into \aleph_1 subintervals in lexicographical order. Let $f(x) = \frac{1}{2}$, then either $f(x) = \frac{1}{2}$ or $f(x) < \frac{1}{2}$.

EXAMPLE 2. Let S be the set of four points in I^2 . Let S consist of four points in I^2 . Figure 1, with the corresponding diagram.

Let K' be a copy of K . Let k_x denote the corresponding copy of k , for $x < z$, be the union of K and k_x . Let \bar{k}_x be the closure of $\{k_x\}$.

Let H be the decomposition of I^2 into elements of which are \bar{k}_x for $x \leq z$.

Figure 2 will aid in understanding the conditions that prevent continuous paths from representing copies of S .

H is a totally nonarchival space.

Presented to the Society of Mathematical Geometers in 1962 and, in revised form, to the American Mathematical Society in 1963.

¹ This work was completed while the author was at the University of California, Berkeley, as an N.S.F. Science Fellow.

² A continuum T is said to be *totally nonarchival* if for each point x in T , the set $T - \{x\}$ contains x in its interior.

³ The author wishes to thank the referee for his attention.

⁴ $g < f$ if, for some t , $g(t) < f(t)$.