

On Bounding the Betti Numbers and Computing the Euler Characteristic of Semi-algebraic Sets

Saugata Basu*

Abstract

In this paper we give a new bound on the sum of the Betti numbers of semi-algebraic sets. This extends a well-known bound due to Oleinik and Petrovsky [19], Thom [23] and Milnor [18]. In separate papers they proved that the sum of the Betti numbers of a semi-algebraic set $S \subset \mathbb{R}^k$, defined by $P_1 \geq 0, \dots, P_s \geq 0, \deg(P_i) \leq d, 1 \leq i \leq s$, is bounded by $(O(sd))^k$. Given a semi-algebraic set $S \subset \mathbb{R}^k$ defined as the intersection of a real variety, $Q = 0, \deg(Q) \leq d$, whose real dimension is k' , with a set defined by a quantifier-free Boolean formula with atoms of the form, $P_i = 0, P_i > 0, P_i < 0, \deg(P_i) \leq d, 1 \leq i \leq s$, we prove that the sum of the Betti numbers of S is bounded by $s^{k'}(O(d))^k$. In the special case, when S is defined by $Q = 0, P_1 > 0, \dots, P_s > 0$, we have a slightly tighter bound of $\binom{s}{k'}(O(d))^k$. This result generalises the Oleinik-Petrovsky-Thom-Milnor bound in two directions. Firstly, our bound applies to arbitrary semi-algebraic sets, not just for basic semi-algebraic sets. Secondly, the combinatorial part (the part depending on s) in our bound, depends on the dimension of the variety rather than that of the ambient space. It also generalizes a result in [7] where a similar bound is proven for the number of connected components. In the second part of the paper we use the tools developed for the above results, as well as some additional techniques, to give the first single exponential time algorithm for computing the Euler characteristic of arbitrary semi-algebraic sets.

1 Introduction

Let $\mathcal{P} = \{P_1, \dots, P_k\} \subset \mathbb{R}[X_1, \dots, X_k]$, be a family of polynomials whose degrees are bounded by d , and let S be a semi-algebraic set defined by a quantifier-free Boolean formula, with atoms of the form $P_i \{>, <, =\} 0, 1 \leq i \leq s$. The Betti numbers, $\beta_i(S)$, which are the ranks of the singular homology groups of S , are a measure of the topological complexity of S and can be bounded in terms of s, d and k . Since by Collin's algorithm for cylindrical algebraic decomposition, there exists a cellular decomposition of S into $(sd)^{2^{O(k)}}$ cells, the same bound applies to $\beta_i(S)$. In the case where the set S is a basic closed semi-algebraic set defined by, $P_1 \geq 0, \dots, P_s \geq 0$, with $\deg(P_i) \leq d$, there is a tighter bound of $(O(sd))^k$ on the sum of the Betti numbers of S . This was proved in separate papers by Oleinik and Petrovsky [19], Thom [23] and Milnor [18]. (Note that this includes the case when S is the real zeros of a set of polynomials.) This bound plays an important role in algorithmic real algebraic geometry [17], and has been used recently in proving lower bounds in the algebraic computation tree model (see [24]).

We extend the same bound to any arbitrary semi-algebraic set. More precisely, we prove the following theorem:

Theorem 1 *Let $S \subset \mathbb{R}^k$, be the intersection of a semi-algebraic set defined by a quantifier-free Boolean formula involving a family, $\mathcal{P} = \{P_1, \dots, P_s\}$, of s polynomials, with the zero set $Z(Q)$, of a polynomial Q . Let the geometric dimension of $Z(Q)$ be k' , and the degrees of the polynomials in $\mathcal{P} \cup \{Q\}$ be bounded by d . Then, the sum of the Betti numbers of S is bounded by $s^{k'}(O(d))^k$. In the special*

*Courant Institute of Mathematical Sciences, New York University, New York, NY 10012, U.S.A.. Supported in part by NSF grant CCR-9122103

case, when the formula defining S is $Q = 0, P_1 > 0, \dots, P_s > 0$, we have a slightly tighter bound of $\binom{s}{k'}(O(d))^k = \binom{s}{k'}^{k'}(O(d))^k$.

If Q is taken as the zero polynomial, then $Z(Q) = R^k$ and $k' = k$, and we get the Oleinik-Petrovsky-Thom-Milnor bound for arbitrary semi-algebraic sets as a special case. Note that a bound of $\binom{s}{k'}(O(d))^k$ on the zero-th Betti number for arbitrary semi-algebraic sets (which is just the number of connected components) was known before [6] and our bound is a generalization of this result to the higher Betti numbers.

Note also that a lower bound of $\binom{sd}{k'}^{k'}$ on the zero-th Betti number is easily obtained by considering the set of non-zeros of s polynomials, each of them a product of d linear polynomials, restricted to a k' dimensional linear subspace.

The dependence on k' instead of k in the combinatorial part (the part depending on s) of the bound in theorem 1 becomes important when we consider low dimensional semi-algebraic sets embedded in a higher dimensional space, and this is sometimes important in applications. For example, the bound on the number of connected components in [6] plays a crucial role in the proof of the main result in [12], where the variety is the real Grassmannian $G_{m,n}$ (the space of m dimensional subspaces of R^n), embedded as a $m(n - m)$ dimensional variety in R^{n^2} .

Remark: In the special case when S is compact, a bound of $(O(sd))^{2k}$ can be deduced from Theorem 1 in Yao's paper [24].

In order to achieve the bound in theorem 1 we prove that an arbitrary semi-algebraic set is homotopic to a compact semi-algebraic set defined by polynomials in general position. This result generalizes a similar result in Canny [8] where it was proved for semi-algebraic sets defined by a single sign condition, and this intermediate result might be of independent interest.

In the second part of the paper, we consider the problem of computing the Euler characteristic $\chi(S)$, of a semi-algebraic set S . The Euler characteristic, which is the alternating sum of the Betti numbers of S , is an important topological invariant and thus can be used as a test to rule out topological equivalence.

We prove the following theorem.

Theorem 2 *Let $S \subset R^k$ be a real semi-algebraic set, defined by a quantifier-free Boolean formula with atoms of the form, $P_i \{<, >, =\} 0, 1 \leq i \leq s$, where $\mathcal{P} = \{P_1, \dots, P_s\} \subset Z[X_1, \dots, X_k]$, is a family of s polynomials whose degrees are bounded by d , and the bit lengths of the coefficients of P_i are bounded by L . Then, there exists an algorithm for computing $\chi(S)$ which performs at most $(ksd)^{O(k)} L^{O(1)}$ bit operations.*

We remark that computing stratifications of semi-algebraic sets, and thus computing their homology groups, in single exponential time, is a central open problem of computational real algebraic geometry (see the survey by Chazelle [9]). Single exponential algorithms for determining certain other (weaker) topological properties of semi-algebraic sets are known. For example, it is possible to compute the number of connected components [8, 13, 14, 15], semi-algebraic description of the connected components [16], as well as to decide whether two points are in the same connected component of a semi-algebraic set [8, 13, 15], in single exponential time.

Collin's algorithm for computing a cylindrical algebraic decomposition [10], gives sufficient topological information for computing the Euler characteristic, and in fact the homology groups of a given semi-algebraic set [22]. However, this algorithm has double exponential complexity $(sd)^{2^{O(k)}}$. Previously, this was the best algorithm for computing the Euler characteristic of general semi-algebraic sets. A single exponential algorithm for computing the Euler characteristic of a smooth algebraic hypersurface is mentioned in [21].

The rest of the paper is organized as follows. In section 2 we give a brief discussion of our use of infinitesimals and the model of computation with appropriate pointers to literature. In section 3 we show how to perturb the polynomials to bring them into general position without changing

the homotopy type of the given semi-algebraic set. In section 4 we prove our bound on the Betti numbers of general semi-algebraic sets. In section 5 we describe our algorithm for computing the Euler characteristic. In subsection 5.1 we describe certain real algebraic algorithms which we use as subroutines. In subsection 5.2 we give an algorithm for computing the Euler characteristic of a basic semi-algebraic set, and in subsection 5.3 we give an algorithm for computing the Euler characteristic of an arbitrary semi-algebraic set.

We omit the proofs of most of the lemmas and propositions in the main text. These can be found in the appendix.

2 Preliminaries

In this paper we give algorithms for computing the Euler characteristic of real semi-algebraic sets, defined by polynomials with integer coefficients. However, our algorithm will work just as well if the coefficients came from some other ordered domain D , and the set defined in some real closed field containing D . The complexity of our algorithm is the number of arithmetic operations, as well as comparisons, performed by the algorithms in D . When $D = \mathbb{Z}$, we also express the complexity in terms of bit operations (see the remarks in section 6.2.2 in the appendix).

Our argument depends on perturbing the polynomials by various *infinitesimals* and to work over the field of Puiseux series in these infinitesimals. We write $R\langle\epsilon\rangle$ for the real closed field of Puiseux series in ϵ with coefficients in R [2]. The sign of an element in this field agrees with the sign of the coefficient of the lowest degree term in ϵ . This order makes ϵ positive and smaller than any positive element of R . We also iterate this notation in the usual way so that $R\langle\epsilon_1, \epsilon_2\rangle = R\langle\epsilon_1\rangle\langle\epsilon_2\rangle$ and thus, $1 \gg \epsilon_1 \gg \epsilon_2$ i.e. ϵ_2 is positive and smaller than any positive element in $R\langle\epsilon_1\rangle$ and ϵ_1 is smaller than any positive element of R . The map $eval_\epsilon : V \rightarrow R$, maps an element of $R\langle\epsilon\rangle$ which is bounded over R , i.e. one that has no negative powers of ϵ , to its constant term.

We also note that the asymptotic complexities of all the real algebraic algorithms that we use, do not change if we introduce a constant number of these infinitesimals [5].

3 Going to general position

Following [5], we say that a family of polynomials \mathcal{P} in k variables is in *general position* if no $k+1$ of them have a common real zero. In this section we show that given a semi-algebraic set, S , defined by a family, \mathcal{P} , of s polynomials with degrees bounded by d , we can define a new compact semi-algebraic set S' , which is homotopy equivalent to S , but which is defined by a family, \mathcal{P}' , of polynomials in general position. Moreover, $|\mathcal{P}'| \leq 6s+1$, and the degrees of the polynomials in \mathcal{P}' are bounded by d' , where d' is the least even number greater than d . Similar results appear in [8], where they are proved for semi-algebraic sets defined by one single sign condition. Our proof techniques are similar to those used in [8], but our results apply to arbitrary semi-algebraic sets.

Given a family of polynomials $\mathcal{P} = \{P_1, \dots, P_s\}$, a *sign condition* σ of \mathcal{P} is an element of $(+, 0, -)^s$. A sign condition σ is *non-empty* if there exists a point $x \in R^k$ such that,

$$\sigma = (\text{sign}(P_1(x)), \dots, \text{sign}(P_s(x))).$$

Given a sign condition σ of the family \mathcal{P} we will denote by $\sigma(\mathcal{P})$ the formula, $\bigwedge_i P_i s_i 0$, where s_i is $>, =, <$ according to whether $\sigma(i)$ is $+, 0, -$, respectively.

We first ensure that a given semi-algebraic set is homotopic to a bounded one.

Lemma 1 *Let S be any semi-algebraic set. Then, for Ω large enough, $S' = S \cap (X_1^2 + \dots + X_k^2 \leq \Omega)$, has the same homotopy type as S .*

Proof: See appendix. □

Let S be a semi-algebraic set defined by the formula, $\bigvee_{1 \leq j \leq L} \sigma_j(\mathcal{P})$, where each σ_j is a non-empty sign condition for the family \mathcal{P} .

For any sign condition σ , and a new variable ϵ we will denote by $\sigma(\mathcal{P}, \epsilon)$ the following formula: For every strict sign in σ we retain the corresponding conjunct in $\sigma(\mathcal{P}, \epsilon)$. For every equality in σ we replace the corresponding conjunct, $P_j = 0$, by the conjunct, $(P_j > -\epsilon) \wedge (P_j < \epsilon)$.

Let $S(\epsilon)$ denote the semi-algebraic set defined by the formula $\bigvee_{1 \leq j \leq L} \sigma_j(\mathcal{P}, \epsilon)$.

The next lemma shows that for sufficiently small $\epsilon > 0$, S and $S(\epsilon)$ has the same homotopy type.

Lemma 2 *Let S and $S(\epsilon)$ be as above. Then, for sufficiently small $\epsilon > 0$, S is a deformation retract of $S(\epsilon)$.*

Proof: See appendix. □

Note that, the set $S(\epsilon)$ is defined by a disjunction of conjunctions of strict inequalities involving $3s$ polynomials, $\bigcup_{P \in \mathcal{P}} \{P, P + \epsilon, P - \epsilon\}$, and that there no equalities in the formula.

We next prove that given a semi-algebraic set defined by a disjunct of conjuncts of the form, $\bigwedge_j Q_j \leq 0$ where $s_j \in \{<, >\}$, then it is possible to define a new semi-algebraic set defined by polynomials in general position, which is of the same homotopy type.

Let S be a semi-algebraic set, defined by the formula, $\bigvee_{1 \leq j \leq L} \sigma_j(Q_j)$, where each σ_j is a sign-condition on a subset Q_j of a family of polynomial Q , and σ_j does not contain equality.

Let $Q = \{Q_1, \dots, Q_s\}$, and $\deg(Q_i) < d$, $1 \leq i \leq s$.

For $1 \leq i \leq 2s$ let, $H_i = 1 + \sum_{1 \leq j \leq k} i^j X_j^{2d}$.

For, $Q_i \in Q$, let $Q_i^+ = (1 - \delta)Q_i + \delta H_{2i-1}$, and $Q_i^- = (1 - \delta)Q_i + \delta H_{2i}$, where δ is a new variable. Let Q' be the family $\bigcup_i \{Q_i^+, Q_i^-\}$.

Lemma 3 *For sufficiently small $\delta > 0$, the family of polynomials Q' is in general position.*

Proof: See [5]. □

For a sign condition σ on Q , without any equalities, define $\sigma(Q, \delta)$ to be the formula obtained as follows:

Every conjunct $Q_i > 0$ is replaced by $Q_i^- \geq 0$, while every conjunct $Q_i < 0$ is replaced by $Q_i^+ \leq 0$.

Let $S^-(\delta)$ be the set defined by, $\bigvee_{1 \leq j \leq L} \sigma_j(Q_j, \delta)$.

Lemma 4 *Let S and $S^-(\delta)$ be as above. Then, for sufficiently small $\delta > 0$, $S^-(\delta)$ is a deformation retract of S .*

Proof: See appendix. □

Combining the previous four lemmas we obtain the following proposition.

Proposition 1 *Let S be any semi-algebraic set defined by a family of s polynomials with degrees bounded by d . Then, there exists a set $S(\Omega, \epsilon, \delta)$ defined by a family of at most $6s + 1$ polynomials, in $X_1, \dots, X_k, \epsilon, \delta, \Omega$ with degrees bounded by $2d$ such that for sufficiently small $\epsilon > 0$, $\frac{1}{\Omega} > 0$, $\delta > 0$, this new family of polynomials is in general position, and $S(\Omega, \epsilon, \delta)$ has the same homotopy type as S . Moreover, the set $S(\Omega, \epsilon, \delta)$ is compact, and defined as a disjunction of conjunctions of non-strict inequalities.*

Proof: Follows easily from the four previous lemmas. □

In our proofs we will need a slightly stronger notion of general position.

Let Q be a polynomial, such that $Z(Q)$ has geometric dimension k' . We call a family of polynomials \mathcal{P} to be in *general position with respect to Q* , if no $k' + 1$ of the polynomials in \mathcal{P} has a real zero in common with Q .

Moreover, consider the intersection of a semi-algebraic set defined by a family of polynomials \mathcal{P} with $Z(Q)$. Using the same arguments as in the proof of proposition 1, always carrying along the extra condition $Q = 0$, and replacing the polynomials H_i by different infinitesimals (see [6] for details), we can replace $S \cap Z(Q)$ by a compact semi-algebraic set contained in $Z(Q)$, having the same homotopy type, but defined by at most $6s + 1$ polynomials in general position with respect to Q .

We also remark that if a family of polynomials \mathcal{P} is in general position with respect to a polynomial Q , then for an infinitesimal ϵ the family $\cup_{P \in \mathcal{P}} \{P, P + \epsilon, P - \epsilon\}$ is also in general position with respect to Q as long as we consider zeros that are bounded over R (see [5] for details). This is implicit in the proofs presented below.

4 Bound on the Betti numbers of a semi-algebraic set

We make use of the following facts from algebraic topology which follows easily from the Mayer-Vietoris sequence (see [1]). Given a semi-algebraic set S we define $r(S) = \sum_i \beta_i(S)$. Let S_1 and S_2 be two compact semi-algebraic sets. Then, from the Mayer-Vietoris sequence,

$$\cdots \rightarrow H_{i+1}(S_1 \cup S_2) \rightarrow H_i(S_1 \cap S_2) \rightarrow H_i(S_1) \oplus H_i(S_2) \rightarrow H_i(S_1 \cup S_2) \rightarrow H_{i-1}(S_1 \cap S_2) \rightarrow \cdots,$$

we can deduce the following:

$$r(S_1) + r(S_2) \leq r(S_1 \cup S_2) + r(S_1 \cap S_2), \quad (1)$$

and,

$$r(S_1 \cup S_2) \leq r(S_1) + r(S_2) + r(S_1 \cap S_2). \quad (2)$$

The following lemma follows easily from inequalities (1) and (2), and the proof of lemma 4.

Lemma 5 *Let S be a semi-algebraic set defined by a conjunct $(Q = 0) \wedge (\sigma_1(\mathcal{P}) \vee \cdots \vee \sigma_L(\mathcal{P}))$, where Q is a polynomial, and σ_j , $1 \leq j \leq L$, are sign conditions on a family of polynomials \mathcal{P} , such that none of the σ_j contain an equality. Then, $\beta_i(S) = \sum_{1 \leq j \leq L} \beta_i(S_j)$ and $\chi(S) = \sum_{1 \leq j \leq L} \chi(S_j)$, where S_j is the set defined by the conjunct $(Q = 0) \wedge \sigma_j(\mathcal{P})$.*

Proof: See appendix. □

4.1 Proof of Theorem 1

Proof: In view of proposition 1 and the remarks following it we can assume without loss of generality that S is a compact set defined by s polynomials in general position with respect to Q .

We next prove two lemmas that will imply the theorem.

Given a polynomial Q and a family of polynomials $\mathcal{P} = \{P_1, \dots, P_s\}$, we define the *combinatorial level* of the system (Q, \mathcal{P}) to be the least integer m such that no $m + 1$ of the polynomials in \mathcal{P} have a common real zero with Q . We refer to the set of real zeros of a polynomial Q by $Z(Q)$.

For example, the combinatorial level of (Q, \mathcal{P}) is bounded by k' if the dimension of $Z(Q)$ is k' and the polynomials in \mathcal{P} are in general position with respect to Q .

Lemma 6 *Let S be a semi-algebraic set, defined by $Q = 0, P_1 > 0, \dots, P_s > 0$, where $Z(Q)$ is bounded and Q is non-negative everywhere. Let $\mathcal{P} = \{P_1, \dots, P_s\}$, and the combinatorial level of the system (Q, \mathcal{P}) be bounded by $m \leq k$, and the degrees of the polynomials Q and P_i be bounded by $2d$ and d respectively. Then, $r(S) = \binom{s}{m} (O(d))^k$.*

Proof: Consider the set $S(\epsilon)$ defined by $Q = 0, P_1 \geq \epsilon, \dots, P_s \geq \epsilon$, where $\epsilon > 0$ is a new variable. Then, it is easy to prove that for sufficiently small $\epsilon > 0$ $S(\epsilon)$ is a deformation retract of S . Let T denote the set defined by, $Q = 0, P_2 \geq \epsilon, \dots, P_s \geq \epsilon$.

Consider the sets $U_1 = T \cap ((P_1 \geq \epsilon) \cup (P_1 \leq -\epsilon))$, $V_1 = T \cap (-\epsilon \leq P_1 \leq \epsilon)$ and $W_1 = T \cap ((P_1 = \epsilon) \cup (P_1 = -\epsilon))$.

From inequality 1 we see that, $r(U_1) + r(V_1) \leq r(T) + r(W_1)$ and from lemma 5 that $r(S) \leq r(U_1)$. It immediately follows that, $r(S) \leq r(T) + r(W_1)$.

Moreover, let $r(s, d, m, k)$ be the maximum possible value of $r(S)$ for any set defined by a system with these parameters. Then, we have the recurrence,

$$r(s, d, m, k) \leq r(s-1, d, m, k) + 2r(s-1, d, m-1, k), m \leq k,$$

$$r(0, d, m, k) = (O(d))^k, r(s, d, 0, k) = (O(d))^k.$$

It follows easily that, $r(s, d, m, k) = \binom{s}{m} (O(d))^k$. □

Lemma 7 *Let S be a compact semi-algebraic set contained in the zero set of a polynomial Q and defined by a family of polynomials $\mathcal{P} = \{P_1, \dots, P_s\}$. Suppose that Q is non-negative everywhere, the combinatorial level of the system (Q, \mathcal{P}) is bounded by $m \leq k$, and the degrees of the polynomials Q and P_i are bounded by $2d$ and d respectively. Then, $r(S) = s^m (O(d))^k$.*

Proof: Let S be defined by the conjunct $(Q = 0) \wedge (\sigma_1 \vee \dots \vee \sigma_L)$ where the σ_i are sign conditions on the family \mathcal{P} .

Let $\epsilon_1 \gg \epsilon_2 \gg \dots \gg \epsilon_s$, be infinitesimals. Consider the sets, T_1, U_1, V_1, W'_1, W_1 defined as follows:

$$T_1 = S \cap ((P_1 \geq \epsilon_1) \cup (P_1 \leq -\epsilon_1)),$$

$$U_1 = S \cap (P_1 = \epsilon_1),$$

$$V_1 = S \cap (P_1 = -\epsilon_1),$$

$$W'_1 = S \cap (-\epsilon_1 \leq P_1 \leq \epsilon_1),$$

and

$$W_1 = S \cap (P_1 = 0).$$

Now, $S = T_1 \cup W'_1$, and it is clear that $T_1 \cap W'_1 = U_1 \cup V_1$, and $U_1 \cap V_1 = \emptyset$.

Using the inequality (2) twice along with the fact that $r(\emptyset) = 0$, we have,

$$r(S) \leq r(T_1) + r(W'_1) + r(U_1) + r(V_1).$$

Moreover, using the same arguments as in lemmas 2 and 4, one can show that W'_1 has the same homotopy type as W_1 . We omit the proof as the arguments are completely analogous.

Thus, we have,

$$r(S) \leq r(T_1) + r(W_1) + r(U_1) + r(V_1).$$

Note that, U_1, V_1, W_1 are defined by the systems, $(Q + (P_1 - \epsilon_1)^2, \mathcal{P})$, $(Q + (P_1 + \epsilon_1)^2, \mathcal{P})$, $(Q + P_1^2, \mathcal{P} \setminus \{P_1\})$, (note that Q is non-negative everywhere) respectively. Moreover, each of the above system has combinatorial level at most $m-1$.

We next consider T_1 which is defined by a set of sign conditions without the atom $P_1 = 0$, and eliminate the atom $P_2 = 0$.

We repeat the process described earlier, replacing $P_2 < 0, P_2 > 0$ by $P_2 \leq -\epsilon_2, P_2 \geq \epsilon_2$ respectively. In this way we obtain the inequality,

$$r(T_1) \leq r(T_2) + r(W_2) + r(U_2) + r(V_2),$$

U_2, V_2, W_2 are sets defined by systems with combinatorial level at most $m - 1$.

The remaining set, T_2 is homotopic to the union of the sets defined by those sign conditions appearing in the definition of S , which contain neither $P_1 = 0$ nor $P_2 = 0$.

We continue this process till we have eliminated $P_s = 0$, and we get the inequality,

$$r(S) \leq r(T_s) + \sum_{1 \leq i \leq s} (r(U_i) + r(V_i) + r(W_i)),$$

The sets U_i, V_i, W_i are defined by systems of at most $2s$ polynomials having combinatorial level at most $m - 1$. Moreover, the remaining term $r(T_s)$ is the bound on the Betti numbers of a semi-algebraic set defined by a union of sign conditions of the form, $Q = 0, P_i \lessgtr 0$ for $1 \leq i \leq s$, with $s_i \in \{<, >\}$.

Again, by lemma 5, the Betti numbers of this set are the sum of the the Betti numbers of the non-empty sets defined by each individual sign condition. Now, consider the set T defined by $Q = 0, P_1^2 > 0, \dots, P_s^2 > 0$. From the above remark it is clear that $r(T_s) \leq r(T)$. Moreover, applying the bound proved in lemma 6 we have, $r(T) = \binom{s}{m} (O(d))^k$.

Let, $r(s, d, k, m)$ denote the maximum Betti number of a semi-algebraic set defined by a system (Q, \mathcal{P}) with $|\mathcal{P}| = s$, combinatorial level of the sytem bounded by m , and the degrees of the polynomial Q and those in \mathcal{P} bounded by d . Then, we have the recurrence,

$$r(s, d, k, m) = \binom{s}{m} (O(d))^k + 3s r(2s, d, k, m - 1), \quad m \leq k$$

$$r(s, d, k, 0) = (O(d))^k.$$

This recurrence solves to $r(s, d, k, m) = s^m (O(d))^k, m \leq k$, which proves the lemma. \square

The theorem now follows since the combinatorial level of a system (Q, \mathcal{P}) with the family \mathcal{P} in general position with respect to Q is bounded by k' . \square

5 Algorithm to compute the Euler Characteristic of a General Semi-Algebraic Set

By proposition 1 we can assume without loss of generality, that the given semi-algebraic set is compact. If the given set is not compact then we make the perturbations described in section 3 and compute the Euler characteristic of the perturbed set. The Euler characteristic of this new set is equal to the Euler characteristic of the original set. The new system will have at most $6s + 1$ polynomials with degrees at most $2d$. Moreover, we now have to compute in a larger ring $Z[\delta, \epsilon, \Omega]$. However, since we have introduced only three infinitesimals, the asymptotic complexity of the algorithm is not affected (see [5]).

5.1 Algorithmic Preliminaries

In our algorithm we will utilize several other algorithms from real algebraic geometry as subroutines. In this section we list the algorithms that we use as subroutines with appropriate pointers to the literature. We use a subroutine [5] that constructs univariate representations of the zeros of a zero dimensional variety. This subroutine takes as input a Gröbner basis of a zero-dimensional ideal, I , of polynomials in k variables and outputs a set consisting of $(k + 2)$ -tuples of univariate polynomials, (f, g_0, \dots, g_k) such that the complex zeroes of I are among the points obtained by evaluating the rational functions $(\frac{g_1}{g_0}, \dots, \frac{g_k}{g_0})$ at the roots of the univariate polynomial f , for all the tuples (f, g_0, \dots, g_k) in the output. We say that the real points corresponding to the tuple (f, g_0, \dots, g_k) , are *associated* to the tuple, and the tuple itself is a *univariate representation* of these points. Moreover, if the degrees of the

polynomials in the input are bounded by d , the degrees of the polynomials in the output as well as the complexity of this subroutine is bounded by $d^{O(k)}$.

We also make use of an algorithm, called the sample points subroutine, that computes at least one points in every connected component of every non-empty sign condition (referred to as *cells* henceforth) of a family of polynomials \mathcal{P} , of size s and degrees bounded by d . The subroutine also outputs the sign vector of the polynomials of \mathcal{P} at each output point. The complexity of this subroutine is $\binom{O(s)}{k} s d^{O(k)} = (s/k)^k s d^{O(k)}$.

Lastly, we make use a multivariate sign determination subroutine [5]. The input is a system T of polynomial equations in k variables, with a finite number of zeros, along with a Gröbner basis for the ideal generated by the polynomials in T , and a set of s polynomials $\mathcal{P} = \{P_1, \dots, P_s\}$. The output is the list of non empty sign-conditions $\sigma_1, \dots, \sigma_M$ of \mathcal{P} at the real zeros of the system T and the numbers c_1, \dots, c_M , where c_i is the number of real zeros of T at which the sign vector of \mathcal{P} is σ_i . Moreover, if the polynomials in the input have degrees bounded by d , the complexity of this subroutine is $d^{O(k)}$.

We will also use this subroutine in the special case of computing the index of a symmetric square matrix of size $k \times k$, with polynomial entries, at the real zeros of a zero-dimensional system. Again, if the degrees of the polynomials in the input are bounded by d , the complexity of the subroutine is bounded by $(kd)^{O(k)}$, (see [21]).

5.2 The algorithm for a semi-algebraic set defined by one sign condition

In this section, we describe an algorithm for computing the Euler characteristic of a semi-algebraic set defined by one single sign condition on a family of polynomials. Using lemma 4 we can assume, without loss of generality that the semi-algebraic set S is defined by, $P_1 \geq 0, \dots, P_s \geq 0$.

The following proposition is crucial for our algorithms. Let S be a compact, basic, semi-algebraic set defined by, $P_1 \geq 0, \dots, P_s \geq 0$. Let the degrees of the polynomials, P_i be bounded by d . The next proposition proves that S is homotopy equivalent to a certain set which is bounded by a smooth hypersurface.

Proposition 2 *Let $Q = \prod_{1 \leq i \leq s} (\zeta + (1 - \zeta)P_i) + \zeta^{s+1}(X_1^{2d'} + \dots + X_k^{2d'} + 1)$, where ζ is a new variable, and $2d' > \sum_{1 \leq i \leq s} \deg(P_i)$. Then, for all sufficiently small $\zeta > 0$, the set defined by $(Q \geq 0) \wedge_{1 \leq i \leq s} ((1 - \zeta)P_i + \zeta \geq 0)$, is homotopy equivalent to the set S .*

Moreover, the above set is bounded by a smooth hypersurface $Q = 0$, which has a finite number of critical points for the projection map onto the X_1 co-ordinate and these critical points are non-degenerate and have distinct X_1 co-ordinates.

Proof: See appendix. □

From proposition 2 it follows that S is homotopic to the union of those connected components of the set defined by $Q \geq 0$, where the polynomials $(\zeta + (1 - \zeta)P_i)$ are non-negative, where $Q = \prod_{1 \leq i \leq s} (\zeta + (1 - \zeta)P_i) + \zeta^{s+1}(X_1^{2d'} + \dots + X_k^{2d'} + 1)$, and ζ is an infinitesimal.

We actually compute the Euler characteristic of the latter set which is equal to $\chi(S)$.

5.2.1 Description of the algorithm

Given a basic semi-algebraic set S defined by, $P_1 \geq 0, \dots, P_s \geq 0$, where $\deg(P_i) < d, 1 \leq i \leq s$, the algorithm computes $\chi(S)$. We then introduce a positive infinitesimal ζ and construct the polynomial, $Q = \prod_{1 \leq i \leq s} (\zeta + (1 - \zeta)P_i) + \zeta^{s+1}(X_1^{2d'} + \dots + X_k^{2d'} + 1)$.

We then solve the zero-dimensional system, $Q = \frac{\partial Q}{\partial X_1} = \dots = \frac{\partial Q}{\partial X_k} = 0$, using the univariate representation subroutine and for each real zero p of this system we check whether the polynomials $\zeta + (1 - \zeta)P_1, \dots, \zeta + (1 - \zeta)P_s$ are all non-negative at the point p . We retain only those real solutions for which this is satisfied. For each critical point $p = (p_1, \dots, p_k)$ retained in the previous step, check

whether the $(p_1 - \epsilon, p_2, \dots, p_k)$ belong to the set defined by $(Q \geq 0) \cap_{1 \leq i \leq s} (\zeta + (1 - \zeta)P_i \geq 0)$, where ϵ is a new infinitesimal. Note that, since the point p is given only as a univariate representation the last sentence is slightly misleading. However, it is clear that using the univariate representation of p and the sign determination subroutine, we can perform this check. Retain only those critical points p for which the above is false.

For each critical point p retained so far, compute the index of the critical point, which is the index of the Hessian matrix, $(\frac{\partial^2 Q}{\partial X_i \partial X_j})_{ij}$, $2 \leq i, j \leq k$, evaluated at p . For $0 \leq i \leq k$ let c_i denote the number of critical points of index i . We output $\chi(S) = \sum_{0 \leq i \leq k} (-1)^i c_i$.

5.2.2 Proof of Correctness

Before giving the proof we need to recall certain facts from Morse theory of stratified sets [11].

A Whitney stratification of a space X is a decomposition of X into submanifolds called strata, which satisfy certain frontier conditions, (see [11] page 37). In particular, given a compact set bounded by a smooth hypersurface, the boundary and the interior form a Whitney stratification.

Now, let X be a compact Whitney stratified subset of R^k , and f a restriction to X of a smooth function. A critical point of f is defined to be a critical point of the restriction of f to any stratum, and a critical value of f is the value of f at a critical point.

The first fundamental result of stratified Morse theory is the following.

Theorem 3 (*Goresky-MacPherson, [11]*) *As c varies in the open interval between two adjacent critical values, the topological type of $X \cap f^{-1}((-\infty, c])$ remains constant.*

Stratified Morse theory actually gives a recipe for describing the topological change in $X \cap f^{-1}((-\infty, c])$ as c crosses a critical value. This is given in terms of Morse data, which consists of a pair of topological spaces (A, B) , $A \supset B$, with the property that as c crosses the critical value $v = f(p)$, the change in $X \cap f^{-1}((-\infty, c])$ can be described by gluing in A along B .

A function is called a Morse function if it has only non-degenerate critical points when restricted to each strata, and all its critical values are distinct. (There is an additional non-degeneracy condition which states that the differential of f at a critical point p of a strata S should not annihilate any limit of tangent spaces to a stratum other than S . However, in our simple situation this will never apply.)

In stratified Morse theory the Morse data is presented as a product of two pairs, called the tangential Morse data and the normal Morse data. The notion of product of pairs is the standard one in topology, namely $(A, B) \times (A', B') = (A \times A', A \times B' \cup B \times A')$.

Instead of describing these in full generality, we restrict ourselves to the very simple situation in which we need them, namely when $X \subset R^k$ is compact and is bounded by a smooth hypersurface. The stratification we consider is the stratification of X into its boundary and interior and the function f is just the projection map onto the X_1 co-ordinate. In this case, all critical points occur on the boundary strata (the interior is open in R^k).

The tangential Morse data at a critical point p is then given by $(D^\lambda \times D^{k-\lambda}, (\partial D^\lambda) \times D^{k-\lambda})$ where D^k is the closed k dimensional disk, ∂ is the boundary map, and λ is the index of the Hessian matrix of f at p .

Let $p = (p_1, \dots, p_k)$. The normal slice $N(p)$ at p is defined to be a sufficiently small segment of length 2ϵ given by, $N(p) = \{(t, p_2, \dots, p_k) \mid -\epsilon \leq t \leq \epsilon\}$. Let $L(p) = \{(-\epsilon, p_2, \dots, p_k)\} \cap X$. Note that, $L(p)$ is empty if X does not lie immediately to the left of p , and a single point otherwise. The normal Morse data at p is then $(N(p), L(p))$.

The interested reader can refer to figure 1 in subsection 6.1 of the appendix for an illustration in the case of a solid torus in R^3 .

The second fundamental theorem of stratified Morse theory states the following:

Theorem 4 (Goresky-MacPherson, [11]) *Let f be a Morse function on a compact Whitney stratified space X . Then, Morse data measuring the change in the topological type of $X \cap f^{-1}((-\infty, c])$ as c crosses the critical value v of the critical point p is the product of the normal Morse data at p and the tangential Morse data at p .*

We are now in a position to prove the main proposition of this section.

Proposition 3 *The above algorithm correctly computes the Euler characteristic of S .*

Proof: By proposition 2 $\chi(S)$ is equal to the Euler characteristic of the set defined by $Q \geq 0$ and $\zeta + (1 - \zeta)P_i \geq 0$, $1 \leq i \leq s$. Denote this set S' .

Moreover, S' is compact, bounded by a smooth hypersurface, and the projection map π onto the X_1 co-ordinate is a Morse function.

By Theorem 3, the topological type of $S' \cap \pi^{-1}((-\infty, c])$ remains constant as c varies in the open interval between two adjacent critical values.

Let, c_1, \dots, c_m be the critical values of $Q = 0$ which lie in S' . And let, S'_i denote the set $S' \cap \pi^{-1}((-\infty, c_i])$, and let $S'_0 = \emptyset$. Let, (A_i, B_i) be the Morse data at the critical value c_i . Then for $1 \leq i \leq m$,

$$\chi(S'_i) = \chi(S'_{i-1}) + \chi(A_i, B_i).$$

This follows from a standard application of the excision property of homology theory.

Let $(A_i, B_i) = (C_i, D_i) \times (M_i, N_i)$, where (C_i, D_i) is the tangential Morse data, and (M_i, N_i) the normal Morse data. Then it is easy to see that $\chi(A_i, B_i) = \chi(C_i, D_i)\chi(M_i, N_i)$.

We next observe that, for a critical point (p_1, \dots, p_k) , $\chi(M_i, N_i) = 0$ if $(p_1 - \epsilon, p_2, \dots, p_k) \in S'$. In this case, N_i , consists of a single point, as S' intersect a sufficiently small segment parallel to the X_1 axis passing through the critical point. Moreover, M_i is the closed segment, and thus, $\chi(M_i, N_i) = \chi(M_i) - \chi(N_i) = 1 - 1 = 0$.

If the test in Step 3 returns false, then $N_i = \emptyset$, and in this case, $\chi(M_i, N_i) = \chi(M_i) = 1$.

Finally, $(C_i, D_i) = (D^\lambda \times D^{k-\lambda}, (\partial D^\lambda) \times D^{k-\lambda})$, where λ is the index of the critical point computed in Step 4. Thus, $\chi(C_i, D_i) = \chi(D^\lambda, \partial D^\lambda) = \chi(D^\lambda) - \chi(\partial D^\lambda)$.

Now, $\chi(D^\lambda) = 1$, and $\chi(\partial D^\lambda) = 1 - (-1)^\lambda$ and hence $\chi(C_i, D_i) = 1 - (1 - (-1)^\lambda) = (-1)^\lambda$.

Since $\chi(S') = \chi(S'_m)$ it is clear that the algorithm correctly computes $\chi(S')$. \square

5.2.3 Complexity of the Algorithm

The polynomial Q has degree bounded by $O(sd)$. The cost of computing the indices of the Hessian matrices in the last step dominates the cost. Using the bound mentioned in section 5.1 for the subroutine used to compute these indices, the total complexity of this algorithm is bounded by $(ksd)^{O(k)}$.

5.3 The case of a general semi-algebraic set

5.3.1 Description of the Algorithm

Using the algorithm in [5] list all non-empty sign conditions, $\sigma_1, \dots, \sigma_m$ of the family \mathcal{P} , such that the set S_j defined by σ_j is contained in S for $1 \leq j \leq m$.

For each j , $1 \leq j \leq m$, do the following. Without loss of generality assume that σ_j is of the form, $P_1 = \dots = P_l = 0, P_{l+1} > 0, \dots, P_s > 0$. Let U_j be the set defined by, $P_1 = \dots = P_l = 0, P_{l+1} \geq 0, \dots, P_s \geq 0$, and let $V_j = U_j \cap Z(\Pi_{l < i \leq s} P_i)$.

Using the algorithm described in section 5.2 for computing the Euler characteristic of basic semi-algebraic sets, compute $\chi(U_j)$ and $\chi(V_j)$.

Output, $\chi(S) = \sum_{1 \leq j \leq m} (\chi(U_j) - \chi(V_j))$.

5.3.2 Proof of Correctness and Complexity

The complexity of this algorithm is again bounded by $(ksd)^{O(k)}$. See subsections 6.2.1 and 6.2.2 of the appendix for the proof of correctness and the complexity analysis.

References

- [1] E. ARTIN, H. BRAUN *Introduction to Algebraic Topology*, Charles E. Merrill Publishing, Columbus, Ohio, (1969).
- [2] J. BOCHNAK, M. COSTE, M.-F. ROY *Géométrie algébrique réelle*. Springer-Verlag (1987).
- [3] M. BEN-OR *Lower Bounds for Algebraic Computation Trees*, *Proc. 15th STOC*, 80-86, (1983).
- [4] S. BASU, R. POLLACK, M.-F. ROY *A New Algorithm to Find a Point in Every Cell Defined by a Family of Polynomials* in “Quantifier Elimination and Cylindrical Algebraic Decomposition”, B. Caviness and J. Johnson Eds., Springer Verlag, to appear.
- [5] S. BASU, R. POLLACK, M.-F. ROY *On the combinatorial and algebraic complexity of Quantifier Elimination*, *Proc. 35th IEEE Symp. on Foundations of Computer Science*, 632-642, (1994) (full version to appear in the JACM).
- [6] S. BASU, R. POLLACK, M.-F. ROY. Computing points meeting every cell on a variety. In *The First Workshop on the Algorithmic Foundations of Robotics*, Randy Wilson and Jean-Claude Latombe Eds., A. K. Peters, Boston, MA, (1994).
- [7] S. BASU, R. POLLACK, M.-F. ROY *On the Number of Cells Defined by a Family of Polynomials on a Variety*, to appear in *Mathematika*.
- [8] J. CANNY, *Computing road maps in general semi-algebraic sets*, the *Computer Journal*, 36: 504-514, (1993).
- [9] B. CHAZELLE *Computational Geometry: A Retrospective*, *Proc. 26th STOC*, 75-94, (1994).
- [10] G. E. COLLINS, *Quantifier elimination for real closed fields by cylindrical algebraic decomposition* Springer Lecture Notes in Computer Science 33, 515-532.
- [11] M. GORESKEY, R. MACPHERSON *Stratified Morse Theory*, Springer-Verlag, (1987).
- [12] J.E. GOODMAN, R. POLLACK, R. WENGER, *Bounding the Number of Geometric Permutations Induced by k -transversals*, *Proc. 10th Annual Symposium on Computational Geometry*, (1994) (journal version to appear in the *Journal of Combinatorial Theory, Series A*).
- [13] D. GRIGOR'EV, N. VOROBOV *Counting connected components of a semi-algebraic set in subexponential time*. *Comput. Complexity*, 2 :133- 186 (1992).
- [14] L. GOURNAY, J. J. RISLER *Construction of roadmaps of semi-algebraic sets*. Technical Report, DMI, Ecole Normale Supérieure, Paris, France (1992).
- [15] J. HEINTZ, M.-F. ROY, P. SOLERNÒ *Single exponential path finding in semi-algebraic sets II : The general case*. *Algebraic geometry and its applications*. C.L. Bajaj editor, Springer Verlag (1993) 467-481.
- [16] J. HEINTZ, M.-F. ROY, P. SOLERNÒ *Description of the Connected Components of a Semi-algebraic Set in Single Exponential Time*, *Discrete and Computational Geometry* 11:121-140, (1994).

- [17] J. HEINTZ, T. RECIO, M.-F. ROY *Algorithms in Real Algebraic Geometry and Applications to Computational Geometry Discrete and Computational Geometry: Selected Papers from the DIMACS Special Year*, Ed. J.E. Goodman, R. Pollack and W. Steiger, AMS and ACM, 6:137-164, (1991).
- [18] J. MILNOR *On the Betti numbers of real varieties*, Proc. AMS 15, 275-280, (1964).
- [19] O. A. OLEINIK, I. B. PETROVSKII *On the topology of real algebraic surfaces*, Izv. Akad. Nauk SSSR 13, 389-402, (1949).
- [20] R. POLLACK, M.-F. ROY *On the number of cells defined by a set of polynomials*, C. R. Acad. Sci. Paris, 316:573-577, (1993).
- [21] P. PEDERSEN, M.-F. ROY, A. SZPIRGLAS *Counting real zeroes in the multivariate case*. Computational algebraic geometry, Eyssette et Galligo ed. Progress in Mathematics 109, 203-224, Birkhauser (1993).
- [22] J. SCHWARTZ, M. SHARIR *On the 'piano movers' problem II. General techniques for computing topological properties of real algebraic manifolds*. Advances in Applied Mathematics 4, 298-351.
- [23] R. THOM *Sur l'homologie des varietes algebriques reelles*, Differential and Combinatorial Topology, Ed. S.S. Cairns, Princeton Univ. Press, 255-265, (1965).
- [24] A. C. YAO *Decision Tree Complexity and Betti Numbers*, *Proceedings of the 26th STOC* 615-624, (1994).

6 Appendix

Proposition 2 *Let*

$$Q = \prod_{1 \leq i \leq s} (\zeta + (1 - \zeta)P_i) + \zeta^{s+1}(X_1^{2d'} + \cdots + X_k^{2d'} + 1),$$

where ζ is a new variable, and $2d' > \sum_{1 \leq i \leq s} \deg(P_i)$. Then, for all sufficiently small $\zeta > 0$, the set defined by $(Q \geq 0) \wedge_{1 \leq i \leq s} ((1 - \zeta)P_i + \zeta \geq 0)$, is homotopy equivalent to the set S .

Moreover, the above set is bounded by a smooth hypersurface $Q = 0$, which has a finite number of critical points for the projection map onto the X_1 co-ordinate and these critical points are non-degenerate and have distinct X_1 co-ordinates.

Proof: Let $S(\zeta) = (Q \geq 0) \cap_{1 \leq i \leq s} ((1 - \zeta)P_i + \zeta \geq 0)$.

We prove that for sufficiently small $\zeta > 0$, S is a deformation retract of $S(\zeta)$ which will prove the first part of the lemma.

Since S is compact there exists a constant R such that, $x \in S \Rightarrow x_1^{2d'} + \cdots + x_k^{2d'} + 1 < R$. For $0 < \zeta < 1/R$, any point $x = (x_1, \dots, x_k)$ satisfying, $P_1(x) \geq 0, \dots, P_s(x) \geq 0$, will also satisfy $Q \geq 0$. This follows directly from the definition of Q and the fact that $x_1^{2d'} + \cdots + x_k^{2d'} + 1 < R$.

Thus, for every connected component C of S there exists a connected component C' of $Q \geq 0$ such that $C \subset C'$. Moreover, the signs of the polynomials, $\zeta + (1 - \zeta)P_i$, $1 \leq i \leq s$ cannot change over C' , because if one of them became zero Q will be negative at that point. But, since C' contains C , and ζ is sufficiently small, it is clear that $\zeta + (1 - \zeta)P_i > 0$, $1 \leq i \leq s$ over C' . Thus, for ζ small enough, $S \subset S(\zeta)$.

Replacing ζ by a new variable t , consider the set $D \subset R^{k+1}$ defined using the same inequalities as $S(\zeta)$ with ζ replaced by t . Let $\pi : R^{k+1} \rightarrow R$, and $\pi_x : R^{k+1} \rightarrow R^k$, denote the projections onto the t and the X co-ordinates, respectively. Then, from the definition of D , it is clear that $\pi_x(D \cap \pi^{-1}(0)) = S$.

Now for sufficiently small $\zeta > 0$ $D \cap \pi^{-1}(0)$ is a deformation retract of $D \cap \pi^{-1}([0, \zeta])$, and there exists a retraction, (see [8] for details) $\phi : D \cap \pi^{-1}([0, \zeta]) \times [0, \zeta] \rightarrow D \cap \pi^{-1}([0, \zeta])$, such that $\phi(D \cap \pi^{-1}([0, \zeta]), a) = D \cap \pi^{-1}([0, a])$, $a \in [0, \zeta]$.

Thus, we have the map, $\psi : S(\zeta) \times [0, \zeta] \rightarrow S(\zeta)$, defined by, $\psi(x, a) = \pi_x(\phi((x, \zeta), a))$, $x \in S(\zeta)$, $a \in [0, \zeta]$, gives a deformation retract of $S(\zeta)$ to S .

We next show that the set $S(\zeta)$ is bounded by connected components of the smooth hypersurface defined by $Q = 0$.

First observe that the set $Q \geq 0$ is bounded. This follows from the fact that $2d' > sd$ and thus the second term in Q dominates the first as $|x|$ becomes large.

Secondly, the polynomials $\zeta + (1 - \zeta)P_i$ are all strictly positive over $S(\zeta)$. Hence, $S(\zeta)$ must be bounded by the smooth hypersurface $Q = 0$.

It remains to show that the hypersurface $Q = 0$ is smooth and has a finite number of critical points for the projection map onto the X_1 co-ordinate, and that these critical points are non-degenerate with distinct X_1 co-ordinate.

Let $Q_t = \prod_{1 \leq i \leq s} (t + (1 - t)P_i) + t^{s+1}(X_1^{2d'} + \cdots + X_k^{2d'} + 1)$, then for $t = 1$, it is clear that $Q_t = 0$ defines a smooth hypersurface, with all its critical points degenerate, and having distinct X_1 co-ordinates. Moreover, these being stable conditions there is an open interval containing 1, such that for t in this interval, these conditions are satisfied. Now, by quantifier elimination over algebraically closed fields, we have that the set of real t for which the above conditions are not met is either a finite set of points or a complement of a finite set of points. It cannot be latter, since it does not contain an open interval around 1, and hence it must be a finite set of points. Thus, there is an open interval $(0, t_0)$ such that for $\zeta \in (0, t_0)$ the conditions of the proposition are satisfied. \square

Lemma 1 *Let S be any semi-algebraic set. Then, for Ω large enough, $S' = S \cap (X_1^2 + \cdots + X_k^2 \leq \Omega)$, has the same homotopy type as S .*

Proof: We consider the set $S(t) \subset R^{k+1}$ defined by the same formulas as S and the inequality, $t(X_1^2 + \dots + X_k^2) \leq 1$. Let π and π_x denote the projections onto the t and the X co-ordinates respectively.

Then, for sufficiently large Ω , there exists a retraction (see [8] for details), of $S'(t) \cap \pi^{-1}((0, \frac{1}{\Omega})) \rightarrow S'(t) \cap \pi^{-1}(\frac{1}{\Omega})$.

Now, $\pi_x(S'(t) \cap \pi^{-1}((0, \frac{1}{\Omega}))) = S$, and $\pi_x(S'(t) \cap \pi^{-1}(\frac{1}{\Omega})) = S'$.

We define a map $d : S \rightarrow S'(t) \cap \pi^{-1}((0, \frac{1}{\Omega}))$ by $d(x) = (x_1, \dots, x_k, \min(x_1^2 + \dots + x_k^2, \Omega))$. Then, d composed with the retraction mentioned above and π_x gives a retraction of S' to S . \square

Lemma 2 *Let S and $S(\epsilon)$ be as in above. Then, for sufficiently small $\epsilon > 0$, S is a deformation retract of $S(\epsilon)$.*

Proof: Let $S'(\epsilon)$ denote the semi-algebraic set defined similarly as $S(\epsilon)$ with the difference that, for every equality in σ we replace the corresponding conjunct, $P_j = 0$, by the conjunct, $P_j \geq -\epsilon \wedge P_j \leq \epsilon$.

Let, $S(t)$ and $S'(t)$ be subsets of R^{k+1} defined by the corresponding formulas for $S(\epsilon)$ and $S'(\epsilon)$ with ϵ replaced by a new variable t .

Then it is clear that,

$$S \subset S(\epsilon) \subset S'(\epsilon).$$

Moreover, for sufficiently small $\epsilon > 0$, there exists a retraction $S'(\epsilon) \rightarrow S$, given by a map,

$$\phi : S'(\epsilon) \times [0, \epsilon] \rightarrow S'(\epsilon),$$

satisfying, $\phi(S'(\epsilon), t) = S'(t)$, $t \in [0, \epsilon]$. Moreover, $S(\epsilon) \supset S'(t)$, $0 < t < \epsilon$.

Now, consider the restriction of ϕ to $S(\epsilon) \times [0, \epsilon]$. It is clear that defines a proper retraction of $S(\epsilon)$ to S . \square

Lemma 4 *Let S and $S^-(\delta)$ be as above. Then, for sufficiently small $\delta > 0$, $S^-(\delta)$ is a deformation retract of S .*

Proof: Let $D \subset R^{k+1}$ denote the set defined by the same formula as $S^-(\delta)$ with δ replaced by a new variable t .

Let π and π_x denote the projections onto the t and the X co-ordinates respectively.

Then for sufficiently small $\delta > 0$ there exists a retraction ϕ of $D \cap \pi^{-1}((0, \delta]) \rightarrow D \cap \pi^{-1}(\delta)$.

Moreover, it is clear that $\pi_x(D \cap \pi^{-1}((0, \delta])) = S$, and $\pi_x(D \cap \pi^{-1}(\delta)) = S^-(\delta)$.

We now define a one-one continuous map, $d : S \rightarrow D \cap \pi^{-1}((0, \delta])$, that composed with the retraction ϕ and π_x will give a retraction of $S \rightarrow S^-(\delta)$.

The details are messy, but the basic idea is that since S is defined by an open condition, if a point $x \in S$ then $x \in S^-(t)$, for all small enough $t > 0$. We map x to (x, t_0) where t_0 is either the maximum t for which $x \in S^-(t)$, or δ if this maximum is greater than δ .

For, $1 \leq j \leq L$, define $r_j : S \rightarrow R$, as follows: Without loss of generality let Q_1, \dots, Q_{l_j} be the polynomials appearing in σ_j , For $1 \leq i \leq l_j$, let $c_{ij} = 1$, if $\sigma_j(i) = -$, and $c_{ij} = 0$, else.

Define,

$$r_j(x) = \min_{1 \leq i \leq l_j} (-1)^{c_{ij}} \frac{Q_i}{H_{2i-c_{ij}}}.$$

Since the polynomials $H_i \geq 1$, it is clear that r_j is a continuous, well defined real valued function. Define, $r(x) = \min(\max_{1 \leq j \leq L} r_j(x), \delta)$. For $x \in S$ it is clear that, $r(x) > 0$, and moreover $r(x)$ is continuous.

Now, define $d(x) = (x, r(x))$. It is clear that d defines a one-one continuous map from S to $D \cap \pi^{-1}((0, \delta])$ and the lemma follows. \square

Lemma 5 Let S be a semi-algebraic set defined by a conjunct $(Q = 0) \wedge (\sigma_1(\mathcal{P}) \vee \dots \vee \sigma_L(\mathcal{P}))$, where Q is a polynomial, and σ_j , $1 \leq j \leq L$, are sign conditions on a family of polynomials \mathcal{P} , such that none of the σ_j contain an equality. Then, $\beta_i(S) = \sum_{1 \leq j \leq L} \beta_i(S_j)$ and $\chi(S) = \sum_{1 \leq j \leq L} \chi(S_j)$, where S_j is the set defined by the conjunct $(Q = 0) \wedge \sigma_j(\mathcal{P})$.

Proof: For a sign condition σ_j on \mathcal{P} , define $\sigma_j(\epsilon, \mathcal{P})$, to be the formula obtained by replacing every inequality in $\sigma_j(\mathcal{P})$ of the type $P_i > 0$ by $P_i \geq \epsilon$, and $P_i < 0$ by $P_i \leq -\epsilon$. Let S' be the set defined by the conjunct, $(Q = 0) \wedge (\sigma_1(\epsilon, \mathcal{P}) \vee \dots \vee \sigma_L(\epsilon, \mathcal{P}))$, and S'_j to be the set defined by, $(Q = 0) \wedge \sigma_j(\epsilon, \mathcal{P})$ for $1 \leq j \leq L$.

Then using the same argument as in the proof of lemma 4 it is possible to show that S' and S has the same homotopy type. Also, by the arguments used in the proof of lemma 2 one can show that, S'_j has the same homotopy type as S_j .

Moreover, S' is the disjoint union of the compact sets S'_j . Therefore, $\beta_i(S') = \sum_{1 \leq j \leq L} \beta_i(S'_j)$.

Hence it follows that,

$$\beta_i(S) = \sum_{1 \leq j \leq L} \beta_i(S_j).$$

Since $\chi(S)$ is the alternating sum of the Betti numbers of S , it immediately follows that $\chi(S) = \sum_{1 \leq j \leq L} \chi(S_j)$. □

6.1 The Example of a Solid Torus




Figure 1: S is the solid torus in 3 space

Consider the set bounded by the smooth torus in R^3 (see figure 1.) There are four critical points p_1, p_2, p_3 and p_4 corresponding to the critical values v_1, v_2, v_3 and v_4 respectively, for the projection map onto the X_1 co-ordinate. The Morse data at these points are given in the following table:

Critical point	Index	Tangential data	Normal data
p_1	0	$(D^0 \times D^2, \emptyset)$	$([0, 1], \emptyset)$
p_2	1	$(D^1 \times D^1, \partial D^1 \times D^1)$	$([0, 1], \{0\})$
p_3	1	$(D^1 \times D^1, \partial D^1 \times D^1)$	$([0, 1], \emptyset)$
p_4	2	$(D^2 \times D^0, \partial D^2 \times D^0)$	$([0, 1], \{0\})$

6.2 Proof of Correctness and Complexity Analysis for the Algorithm in the General Case

6.2.1 Proof of Correctness

First note that the set S is compact.

We utilize several results from [24]. We assume that the reader is familiar with algebraic decision trees ([24]). Consider the following fixed degree algebraic decision tree denoted by T . Let the polynomial associated to the root be the polynomial, $P_1 = X_1^2 + \dots + X_k^2 - \Omega$. The associated polynomial to each node at level i , $2 \leq i \leq s$ is P_i . The sets associated to each of the leaf nodes correspond to the 3rd possible sign-conditions on the family \mathcal{P} . Let the set of leaves corresponding to the sign conditions which are non-empty and included in S be denoted by L_S . For a leaf $l \in L_S$, let S_l denote the set associated with it.

For any set A , following [24], we define $\beta'_i(A)$ to be the rank of the i -th homology group $H_i(\bar{A}, \partial A)$, where \bar{A} is the closure of A in the topology of R^k and $\partial A = \bar{A} - A$.

Similarly we define, $\chi'(A) = \sum_{0 \leq i \leq k} (-1)^i \beta'_i(A)$. If A is compact then $\beta'_i(A) = \beta_i(A)$, and $\chi'(A) = \chi(A)$.

We first prove a preliminary lemma.

Lemma 8 *Let X be a semi-algebraic set in R^k which is bounded and semi-closed. For some polynomial $f \in R[X_1, \dots, X_k]$, let $A = X \cap (f \geq 0)$, and $B = X - A$. Then, $\chi'(X) = \chi'(A) + \chi'(B)$.*

Proof: The proof is quite similar to the proof of lemma 5 in [24].

From the exact sequence,

$$\dots \rightarrow H_i(\bar{A} \cup \partial X, \partial X) \rightarrow H_i(\bar{X}, \partial X) \rightarrow H_i(\bar{X}, \bar{A} \cup \partial X) \rightarrow \dots,$$

we have that, $\chi(\bar{X}, \partial X) = \chi(\bar{X}, \bar{A} \cup \partial X) + \chi(\bar{A} \cup \partial X, \partial X)$. Moreover, $\beta_i(\bar{X}, \bar{A} \cup \partial X) = \beta'_i(B)$, and $\beta_i(\bar{A} \cup \partial X, \partial X) = \beta'_i(A)$ (see [24] for a proof). It follows easily that, $\chi'(X) = \chi'(A) + \chi'(B)$. \square

Next, we prove that,

Lemma 9 $\chi(S) = \sum_{l \in L_S} \chi'(S_l)$.

Proof: Note that since S is compact, $\chi'(S) = \chi(S)$. The lemma now follows by a simple induction on the levels of T , starting from the leaf nodes and going up, and using lemma 8 at each node. We omit the details. \square

We next prove that, following the notation introduced in the algorithm, that $\beta'_i(S_j) = \beta_i(U_j, V_j)$, and thus $\chi'(S_j) = \chi(U_j, V_j)$. The proof of this appears in [24] (page 621) and is omitted.

It follows from the exact sequence,

$$\dots \rightarrow H_i(V_j) \rightarrow H_i(U_j) \rightarrow H_i(U_j, V_j) \rightarrow H_{i-1}(V_j) \rightarrow \dots,$$

we immediately deduce that, $\chi(U_j, V_j) = \chi(U_j) - \chi(V_j)$.

This in conjunction with lemma 9 shows that, $\chi(S) = \sum_{1 \leq j \leq m} (\chi(U_j) - \chi(V_j))$, and this proves the correctness of the algorithm.

6.2.2 Complexity Analysis

The cost of computing all the non-empty sign conditions of the family is \mathcal{P} is $s^{k+1}d^{O(k)}$ (see [5]). Moreover, there can be only $\binom{s}{k}(O(d))^k$ such non-empty sign conditions. For each such sign-condition included in S , we call the algorithm for computing the Euler characteristics of basic semi-algebraic sets twice. The sum of the degrees of the polynomials involved in each such call is $O(sd)$. Thus each call costs $(ksd)^{O(k)}$. Hence, the total complexity of the algorithm is bounded by $(ksd)^{O(k)}$.

The bound in the bit model follows easily once we note that bit sizes of the intermediate values are bounded by $L(ksd)^{O(k)}$.

This proves theorem 2.