

# RESOLUTIONS AND THE HOMOLOGY OF MATCHING AND CHESSBOARD COMPLEXES

VICTOR REINER AND JOEL ROBERTS

University of Minnesota

Preprint version, March 1997

ABSTRACT. We generalize work of Lascoux and Joźefiak-Pragacz-Weyman on Betti numbers for minimal free resolutions of ideals generated by  $2 \times 2$  minors of generic matrices and generic symmetric matrices, respectively. Quotients of polynomial rings by these ideals are the classical Segre and quadratic Veronese subalgebras, and we compute the analogous Betti numbers for some natural modules over these Segre and quadratic Veronese subalgebras. The motivation for these results are two-fold:

- Using an old observation on Betti numbers of semigroup modules over semigroup rings in terms of simplicial complexes, we immediately deduce from these results the irreducible decomposition for the symmetric group action on the rational homology of all *chessboard complexes* and complete graph *matching complexes* as studied by Björner, Lovasz, Vrećica and Živaljević
- The class of modules over the Segre rings and quadratic Veronese rings which we consider is closed under the operation of taking *canonical modules*, and hence exposes a pleasant symmetry inherent in these Betti numbers.

## 1. Introduction and Main Results.

Hilbert's Syzygy theorem says that every finitely generated module  $M$  over a polynomial ring  $A = k[x_1, \dots, x_n]$  has a finite resolution by free  $A$ -modules, i.e. an exact sequence

$$(1.1) \quad 0 \rightarrow A^{\beta_h} \rightarrow \dots \rightarrow A^{\beta_1} \rightarrow A^{\beta_0} \rightarrow M \rightarrow 0.$$

In the case where each  $\beta_i$  is as small as possible, this is called a *minimal free resolution*, and the numbers  $\beta_i$  are called the *Betti numbers* of  $M$  over  $A$ . If  $M$  is a graded module over  $A$  it is known that  $\beta_i = \dim_k \operatorname{Tor}_i^A(M, k)$ , where  $k$  is regarded as the trivial  $A$ -module  $k = A/(x_1, \dots, x_n)$ .

In a seminal work, Lascoux [La] computed  $\operatorname{Tor}_i^A(M, k)$  in the case where  $A = k[z_{ij}]$  is the polynomial ring in the entries of a generic  $m \times n$  matrix  $(z_{ij})$ ,  $k$  is a field of characteristic zero, and  $M$  is the quotient ring  $A/I$  where  $I$  is the ideal generated by all  $t \times t$  minors of the matrix  $(z_{ij})$ . In this situation, there is an action of  $GL_m(k) \times GL_n(k)$  on  $\operatorname{Tor}_i^A(M, k)$  which is crucial for Lascoux's

---

1991 *Mathematics Subject Classification.* 14M12, 18G10, 13C14, 14M25, 05E15, 05E25.

analysis, and his result actually describes the decomposition of  $\text{Tor}^A(M, k)$  into  $GL_m(k) \times GL_n(k)$ -irreducibles. Józefiak, Pragacz, and Weyman [JPW] used similar methods to compute  $\text{Tor}^A(M, k)$  where  $A$  is the polynomial ring  $k[z_{ij}]$  in the entries of a generic  $n \times n$  symmetric matrix ( $z_{ij} = z_{ji}$ ),  $I$  is the ideal generated by all  $t \times t$  minors, and  $M$  is the quotient  $A/I$  (again  $k$  has characteristic zero). Their results also rely heavily on the inherent  $GL_n(k)$ -action, and describe the irreducible  $GL_n(k)$ -decomposition of  $\text{Tor}^A(M, k)$ .

The main results of this paper will generalize the results for  $2 \times 2$  minors from [La, JPW], as we now explain. Let  $k[\mathbf{x}, \mathbf{y}] := k[x_1, \dots, x_m, y_1, \dots, y_n]$  be a polynomial ring in two sets of variables of sizes  $m, n$  respectively. The Segre subalgebra  $\text{Segre}(m, n, 0)$  is the subalgebra generated by all monomials  $x_i y_j$  with  $1 \leq i \leq m$  and  $1 \leq j \leq n$ . Letting  $A_{m,n}$  be the polynomial ring  $k[z_{ij}]$  in the entries of a generic  $m \times n$  matrix ( $z_{ij}$ ) as above, there is a surjection

$$\begin{aligned} \phi : A_{m,n} &\rightarrow \text{Segre}(m, n, 0) \\ z_{ij} &\mapsto x_i y_j \end{aligned}$$

The kernel of this surjection is well-known to be the ideal  $I_{m,n}$  generated by the  $2 \times 2$  minors of the matrix  $(x_{ij})$ , and hence  $\text{Segre}(m, n, 0) \cong A_{m,n}/I_{m,n}$ . Identifying  $x_1, \dots, x_m$  and  $y_1, \dots, y_n$  with the bases of two  $k$ -vector spaces  $V \cong k^m$  and  $W \cong k^n$ , then  $k[\mathbf{x}, \mathbf{y}]$  may be viewed as the symmetric algebra

$$\text{Sym}(V \oplus W) = \bigoplus_{a,b \geq 0} \text{Sym}^a V \otimes \text{Sym}^b W.$$

If we define

$$\text{Segre}(m, n, r) = \bigoplus_{a,b \geq 0, a=b+r} \text{Sym}^a V \otimes \text{Sym}^b W$$

for any integer  $r$ , then it is easy to check that  $\text{Segre}(m, n, 0)$  agrees with our earlier definition, and in general  $\text{Segre}(m, n, r)$  is a finitely-generated module over  $\text{Segre}(m, n, 0)$ . Therefore the surjection  $\phi$  endows  $\text{Segre}(m, n, r)$  with the structure of a finitely-generated  $A_{m,n}$ -module. Furthermore, if we identify  $z_{ij}$  with  $x_i \otimes y_j$ , then  $A_{m,n} \cong \text{Sym}(V \otimes W)$ . As a consequence, the product of general linear groups  $GL(V) \times GL(W) \cong GL_m(k) \times GL_n(k)$  acts compatibly on  $A_{m,n}$  and  $\text{Segre}(m, n, r)$  and hence also acts  $\text{Tor}^{A_{m,n}}(\text{Segre}(m, n, r), k)$ . The results of [La] for  $2 \times 2$  minors therefore describe the irreducible decomposition of  $\text{Tor}^{A_{m,n}}(\text{Segre}(m, n, 0), k)$  when  $k$  has characteristic zero, and our first main result generalizes this to  $\text{Segre}(m, n, r)$ . Recall that the irreducible representations  $V^\lambda$  of  $GL_n(k) = GL(V)$  are indexed by partitions  $\lambda = (\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_n \geq 0)$ , and  $|\lambda| := \sum_i \lambda_i$ . Similarly, we denote by  $W^\mu$  the irreducible representation of  $GL_m(k) \cong GL(W)$  indexed by the partition  $\mu$ .

**Theorem 1.1.** *For fields  $k$  of characteristic zero and all  $r \in \mathbb{Z}$ , as a  $GL_m(k) \times GL_n(k)$ -representation,  $\text{Tor}^{A_{m,n}}(\text{Segre}(m, n, r), k)$  is the direct sum of irreducible representations  $V^\lambda \otimes W^\mu$  where  $(\lambda, \mu)$  runs through all pairs of partitions pictured in Figure 1, with*

- $s$  arbitrary,
- $\lambda, \mu$  having at most  $m, n$  parts respectively,

and with the pair  $(\lambda, \mu)$  occurring in homological degree  $s(s-r) + |\alpha| + |\beta|$ . i.e. in  $\text{Tor}_{s(s-r)+|\alpha|+|\beta|}^{A_{m,n}}(\text{Segre}(m, n, 0), k)$ , where  $\alpha, \beta$  are as shown in the figure.

Similarly, if we let  $k[\mathbf{x}] := k[x_1, \dots, x_n]$  then the  $d^{\text{th}}$  Veronese subalgebra  $\text{Veronese}(n, d, 0)$  is the subalgebra of  $k[\mathbf{x}]$  generated by all monomials of degree  $d$ . Letting  $A_n$  be the polynomial ring  $k[z_{ij}]$  in the entries of a generic symmetric  $n \times n$  matrix  $(z_{ij})$  (so  $z_{ij} = z_{ji}$ ) as above, there is a surjection

$$\begin{aligned} \phi : A_n &\rightarrow \text{Veronese}(n, 2, 0) \\ z_{ij} &\mapsto x_i x_j \end{aligned}$$

The kernel of this surjection is well-known to be the ideal  $I_n$  generated by the  $2 \times 2$  minors of the symmetric matrix  $(z_{ij})$ , and hence  $\text{Veronese}(n, 2, 0) \cong A_n/I_n$ . If we identify  $x_1, \dots, x_n$  with the basis of the  $k$ -vector space  $V \cong k^n$ , then  $k[\mathbf{x}]$  may be viewed as the symmetric algebra

$$\text{Sym}V = \bigoplus_{a \geq 0} \text{Sym}^a V.$$

Defining

$$\text{Veronese}(n, d, r) := \bigoplus_{a \equiv r \pmod d} \text{Sym}^a V$$

for any  $r \in \mathbb{Z}/d\mathbb{Z}$ , it is easy to check that  $\text{Veronese}(n, 2, 0)$  agrees with our earlier definition, and in general  $\text{Veronese}(n, d, r)$  is a finitely-generated module over  $\text{Veronese}(n, d, 0)$ . Therefore the surjection  $\phi$  endows  $\text{Segre}(n, 2, r)$  for  $r \equiv 0, 1 \pmod 2$  with the structure of a finitely-generated  $A_n$ -module. Furthermore,  $A_n \cong \text{Sym}(\text{Sym}^2 V)$  so that  $GL(V) \cong GL_n(k)$  acts compatibly on  $A_n$  and  $\text{Veronese}(n, 2, r)$ , and hence also acts on  $\text{Tor}^{A_n}(\text{Veronese}(n, 2, r), k)$ . The results of [JPW] for  $2 \times 2$  minors describe the irreducible decomposition of  $\text{Tor}^{A_n}(\text{Veronese}(n, 2, 0), k)$  when  $k$  has characteristic zero, and our second main result generalizes this to  $\text{Veronese}(n, 2, r)$ .

**Theorem 1.2.** *For fields  $k$  of characteristic zero, and for  $r \equiv 0, 1 \pmod 2$ , as a  $GL(V)$ -representation,  $\text{Tor}^{A_n}(\text{Veronese}(n, 2, r), k)$  is the direct sum of irreducible  $GL(V)$ -representations  $V^\lambda$  where  $\lambda$  runs through all self-conjugate partitions  $\lambda$ , as shown in Figure 2, with*

- $r \equiv |\lambda| \pmod 2$ ,
- $\lambda$  having at most  $n$  parts,

and with  $V^\lambda$  occurring in homological degree  $\binom{s}{2} + |\alpha|$  (i.e. in  $\text{Tor}_{\binom{s}{2} + |\alpha|}^{A_n}(\text{Veronese}(n, 2, r), k)$ ) where  $s$  is the size of the Durfee square of  $\lambda$ , and  $\alpha$  is as shown in Figure 2.

Our original motivation for performing these computations comes from an old observation (Proposition 5.1) that has been re-discovered many times (see e.g. [Sta1, Theorem 7.9], [BH, Proposition 1.1], [CM]). The observation is that in the case where  $M$  is a finitely generated semigroup module over an affine semigroup ring  $S$ , and  $A$  is the polynomial ring in the generators of  $S$ , the groups  $\text{Tor}^A(M, k)$  are the same as the homology groups with coefficients in  $k$  for certain simplicial complexes derived from  $S, M$ . As will be shown in Section 5 (and was alluded to briefly in








FIGURE 2. The self-conjugate partitions  $\lambda$  indexing  $V^\lambda$  which occur in  $\text{Tor}^{A^{m,n}}(\text{Veronese}(n, 2, r), k)$  for  $r = 0, 1$ .

[BH]), this result applies to both  $\text{Segre}(m, n, r)$  and  $\text{Veronese}(n, 2, r)$ . Furthermore, the relevant simplicial complexes include as special cases the  $m \times n$  chessboard complexes  $\Delta_{m,n}$  and the matching complex  $\Delta_n$  for the complete graph on  $n$ -vertices, as defined and studied in [BLVZ]. As a consequence of Theorems 1.1 and 1.2, we will deduce in Section 5 the following result about the irreducible decomposition of the rational homology of the complexes  $\Delta_{m,n}$  and  $\Delta_n$ . For its statement, recall that the irreducible representations  $S^\lambda$  of the symmetric group  $\Sigma_n$  are indexed by partitions  $\lambda$  with  $|\lambda| = n$ .

**Theorem 1.3.** *For fields  $k$  of characteristic zero, as an  $\Sigma_m \times \Sigma_n$ -representation, the reduced homology  $\tilde{H}(\Delta_{m,n}; k)$  is the direct sum of irreducible representations  $S^\lambda \otimes S^\mu$  where  $(\lambda, \mu)$  runs through all pairs of partitions pictured in Figure 1 with*

- $s$  arbitrary,
- $|\lambda| = m, |\mu| = n$  (so that  $r = m - n$ ),

and with the pair  $(\lambda, \mu)$  occurring in  $\tilde{H}_{s(s-r)+|\alpha|+|\beta|}(\Delta_{m,n}; k)$  where  $\alpha, \beta$  are as shown in Figure 1.

Also for fields  $k$  of characteristic zero, as an  $\Sigma_n$ -representation, the reduced homology  $\tilde{H}(\Delta_n; k)$  for  $r = 0, 1$  is the direct sum of irreducible representations  $S^\lambda$  where  $\lambda$  runs through all self-conjugate partitions  $\lambda$ , as shown in Figure 1, with

- $|\lambda| = n$ ,
- $|\lambda| \equiv r \pmod{2}$ ,

and with  $S^\lambda$  occurring in  $\tilde{H}_{\binom{s}{2}+|\alpha|-1}(\Delta_n; k)$ , where  $s$  is the size of the Durfee square of  $\lambda$ , and  $\alpha$  as shown in the figure.

We should point out that although we were not originally aware of it, the results in Theorem 1.3 are not new. In a recent preprint [FrH], Friedman and Hanlon obtain exactly the same description as in Theorem 1.3 for the rational homology of the chessboard complex  $\Delta_{m,n}$ , using a beautiful, but entirely different method involving

the spectral decomposition of *discrete Laplacians* on  $\Delta_{m,n}$ . Their method uncovers further information about the irreducible decompositions of eigenspaces for these Laplacians. Also, the same description as in Theorem 1.3 for the rational homology of the matching complex  $\Delta_n$  obtained in the doctoral thesis of Karagueusian [Ka]. On the other hand, our method uncovers further information of a different sort: Theorem 5.2 describes the group action on the rational homology for all *chessboard complexes with multiplicities*, as defined in [BH, Remark 3.5], and for the class of complexes generalizing  $\Delta_n$  which we call *bounded-degree graph complexes*.

There is another recent motivation for the computation of the rational homology of the complete graph matching complex, ensuing from work of Vassiliev, which is discussed in [BBLSW, §3]. In particular, the table on page 6 of that reference lists homology calculations of  $\tilde{H}_i(\Delta(m, n); k)$  for small values of  $i$ ,  $\text{char}(k)$  and Theorem 1.3 (or the results of [Ka]) accurately predict all of the non-torsion data which occurs in this table.

Finally, we explain one further motivation for generalizing the  $2 \times 2$  minor results of [La, JPW] as in Theorems 1.1 and 1.2. It turns out that the class of modules  $\text{Segre}(m, n, r)$  are closed under the operation  $M \mapsto \Omega(M)$  of taking the *canonical module* of a Cohen-Macaulay module  $M$ , and similarly for the class of modules  $\text{Veronese}(n, d, r)$ . This means that Theorems 1.1 and 1.2 contain a certain duality symmetry (Proposition 2.4) which is not as apparent when considering only the subalgebras  $\text{Segre}(m, n, 0)$  and  $\text{Veronese}(n, 2, 0)$ . We remark that since

$$k[\mathbf{x}, \mathbf{y}] = \bigoplus_{r \in \mathbb{Z}} \text{Segre}(m, n, r)$$

$$k[\mathbf{x}] = \bigoplus_{r \in \mathbb{Z}/d\mathbb{Z}} \text{Veronese}(n, d, r)$$

as  $A_{m,n}$ -modules and  $A_n$ -modules, respectively, it is perhaps not surprising that consideration of *all* of the modules  $\text{Segre}(m, n, r)$  and  $\text{Veronese}(n, 2, r)$  simultaneously leads to a unified result.

The paper is structured as follows. Section 2 discusses the canonical modules of  $\text{Segre}(m, n, r)$  and  $\text{Veronese}(n, 2, r)$ , and explains how Theorems 1.1 and 1.2 respect canonical module duality. It then uses this duality to prove Theorems 1.1 and 1.2. Section 3 sketches the proof of the old observation on Betti numbers of semigroup modules over semigroup rings needed to deduce Theorem 1.3. This section also states a result (Theorem 3.3) generalizing Theorem 1.3 concerning the rational homology of chessboard complexes with multiplicities and bounded-degree graph complexes. Section 4 is devoted to remarks and open problems.

## 2. Canonical modules and the proof of Theorems 1.1, 1.2.

The goal of this section is two-fold. First we review the definition of Cohen-Macaulayness and canonical modules. A general reference for some of this material is [Sta1]. Then we determine when  $\text{Segre}(m, n, r)$  and  $\text{Veronese}(n, d, r)$  are Cohen-Macaulay and identify their canonical modules. We then explain how Theorems 1.1 and 1.2 respect canonical module duality and show how this implies the theorems.

Recall that for a finitely generated graded module  $M$  over the polynomial ring  $A = k[x_1, \dots, x_n]$ , the *homological dimension*  $h = \text{hd}_A(M)$  is the length of a

minimal free resolution for  $M$ , i.e. it is the largest index  $h$  such that  $\mathrm{Tor}_h^A(M, k) \neq 0$ . If we denote by  $d$  the *Krull dimension* of quotient  $A/\mathrm{Ann}_A M$ , then  $A$  is said to be *Cohen-Macaulay* if  $\mathrm{hd}_A(M) = n - d$ . If  $M$  is a module over a finitely generated graded  $k$ -algebra  $R$  which is not a polynomial ring, then one usually takes  $A$  to be a polynomial ring in indeterminates which map to a minimal set of algebra generators for  $R$ , and say that  $M$  is a Cohen-Macaulay  $R$ -module if it is Cohen-Macaulay as an  $A$ -module.

When  $M$  is Cohen-Macaulay, the groups  $\mathrm{Ext}_i(M, A)$  are known to vanish for  $i < h$ , and the *canonical module*  $\Omega(M)$  is defined to be the  $A$ -module  $\mathrm{Ext}_h(M, A)$ . Because of the vanishing of the lower Ext groups, applying the functor  $\mathrm{Hom}_A(A, \cdot)$  to the minimal free resolution (1.1) gives an exact sequence (and hence a minimal free resolution)

$$0 \leftarrow \Omega(M) \leftarrow (A^*)^{\beta_h} \leftarrow \cdots \leftarrow (A^*)^{\beta_1} \leftarrow (A^*)^{\beta_0} \leftarrow 0$$

of  $\Omega(M)$ . We conclude from this resolution that  $\mathrm{Tor}_i^A(M, k)$  and  $\mathrm{Tor}_{h-i}^A(\Omega(M), k)$  are dual as  $k$ -vector spaces for all  $i$ .

**Theorem 2.1.** *Segre( $m, n, r$ ) is a Cohen-Macaulay  $A_{m,n}$ -module if and only if either*

- $0 \leq r \leq m - 1$ , or
- $0 \leq -r \leq n - 1$ , or
- $m = n = 1$  and  $r$  is arbitrary.

*Proof.* Exchanging the roles of  $V$  and  $W$  shows that

$$\mathrm{Segre}(m, n, r) \cong \mathrm{Segre}(n, m, -r),$$

so without loss of generality we may assume that  $r \geq 0$ .

Our approach will utilize some standard geometry of projective spaces, so we begin by reviewing these facts. Let  $X = \mathrm{Proj}(R)$  be a projective variety over the field  $k$ . Given a finitely generated graded  $R$ -module  $M$ , we can construct a coherent sheaf  $\mathcal{L} = \tilde{M}$  on  $X$ ; if  $\mathcal{L}$  is locally free, this sheaf is called a *vector bundle* (see [Har], Chapter II). If  $R = \mathrm{Segre}(m, n, 0)$ , then  $X$  is the Cartesian product  $\mathbb{P}^{m-1} \times \mathbb{P}^{n-1}$  embedded in  $\mathbb{P}^{mn-1}$  via the Segre embedding

$$([x_1 : \cdots : x_m], [y_1 : \cdots : y_n]) \mapsto [x_1 y_1 : x_1 y_2 : \cdots : x_i y_j : \cdots : x_m y_n].$$

This same construction applied to the module  $M = \mathrm{Segre}(m, n, r)$  yields the vector bundle

$$\mathcal{L} = \begin{cases} \mathcal{O}_{\mathbb{P}^{m-1}}(r) \otimes \mathcal{O}_{\mathbb{P}^{n-1}} & \text{if } r \geq 0 \\ \mathcal{O}_{\mathbb{P}^{m-1}} \otimes \mathcal{O}_{\mathbb{P}^{n-1}}(-r) & \text{if } r \leq 0 \end{cases}$$

Conversely, a coherent sheaf  $\mathcal{L}$  yields a finitely generated graded  $R$ -module

$$\Gamma_*(\mathcal{L}) = \bigoplus_{t \in \mathbb{Z}} H^0(X, \mathcal{O}_X(t))$$

and there is a natural homomorphism of graded modules  $\alpha : M \rightarrow \Gamma_*(\tilde{M})$  which maps  $M_t$  to  $H^0(X, \mathcal{L} \otimes \mathcal{O}_X(t))$  for every  $t$ .



This leads to the following criterion for Cohen-Macaulayness, which may be deduced from ([EGA],[Mumford, Lectures on Curves on an algebraic surface]??):  $M$  is a Cohen-Macaulay  $R$ -module if and only if

- (CM1)  $\alpha$  is an isomorphism, so that  $M_t \cong H^0(X, \mathcal{L} \otimes \mathcal{O}_X(t))$  for all  $t \in \mathbb{Z}$ , and  
 (CM2)  $H^s(X, \mathcal{L} \otimes \mathcal{O}_X(t)) = 0$  for all  $t \in \mathbb{Z}$  and  $0 < s < \dim(X)$ .

Here  $\dim(X)$  is the dimension of  $X$  as a projective variety.

We have used above, and will use further below, the following well-known cohomology calculations for a projective space  $\mathbb{P}(U)$  which is the projectivization of a  $k$ -vector space  $U$ :

$$(2.1) \quad \begin{aligned} & H^i(\mathbb{P}(U), \mathcal{O}_{\mathbb{P}(U)}(d)) \\ &= \begin{cases} \text{Sym}^d U & \text{if } i = 0 \text{ and } d \geq 0 \\ (\text{Sym}^{-d - \dim_k U} U)^* & \text{if } i = \dim_k U - 1 \text{ and } d \leq -\dim_k U \\ 0 & \text{otherwise.} \end{cases} \end{aligned}$$

We next apply the above criterion for Cohen-Macaulayness to  $X = \text{Proj}(\text{Segre}(m, n, 0))$  and  $M = \text{Segre}(m, n, r)$ . We first verify that condition CM1 is satisfied for all  $m, n, r$ :

$$\begin{aligned} & H^0(X, \mathcal{L} \otimes \mathcal{O}_X(t)) \\ & \cong H^0(\mathbb{P}^{m-1} \times \mathbb{P}^{n-1}, \mathcal{O}_{\mathbb{P}^{m-1}}(r) \otimes \mathcal{O}_{\mathbb{P}^{n-1}} \otimes \mathcal{O}_{\mathbb{P}^{m-1} \times \mathbb{P}^{n-1}}(t)) \\ & \cong H^0(\mathbb{P}^{m-1} \times \mathbb{P}^{n-1}, \mathcal{O}_{\mathbb{P}^{m-1}}(t+r) \otimes \mathcal{O}_{\mathbb{P}^{n-1}}(t)) \\ & \cong \text{Sym}^{t+r} V \otimes \text{Sym}^t W \\ & \cong M_t. \end{aligned}$$

To verify condition CM2, note that it is vacuously satisfied if  $m = n = 1$  since  $\dim(X) = (m-1) + (n-1) = 0$  in this case. On the other hand, for  $m, n \geq 2$  we can compute

$$\begin{aligned} H^s(X, \mathcal{L} \otimes \mathcal{O}_X(t)) & \cong H^s(\mathbb{P}^{m-1} \times \mathbb{P}^{n-1}, \mathcal{O}_{\mathbb{P}^{m-1}}(t+r) \otimes \mathcal{O}_{\mathbb{P}^{n-1}}(t)) \\ & \cong \bigoplus_{i+j=s} H^i(\mathbb{P}^{m-1}, \mathcal{O}_{\mathbb{P}^{m-1}}(t+r)) \otimes H^j(\mathbb{P}^{n-1}, \mathcal{O}_{\mathbb{P}^{n-1}}(t)), \end{aligned}$$

where the last isomorphism follows from ([Mumford, Lectures on curves on an algebraic surface]??). Using the cohomology calculations (2.1), we conclude that the terms in the last expression which might not vanish are when

- $(i, j) = (0, 0)$ , in which case  $s = i + j = 0$ , so we will never contradict condition CM2.
- $(i, j) = (m-1, n-1)$ , in which case  $s = i + j = (m-1) + (n-1) = \dim X$ , so we will never contradict condition CM2.
- $(i, j) = (m-1, 0)$ , in which case we must have  $t+r \leq -m$  and  $t \geq 0$ . But this is impossible since we assumed  $r \geq 0$ .
- $(i, j) = (0, n-1)$ , in which case we must have  $t \leq -n$  and  $t+r \geq 0$ . So this case would contradict condition CM2, and it will not occur if and only if  $r \leq n-1$ .

The conclusion we reach is that the criterion for condition CM2 is always satisfied if and only if  $r \leq n - 1$ . This completes the proof of the proposition.  $\square$

When  $k$  has characteristic zero, the modules  $\text{Veronese}(n, d, r)$  are always Cohen-Macaulay. One way to see this fact, and also to identify the canonical modules of  $\text{Segre}(n, d, r)$  and  $\text{Veronese}(n, d, r)$ , is through some facts from the invariant theory of finite (or compact) groups which we now review (see [Sta2] for a nice survey).

Recall that if  $G$  is any subgroup of  $GL(V) \cong GL_n(k)$ , then identifying  $R = k[x_1, \dots, x_n]$  with  $\text{Sym}(V)$  defines a  $G$ -action on  $R$ . For the remainder of this section, assume that  $k = \mathbb{C}$ , and we will assume that  $G$  is a compact subgroup of  $GL_n(\mathbb{C})$ . When  $G$  is compact, the subring  $R^G$  of  $G$ -invariant polynomials is finitely generated and Cohen-Macaulay by the methods of Hochster and Eagon [HE]. More generally, for any irreducible character  $\chi$  of  $G$ , one can define the module of  $\chi$ -relative invariants  $R^{G,\chi}$  to be the  $\chi$ -isotypic component of  $R$ . It is shown in [Sta2, Theorem 3.10] that for  $G$  finite,  $R^{G,\chi}$  is a finitely generated Cohen-Macaulay module over  $R^G$ , although Theorem 2.1 will show that that Cohen-Macaulayness can fail for compact groups  $G$  and non-trivial characters  $\chi$ . One can furthermore identify the canonical module  $\Omega(R^{G,\chi})$  in the cases where  $R^{G,\chi}$  is Cohen-Macaulay.

**Lemma 2.2** [Sta2, Remark on p. 502]. *Let  $G \subset GL_n(\mathbb{C})$  be compact,  $\chi$  an irreducible character of  $G$ , det the determinant character of  $G$ , and  $\bar{\chi}$  the conjugate character to  $\chi$ , i.e.  $\bar{\chi}(g) = \overline{\chi(g)}$ . Assume  $R^{G,\chi}$  is a Cohen-Macaulay  $R^G$ -module. Then we have the following isomorphism of graded  $R^G$ -modules*

$$\Omega(R^{G,\chi}) \cong R^{G,\overline{\chi \cdot \det}}$$

up to an overall shift in grading.

We now apply these facts to  $\text{Segre}(m, n, r)$ ,  $\text{Veronese}(n, d, r)$ . Let  $\mathbb{S}^1$  be the circle group

$$\mathbb{S}^1 = \{e^{i\theta}\}_{\theta \in \mathbb{R}/2\pi\mathbb{Z}}$$

embedded as a subgroup  $G \hookrightarrow GL(V \oplus W) \cong GL_{n+m}(\mathbb{C})$  as follows:

$$e^{i\theta} \mapsto \begin{pmatrix} e^{i\theta} \cdot I_V & 0 \\ 0 & e^{-i\theta} \cdot I_W \end{pmatrix}.$$

Here  $I_V, I_W$  denote the identity matrices acting on  $V, W$  respectively. If we let  $R = \text{Sym}(V \oplus W)$  and let  $\chi_r$  denote the character  $\chi(e^{i\theta}) = e^{ri\theta}$  of  $G$ , then it is clear  $\text{Segre}(m, n, 0)$  is the invariant subring  $R^G$ , and  $\text{Segre}(m, n, r)$  is the module of relative invariants  $R^{G,\chi_r}$ .

Similarly, embed the cyclic group  $\mathbb{Z}/d\mathbb{Z}$  as a subgroup  $G \subseteq GL(V) \cong GL_n(\mathbb{C})$  as follows:

$$\zeta \mapsto e^{\frac{2\pi i}{d}} \cdot I_V$$

where  $\zeta$  is a generator of  $\mathbb{Z}/d\mathbb{Z}$ . If we let  $R = \text{Sym}(V)$  and let  $\chi_r$  be the character  $\chi(\zeta) = e^{\frac{2\pi ir}{d}}$  of  $G$ , then it is clear  $\text{Veronese}(n, d, 0)$  is the invariant subring  $R^G$ , and  $\text{Veronese}(n, d, r)$  is the module of relative invariants  $R^{G,\chi_r}$ .

**Corollary 2.3.** *When  $k = \mathbb{C}$ , the Veronese  $(n, d, 0)$ -modules  $\text{Veronese}(n, d, r)$  are always Cohen-Macaulay. Furthermore, when  $k = \mathbb{C}$  and whenever  $\text{Segre}(m, n, n - m - r)$ ,  $\text{Veronese}(n, d, -n - r)$  are Cohen-Macaulay, their canonical modules are described as follows:*

$$\begin{aligned}\Omega(\text{Segre}(m, n, r)) &\cong \text{Segre}(m, n, n - m - r) \\ \Omega(\text{Veronese}(n, d, r)) &\cong \text{Veronese}(n, d, -n - r)\end{aligned}$$

*Proof.* Apply Lemma 2.2 to the previous discussion.  $\square$

As a consequence, the duality between the opposite Tor groups for  $\Omega(M)$  and  $M$  manifest itself in a combinatorial/representation theoretic duality inherent in Theorems 1.1 and 1.2. The next result is the combinatorial manifestation of that duality.

**Proposition 2.4.**

*For  $0 \leq r \leq n - 1$  or  $0 \geq -r \geq m - 1$ , consider the operation of complementing the shapes  $(\lambda, \mu)$  within the rectangular shapes  $((n-1)^m, (m-1)^n)$  and then rotating both shapes 180 degrees. This operation gives an involution which pairs the shapes predicted by Theorem 1.1 to occur in*

$$\text{Tor}_i^{A^{m,n}}(\text{Segre}(m, n, r), \mathbb{C})$$

*with those predicted to occur in*

$$\text{Tor}_j^{A^{m,n}}(\text{Segre}(m, n, n - m - r), \mathbb{C})$$

*where  $i + j = (m - 1)(n - 1)$ .*

*For  $r \equiv 0, 1 \pmod{2}$ , consider the operation of complementing the self-conjugate shape  $\lambda$  within the square shape  $n^n$ , and then rotating 180 degrees. This operation gives an involution which pairs the shapes predicted by Theorem 1.2 to occur in*

$$\text{Tor}_i^{A_n}(\text{Veronese}(n, 2, r), \mathbb{C})$$

*with those predicted to occur in*

$$\text{Tor}_j^{A_n}(\text{Veronese}(n, 2, -n - r), \mathbb{C})$$

*where  $i + j = \binom{n}{2}$ .*

**Remark.** We note that since  $M = \text{Segre}(m, n, r)$ ,  $\text{Veronese}(n, d, r)$  are torsion free modules over the subalgebras  $\text{Segre}(m, n, 0)$ ,  $\text{Veronese}(n, d, 0)$  respectively, in both cases the quotient  $A/\text{Ann}_A(M)$  is isomorphic to the corresponding subalgebra. Since we can compute the Krull dimensions of these subalgebras from the dimensions of the the known dimensions of the Segre and Veronese varieties, we conclude from Cohen-Macaulayness that

$$\begin{aligned}\text{hd}_{A_{m,n}}(\text{Veronese}(m, n, r)) &= mn - (m + n - 1) = (m - 1)(n - 1) \\ \text{hd}_{A_n}(\text{Veronese}(n, 2, r)) &= \binom{n+1}{2} - n = \binom{n}{2}.\end{aligned}$$

Therefore in the dual pairing we should expect  $\text{Tor}_i, \text{Tor}_j$  to pair when  $i + j = h$ , with exactly the values of  $h$  as stated in the Proposition.

*Proof of Proposition 2.4.* Figures 4(a) and 4(b) depict the relevant shapes  $(\lambda, \mu)$  and  $\lambda$  along with their complementary partners within the appropriately sized boxes. As shown, the complementary shapes also fit the format of Figures 1 and 2, with their parameters related to the original parameters as follows. For  $(\lambda, \mu)$  with parameters  $r, s$  the complements  $(\lambda', \mu')$  have parameters  $r' = n - m - r, s' = n - s - 1$  as shown. For  $\lambda$  with Durfee square of size  $s$ , the complement  $\lambda'$  has Durfee square of size  $n - s$ . To see that the homological degrees  $i, j$  of the original shapes and their complements, respectively, add up to the appropriate homological dimension  $h$ , one has two alternatives. One can either do a direct calculation in the two cases, or one can note that in both cases,  $i + j$  is the same as the number of shaded squares depicted in Figures 4(c) and 4(d), and count that the number of shaded squares is the appropriate value  $(m - 1)(n - 1)$  or  $\binom{n}{2}$ .  $\square$

The pairing of shapes inside rectangular boxes as in the previous proposition really *is* a pairing of dual vector spaces, and in fact a pairing of *contragredient* representations, due to the following well-known result.

**Proposition 2.5** [PW, §0.2(c)]. *Let  $\lambda$  be a partition with at most  $n$  parts and all parts of size at most  $m$ . Let  $B$  be a rectangular box with  $n$  rows and  $m$  columns, and let  $\lambda'$  be the complement of  $\lambda$  within the box  $B$ , after rotating 180 degrees. Then as  $GL_n(\mathbb{C})$  representations we have*

$$V_{\lambda'} \cong V_{\lambda}^* \otimes (\det)^{\otimes m}$$

where  $V_{\lambda}^*$  denotes the contragredient representation to  $V_{\lambda}$ , and  $\det \cong \wedge^m(V)$  is the one-dimensional determinant representation of  $GL(V)$ .

As a consequence of this proposition and from the dimensions of the rectangular boxes which occur in Proposition 6.4, we can see what shift in grading is necessary to turn some of the isomorphisms in Corollary 6.2 into graded isomorphisms:

$$\Omega(\text{Segre}(m, n, r)) \cong \text{Segre}(m, n, n - m - r)[(x_1 \cdots x_m)^{n-1}(y_1 \cdots y_n)^{m-1}]$$

$$\Omega(\text{Veronese}(n, 2, r)) \cong \text{Veronese}(n, 2, -n - r)[(x_1 \cdots x_n)^n]$$

where  $M[\mathbf{x}^{\alpha}]$  indicates the module  $M$  with multidegrees shifted up by  $\alpha$ . If  $r = 0$ , we can verify that these conjectural shifts in grading are actually correct: First assume without loss of generality that  $m \leq n$ , and compute the representations

$$\text{Tor}_{(m-1)(n-1)}^{A_{m,n}}(\text{Segre}(m, n, 0), k) = V_{((n-1)^{m-1}, m-1)} \otimes W_{((m-1)^n)}$$

$$\text{Tor}_{\binom{n}{2}}^{A_n}(\text{Veronese}(n, 2, 0), k) = \begin{cases} V_{(n^n)} & \text{if } n \text{ is even} \\ V_{(n^{n-1}, n-1)} & \text{if } n \text{ is odd} \end{cases}$$

known from the results of [La, JPW]. Then compare these with the easily computable representations

$$\text{Tor}_0^{A_{m,n}}(\text{Segre}(m, n, n - m), k) = V_{(n-m)} \otimes W_{\emptyset}$$

$$\text{Tor}_0^{A_n}(\text{Veronese}(n, 2, -n), k) = \begin{cases} V_{\emptyset} & \text{if } n \text{ is even} \\ V_{(1)} & \text{if } n \text{ is odd} \end{cases}$$

with which they are supposed to be paired. As a consequence, we immediately deduce from Theorem 2.1, Proposition 2.3, and Proposition 2.4 the following:



**Corollary 2.6.** *Theorem 1.1 is correct when  $r = 0$  and when  $n - m - r = 0$ . Theorem 1.2 is correct when  $r \equiv 0 \pmod{2}$  and when  $-n - r \equiv 0 \pmod{2}$ .  $\square$*

Finally, from this we can deduce Theorems 1.1, 1.2:

*Proof of Theorems 1.1 and 1.2.* Since Theorems 1.1 and 1.2 both assert that groups  $\text{Tor}_i^A(M, \mathbb{C})$  have certain decompositions as  $GL(V)$ - or  $GL(V) \times GL(W)$ -representations, we first claim they are *polynomial* representations, and hence it suffices to check that they have the correct *characters*, i.e. that the dimensions of weight-spaces  $\text{Tor}_i^A(M, \mathbb{C})_\gamma$  are correct for each weight  $\gamma$ . To see this claim, we use the fact that

$$\text{Tor}_i^A(M, \mathbb{C}) \cong \text{Tor}_i^A(\mathbb{C}, M),$$

and we can compute the latter by tensoring the Koszul resolution of  $\mathbb{C}$  as an  $A$ -module with  $M$  and taking homology of the resulting complex. The terms in the Koszul resolution are exterior powers of  $\mathbb{C}$ -vector spaces tensored with  $A$ , and hence are polynomial representations. Since  $M$  is always a polynomial representation, tensoring with it preserves polynomiality. Then the homology groups of the resulting tensored complex are quotients of submodules of these polynomial representations, and hence also polynomial.

It remains to show that the weight spaces  $\text{Tor}_i^A(M, \mathbb{C})_\gamma$  always have the correct dimension asserted in Theorems 1.1 and 1.2. We start with Theorem 1.2, so  $A = A_n, M = \text{Veronese}(n, 2, r)$  and the group acting is  $GL(V)$ . If  $n, r$  are not already in the cases covered by Corollary 2.6, then  $n$  is even and  $r$  is odd. But then  $n + 1$  is odd, so we know that Theorem 1.2 is correct for  $\text{Veronese}(n + 1, 2, r)$ . Therefore the dimension of each weight space  $\text{Veronese}(n + 1, 2, r)_{\tilde{\gamma}}$  for  $\tilde{\gamma} \in \mathbb{N}^{n+1}$  has the correct dimension predicted by Theorem 1.2. Given a weight  $\gamma \in \mathbb{N}^n$ , we can append an extra coordinate at the end equal to zero to obtain a weight  $\tilde{\gamma} \in \mathbb{N}^{n+1}$ . Proposition 3.2 shows that

$$\begin{aligned} \text{Tor}_i^{A_n}(\text{Veronese}(n, 2, r), \mathbb{C}) &\cong \tilde{H}_{i-1}(\Delta_\gamma; \mathbb{C}) \\ &\cong \tilde{H}_{i-1}(\Delta_{\tilde{\gamma}}; \mathbb{C}) \\ &\cong \text{Tor}_i^{A_{n+1}}(\text{Veronese}(n + 1, 2, r), \mathbb{C}) \end{aligned}$$

where the second isomorphism comes from the crucial (but trivial) fact that  $\Delta_\gamma$  and  $\Delta_{\tilde{\gamma}}$  are isomorphic simplicial complexes. Theorem 1.2 for  $\text{Veronese}(n, 2, r)$  then follows from the well-known fact that the dimension of the weight-space  $V_\gamma^\lambda$  in the irreducible  $GL_n(\mathbb{C})$ - representation  $V^\lambda$  is the same as for the weight space  $V_{\tilde{\gamma}}^\lambda$  in the irreducible  $GL_{n+1}(\mathbb{C})$ - representation  $V^\lambda$ .

A similar argument works for  $\text{Segre}(m, n, r)$ . If  $m, n, r$  are not already in the cases covered by Corollary 2.6, then we can always choose  $m' \geq m$  and  $n' \geq n$  such that  $n - m - r = 0$  and either  $0 \leq r \leq n - 1$  or  $0 \leq -r \leq m - 1$ . Then Theorem 1.1 is correct for  $\text{Segre}(m', n', r)$ , so the dimensions of each of the weight spaces  $\text{Tor}_i^{A^{m,n}}(\text{Segre}(m', n', r), \mathbb{C})_{(\gamma, \delta)}$  are as predicted by Theorem 1.1. A similar argument using Proposition 3.2 then finishes the proof.  $\square$

### 3. Rational homology.

The goal of this section is to sketch the proof of an old observation on Betti numbers of semigroup modules over semigroup rings, and then apply this to deduce Theorem 1.3 and other consequences.

To this end, we introduce some terminology. Let  $\Lambda$  be a finitely generated additive sub-semigroup of  $\mathbb{N}^d$ , and let  $\mathcal{M} \subseteq \mathbb{N}^d$  be a finitely-generated  $\Lambda$ -module, i.e.  $\lambda + \mu \in \mathcal{M}$  for all  $\lambda \in \Lambda$  and  $\mu \in \mathcal{M}$ . The semigroup ring  $k[\Lambda]$  may be identified with a subalgebra of  $k[z_1, \dots, z_d]$  generated by some minimal generating set of monomials  $m_1, \dots, m_n$ . Then  $\mathcal{M}$  gives rise to a finitely generated module  $M = k\mathcal{M}$  over  $k[\Lambda]$  inside  $k[\mathbf{z}]$ , simply by taking the  $k$ -span of all monomials of the form  $\mathbf{z}^\mu$  where  $\mu \in \mathcal{M}$ . Surjecting  $A = k[x_1, \dots, x_n]$  onto  $k[\Lambda]$  by  $x_i \mapsto m_i$ , we endow  $k[\Lambda]$  and  $M$  with the structure of finitely generated  $A$ -modules. Furthermore, all the rings and modules just defined carry an  $\mathbb{N}^d$ -grading, and hence so does  $\text{Tor}_i^A(M, k)$ . We will refer to the  $\alpha^{\text{th}}$ -graded piece of  $\text{Tor}_i^A(M, k)$  by  $\text{Tor}_i^A(M, k)_\alpha$  for  $\alpha \in \mathbb{N}^d$ .

Given  $\mu \in \mathcal{M}$ , define a simplicial complex  $K_\mu$  on vertex set  $[n] := \{1, 2, \dots, n\}$  as follows:

$$K_\mu := \left\{ F \subseteq [n] : \frac{\mathbf{z}^\mu}{\prod_{i \in F} m_i} \in M \right\}$$

**Proposition 3.1**(cf. [BH, Propostion 1.1], [Sta1, Theorem 7.9], [CM], [Stu, Theorem 12.12]). *For  $\Lambda, \mathcal{M}, A, M$  and  $\mu \in \mathcal{M}$  as above, we have*

$$\text{Tor}_i^A(M, k)_\mu \cong \tilde{H}_{i-1}(K_\mu; k)$$

where here  $\tilde{H}$  denotes reduced (simplicial) homology, and all other graded pieces  $\text{Tor}_i^A(M, k)_\alpha$  for  $\alpha \notin \mathcal{M}$  vanish.

*Proof.* For completeness, we sketch the proof as in [BH, Proposition 1.1].

First note that  $\text{Tor}_i^A(M, k)_\mu \cong \text{Tor}_i^A(k, M)_\mu$ . We can compute the right-hand side starting with the well-known Koszul complex  $\mathcal{K}$  resolving  $k$  as an  $A$ -module. This complex has as its  $t^{\text{th}}$  term  $\mathcal{K}_t$  the module  $\bigwedge^t A^n$  which is the free  $A$ -module with  $A$ -basis

$$\{e_{i_1} \wedge \cdots \wedge e_{i_t}\}_{1 \leq i_1 < \cdots < i_t \leq n}$$

and where  $e_i$  carries the same  $\mathbb{N}^d$ -grading as the monomial generator  $m_i$  of  $k[\Lambda]$ . Tensoring the resolution  $\mathcal{K}$  with the  $A$ -module  $M$  gives a complex  $\mathcal{K} \otimes M$ . Fix  $\mu \in \mathbb{N}^d$  and restrict attention to the  $\mu^{\text{th}}$ -graded piece  $(\mathcal{K} \otimes M)_\mu$ , which is a complex of  $k$ -vector spaces. The  $t^{\text{th}}$  term  $(\mathcal{K} \otimes M)_{t, \mu}$  in this complex has typical  $k$ -basis element of the form

$$\mathbf{z}^\gamma e_{i_1} \wedge \cdots \wedge e_{i_t}$$

where  $\mathbf{z}^\gamma \in M$ , and

$$(3.1) \quad \mathbf{z}^\gamma \cdot m_{i_1} \cdots m_{i_t} = \mathbf{z}^\mu.$$

Equation (3.1) implies that  $(\mathcal{K} \otimes M)_\mu$  vanishes unless  $\mu \in \mathcal{M}$ . Furthermore, when  $\mu \in \mathcal{M}$ , note that in the above basis vector,  $\gamma$  is uniquely determined by  $\mu$  and  $\{i_1, \dots, i_t\}$  from equation (3.1). If we identify the above basis vector with the

oriented simplex  $[i_1, \dots, i_t]$  in  $K_\mu$ , one can check that  $(\mathcal{K} \otimes M)_\mu$  is identified with the (augmented) simplicial chain complex  $\tilde{C}.(K_\mu; k)$  up to a shift in grading by 1. The proposition then follows.  $\square$

To apply this result along with Theorems 1.1 and 1.2, we note that  $\text{Segre}(m, n, 0)$  is the semigroup ring for the submonoid of  $\mathbb{N}^m \times \mathbb{N}^n$  generated by  $\{(e_i, e_j)\}_{1 \leq i \leq m, 1 \leq j \leq n}$  where  $e_i$  is the  $i^{\text{th}}$  standard basis vector, and  $\text{Segre}(m, n, r)$  is the semigroup module generated over this semigroup by  $\{(v, 0)\}$  as  $v$  runs over all vectors in  $\mathbb{N}^m$  with  $\sum_i v_i = r$  if  $r > 0$  (and similarly  $\{(0, w)\}$  if  $r < 0$ ). For any multidegree  $(\gamma, \delta)$  occurring in  $\text{Segre}(m, n, r)$ , the complex  $K_{(\gamma, \delta)}$  from Proposition 5.1 is isomorphic to the *chessboard complex with multiplicities*  $\Delta_{\gamma, \delta}$  defined in [BH, Remark 3.5]:  $\Delta_{\gamma, \delta}$  is the simplicial complex whose vertex set is the set of squares on an  $m \times n$  chessboard, and whose simplices are the sets  $F$  of squares having no more than  $\gamma_i$  squares from row  $i$  and no more than  $\delta_j$  squares from row  $j$  for all  $i, j$ . The isomorphism  $K_{(\gamma, \delta)} \cong \Delta_{\gamma, \delta}$  comes from identifying the generator  $(e_i, e_j)$  of the semigroup with the square in row  $i$  and column  $j$  of the chessboard. Note that in the square-free multidegree  $(\gamma, \delta) = ((1, \dots, 1), (1, \dots, 1))$ , this complex  $\Delta_{\gamma, \delta} = \Delta_{m, n}$  is the  $m \times n$  *chessboard complex* considered in [BLVZ], whose vertices are the squares of the chessboard, and whose simplices are the sets of squares which correspond to a placement of rooks on the board so that no two rooks lie in the same row or column. The complex  $\Delta_{3,3}$  is depicted in Figure 4(a).

Similarly,  $\text{Veronese}(n, 2, 0)$  is the semigroup ring for the submonoid of  $\mathbb{N}^n$  generated by  $\{(e_i + e_j)\}_{1 \leq i < j \leq n}$ , and  $\text{Veronese}(n, 2, 1)$  is the semigroup module over this semigroup generated by  $\{e_i\}_{1 \leq i \leq n}$ . For any multidegree  $\gamma$  occurring in  $\text{Veronese}(n, 2, r)$ , the complex  $K_\gamma$  from Proposition 5.1 may be identified with what we will call a *bounded-degree graph complex*  $\Delta_\gamma$ . In the square-free multidegree  $\gamma = (1, \dots, 1)$ , this complex  $\Delta_\gamma$  is the *matching complex*  $\Delta_n$  for a complete graph on  $n$  vertices, as considered in [BLVZ]. The matching complex for a graph  $G$  is the simplicial complex whose vertex set is the set of edges of  $G$ , and whose simplices are the subsets of edges which form a *partial matching*, i.e. an edge-subgraph in which every vertex lies on at most one edge. The isomorphism  $\Delta_{(1, \dots, 1)} \cong \Delta_n$  comes from the fact that  $\Delta_{(1, \dots, 1)}$  cannot use any vertices corresponding to the generators  $\{2e_i\}$  of the semigroup because of the square-free multidegree  $(1, \dots, 1)$ , and the vertex corresponding to the generator  $e_i + e_j$  may be identified with the edge between vertices  $i$  and  $j$  in the complete graph. The matching complex  $\Delta_5$  is depicted in Figure 4(b). For more general  $\gamma$  which are not square-free,  $\Delta_\gamma$  may be described as the bounded-degree graph complex, whose vertices correspond to the possible loops and edges in a complete graph on  $n$  vertices, and whose faces are the subgraphs (with loops allowed) in which the degree of vertex  $i$  is bounded by  $\gamma_i$ . Here a loop on a vertex is counted as adding 2 to the degree of the vertex.

We record the preceding observations in the following Proposition:

**Proposition 3.2.** *For any field  $k$  there are isomorphisms*

$$\begin{aligned} \text{Tor}_i^{A^{m,n}}(\text{Segre}(m, n, r), k)_{(\gamma, \delta)} &\cong \tilde{H}_{i-1}(\Delta_{\gamma, \delta}; k) \\ \text{Tor}_i^{A^n}(\text{Veronese}(n, 2, r), k)_\gamma &\cong \tilde{H}_{i-1}(\Delta_\gamma; k). \end{aligned}$$






FIGURE 4. (a) The chessboard complex  $\Delta_{3,3} = \Delta_{(1,1,1),(1,1,1)}$ . The vertices are labelled by the generators  $x_i y_j$  of Segre(3,3,0). The triangular face with vertices  $x_2 y_1, x_3 y_2, x_1 y_3$  is shown transparent so as not obscure the faces underneath.

(b) The matching complex  $\Delta_5 = \Delta_{(1,1,1,1,1)}$  with vertices labelled by some of the generators  $x_i x_j$  of Veronese(5,2,0). Note that the generators  $x_i^2$  do not appear as vertices, since they do not divide into  $\mathbf{x}^{(1,1,1,1,1)} = x_1 x_2 x_3 x_4 x_5$ .

We next consider symmetries which lead to group actions on these complexes. Notice that one can re-index the rows and columns of the chessboard (which corresponds to permuting the coordinates of  $(\gamma, \delta)$  independently via an element of  $\Sigma_m \times \Sigma_n$ ), without changing the chessboard complex  $\Delta_{\gamma, \delta}$  up to isomorphism. Consequently, we may assume without loss of generality that  $\gamma, \delta$  are partitions, i.e. that their coordinates appear in weakly decreasing order. Therefore  $\gamma, \delta$  are completely determined by the multiplicities of the parts which occur in them, so we can write  $\gamma = 1^{a_1} 2^{a_2} \dots$  and  $\delta = 1^{b_1} 2^{b_2} \dots$ . With this notation, define the *Young or parabolic subgroup*

$$\Sigma_{\mathbf{a}} \times \Sigma_{\mathbf{b}} \hookrightarrow \Sigma_m \times \Sigma_n$$

where  $\Sigma_{\mathbf{a}} = S_{a_1} \times S_{a_2} \times \dots$  and similarly for  $\Sigma_{\mathbf{b}}$ . Then  $\Sigma_{\mathbf{a}} \times \Sigma_{\mathbf{b}}$  acts as a group of simplicial automorphisms of  $\Delta_{\gamma, \delta}$ . Note that in the square-free case case, it is the entire group  $\Sigma_m \times \Sigma_n$  which acts on  $\Delta_{m, n}$ .

Similarly, one can re-index the vertices  $[n]$  of the complete graph (which corre-

sponds to permuting the coordinates of  $\gamma$  via an element of  $\Sigma_n$ , without changing the multi-matching complex  $\Delta_\gamma$  up to isomorphism. Consequently, we may assume without loss of generality that  $\gamma$  is a partition, and completely determined by the multiplicities of the parts which occur, and we can write  $\gamma = 1^{a_1} 2^{a_2} \dots$ . There is then a Young subgroup  $\Sigma_{\mathbf{a}} \hookrightarrow \Sigma_n$  acting as a group of simplicial automorphisms of  $\Delta_\gamma$ , in the square-free case it is the entire symmetric group  $\Sigma_n$  which acts on  $\Delta_n$ .

In order to state our next result, we need to recall the notion of a *weight space* in a  $GL_n(k)$ -representation (see [FuH] for this and other facts from the representation theory of  $GL_n(k)$ ). Let  $\text{diag}(\mathbf{x})$  denote the diagonal matrix in  $GL_n(k)$  having eigenvalues  $x_1, \dots, x_n$ . It is known that when  $k$  has characteristic zero, any finite-dimensional (rational) representation  $U$  of  $GL_n(k)$  decomposes as a direct sum of  $k$ -vector spaces

$$U = \bigoplus_{\gamma \in \mathbb{N}^n} U_\gamma$$

where  $U_\gamma$  is the  $\mathbf{x}^\gamma$ -eigenspace for  $\text{diag}(\mathbf{x})$ , and  $U_\gamma$  is usually called the *weight space* of  $U$  corresponding to the *weight*  $\gamma$ . It is well-known and easy to see that when we act on  $\gamma$  by an element of  $\Sigma_n$  by permuting coordinates we obtain a weight  $\gamma'$  whose weight space  $U_{\gamma'}$  is isomorphic to  $U_\gamma$ . As a consequence, in studying weight spaces we may restrict attention to those with  $\gamma$  a partition (i.e. a *dominant weight*), so  $\gamma = 1^{a_1} 2^{a_2} \dots$ . As in the previous two paragraphs, the Young (parabolic) subgroup  $\Sigma_{\mathbf{a}} \hookrightarrow \Sigma_n \hookrightarrow GL_n(k)$  acts on  $U$  and preserves  $U_\gamma$ , so that  $U_\gamma$  is a  $\Sigma_{\mathbf{a}}$ -representation.

**Theorem 3.3.** *Let  $(\gamma, \delta) \in \mathbb{N}^m \times \mathbb{N}^n$  be partitions,  $r =: |\gamma| - |\delta|$ ,  $\Sigma_{\mathbf{a}} \times \Sigma_{\mathbf{b}}$  the group described above, and  $k$  a field of characteristic zero. Then as a  $\Sigma_{\mathbf{a}} \times \Sigma_{\mathbf{b}}$ -representation, the reduced homology  $\tilde{H}(\Delta_{\gamma, \delta}; k)$  of the chessboard complex with multiplicity  $\Delta_{\gamma, \delta}$  is isomorphic to the direct sum of the  $(\gamma, \delta)$ -weight spaces*

$$\bigoplus_{(\lambda, \mu)} (V^\lambda \otimes W^\mu)_{(\gamma, \delta)}$$

as  $(\lambda, \mu)$  runs through the same indexing set as in Theorem 1.1, and where  $(\lambda, \mu)$  occurs in  $\tilde{H}_{s(s-r)+|\alpha|+|\beta|-1}(\Delta_{\gamma, \delta}; k)$ .

Let  $\gamma \in \mathbb{N}^n$  be a partition,  $r := |\gamma| \bmod 2$ , and  $\Sigma_{\mathbf{a}}$  the permutation group as described above. Then as a  $\Sigma_{\mathbf{a}}$ -representation, the reduced homology  $\tilde{H}(\Delta_\gamma; k)$  of the complete graph matching complex  $\Delta_\gamma$  is isomorphic to the direct sum of the  $\gamma$ -weight spaces

$$\bigoplus_{\lambda} V_\gamma^\lambda$$

as  $\lambda$  runs through the same indexing set as in Theorem 1.2, and where  $\lambda$  occurs in  $\tilde{H}_{\binom{s}{2}+|\alpha|-1}(\Delta_\gamma; k)$ .

*Proof.* By Proposition 3.2 we have

$$\tilde{H}_{i-1}(\Delta_{\gamma, \delta}; k) \cong \text{Tor}_i^{A^{m, n}}(\text{Segre}(m, n, r), k)_{(\gamma, \delta)}$$

where  $r := |\gamma| - |\delta|$ . Since the grading by multidegrees  $(\gamma, \delta) \in \mathbb{N}^m \times \mathbb{N}^n$  is easily seen to coincide with the decomposition of  $\text{Tor}_i^{A^{m,n}}(\text{Segre}(m, n, r), k)$  into  $GL_n(k) \times GL_m(k)$ -weight spaces, the assertion for  $\Delta_{\gamma, \delta}$  then follows from Theorem 1.1.

Similarly, by Proposition 3.2 we have

$$\tilde{H}_{i-1}(\Delta_\gamma; k) = \text{Tor}_i^{A^n}(\text{Veronese}(n, 2, r), k)_\gamma$$

where  $r := |\gamma| \bmod 2$ , and hence the assertion for  $\Delta_\gamma$  follows from Theorem 1.2.  $\square$

*Proof of Theorem 1.3.* We simply recall the fact that the  $(1, \dots, 1)$  weight-space  $V_{(1, \dots, 1)}^\lambda$  of the irreducible  $GL_n(k)$ -representation  $V^\lambda$  affords the irreducible  $S_n$ -representation  $\mathcal{S}^\lambda$ . This fact follows, for example, from a comparison of Weyl's construction of  $V^\lambda$  with the Specht construction of  $\mathcal{S}^\lambda$  (see [FuH, Part I §§4 and 6]).  $\square$

#### Remark 3.4

The reader may be unsatisfied with our general description of the rational homologies  $\tilde{H}_i(\Delta_{\gamma, \delta}; k)$ ,  $\tilde{H}_i(\Delta_\gamma; k)$ , since the answers are stated in terms of the mysterious  $\Sigma_{\mathbf{a}}$ -representations on the weight-spaces  $V_\gamma^\lambda$  of the irreducible  $GL_n(k)$ -representations  $V^\lambda$ . However, we would like to point out that from this description one can deduce their decompositions into irreducible  $\Sigma_{\mathbf{a}}$ -representations, once one knows the irreducible  $\Sigma_{\mathbf{a}}$ -decomposition of  $V_\gamma^\lambda$ . The latter decomposition can be reduced to computations of *Littlewood-Richardson coefficients* and some instances of the *plethysm problem*, as we now explain. The authors would like to thank Mark Shimozono and William Doran for explaining this reduction to us.

Let  $\gamma = 1^{a_1} 2^{a_2} \dots t^{a_t}$ , and let  $GL_{\mathbf{a}}$  be the subgroup

$$GL_{a_1} \times \dots \times GL_{a_t} \hookrightarrow GL_n(k).$$

By restriction,  $\text{Res}_{GL_{\mathbf{a}}}^{GL_n} V_\lambda$  becomes a  $GL_{\mathbf{a}}$ -representation, and as such has a decomposition into  $GL_{\mathbf{a}}$ -irreducibles

$$\text{Res}_{GL_{\mathbf{a}}}^{GL_n} V^\lambda \cong \bigoplus_{(\rho_1, \dots, \rho_t)} (V^{\rho_1} \otimes \dots \otimes V^{\rho_t})^{\oplus c_{\rho_1, \dots, \rho_t}^\lambda}$$

where  $c_{\rho_1, \dots, \rho_t}^\lambda$  is a nonnegative integer which can be computed using the *Littlewood-Richardson rule* [FuH, p. 455]. In fact,  $c_{\rho_1, \dots, \rho_t}^\lambda$  has the following combinatorial interpretation: it is the number of *column-strict tableaux* of shape  $\lambda$  and content  $(\rho_1, \dots, \rho_t)$  which are *Yamanouchi* with respect to each of the alphabets  $1, 2, \dots, a_1$  and  $a_1 + 1, \dots, a_1 + a_2$  and  $a_1 + a_2 + 1, \dots, a_1 + a_2 + a_3$ , etc. We refer the reader to [FH] for the definition of column-strict tableaux and contents. A tableaux is said to be Yamanouchi with respect to an alphabet  $a, a + 1, \dots, b - 1, b$  if when one restricts attention to the entries of the tableaux that lie in this alphabet, and read these entries from right to left in a row, proceeding from the top row and moving down, one obtains a word that has more occurrences of the letter  $i$  than the letter  $i + 1$  in any initial segment, for all  $i$ .

Now using the inclusions

$$\Sigma_{\mathbf{a}} \hookrightarrow GL_{\mathbf{a}} \hookrightarrow GL_n(k)$$

it is easy to see that we have the following isomorphism of  $\Sigma_{\mathbf{a}}$ -representations:

$$V_{\gamma}^{\lambda} \cong \bigoplus_{(\rho_1, \dots, \rho_t)} (V_{1^{a_1}}^{\rho_1} \otimes V_{2^{a_2}}^{\rho_2} \cdots \otimes V_{t^{a_t}}^{\rho_t})^{\oplus c_{\rho_1, \dots, \rho_t}^{\lambda}}$$

where  $\Sigma_{a_i}$  acts on  $V_{i^{a_i}}^{\rho_i}$  for each  $i$ .

Consequently, we have reduced the original description to the following problem: given  $a, b$  nonnegative integers, and  $\rho$  a partition with  $|\rho| = ab$ , how do we decompose the  $\Sigma_a$ -representation  $V_{b^a}^{\rho}$  into  $\Sigma_a$  irreducibles  $\mathcal{S}^{\nu}$  with  $|\nu| = a$ ? By a result of Gay [Ga], this is an instance of the *plethysm problem* [FH, Part I §6]: the multiplicity of  $\mathcal{S}^{\nu}$  as a  $\Sigma_a$ -representation in  $V_{b^a}^{\rho}$  is the same as the multiplicity of  $V^{\rho}$  in  $S_{\nu}(\text{Sym}^b V)$  where  $S_{\nu}$  is the *Schur functor* [ABW] corresponding to  $\nu$ . Alternatively, the above multiplicity is the same as the coefficient of the *Schur function*  $s_{\rho}$  in the *plethysmic composition*  $s_{\nu}[s_{(b)}]$ . Algorithms for computing these multiplicities are contained in [CGR].

#### 4. Remarks and open problems.

Most of the results in this paper have been limited to the case where  $k$  has characteristic zero, since some of the methods involved break down in positive characteristic. The question of how  $\text{Tor}^A(\cdot, k)$  varies with the characteristic of  $k$  for the Segre and quadratic Veronese modules, and consequently what torsion can occur in the homology of the chessboard and matching complexes  $\Delta_{m,n}, \Delta_n$  is wide open, and of great interest. For this reason, we review some of what little is known here.

Hashimoto [Ha1] was the first to show that  $\text{Tor}_3^{A_{5,5}}(\text{Segre}(5, 5, 0), k)$  depends upon whether  $k$  has characteristic 3, and consequently that  $\Delta_{5,5}$  has 3-torsion in its 2-homology (see also [BLVZ, Proposition 2.3] which contains an error that was later corrected). Anderson [An] showed that  $\text{Tor}_5^{A_7}(\text{Veronese}(7, 2, 0), k)$  depends upon whether  $k$  has characteristic 5, by an explicit calculation of  $\tilde{H}_4(\Delta_{\gamma}, \mathbb{Z})$  for the multidegree  $\gamma = (2, 2, 2, 2, 2, 2)$ . The torsion numbers computed in the table on page 6 of [BBLSW] show that  $\tilde{H}_2(\Delta_9)$  has 3-torsion, and hence  $\text{Tor}_3^{A_9}(\text{Veronese}(9, 2, 1), k)_{\gamma}$  for  $\gamma = (1, 1, \dots, 1)$  will depend upon whether the characteristic of  $k$  is 3.

On the other hand, the resolutions of determinantal ideals generated by  $t \times t$  minors of an  $m \times n$  matrix are known to be characteristic-free when  $t = m, m-1, m-2$  by results of Eagon and Northcott [EN], Akin, Buchsbaum, and Weyman [ABW2], and Hashimoto [Ha2], respectively. This implies (using the  $2 \times 2$  minor case) that  $\text{Segre}(m, n, 0)$  has a characteristic free resolution whenever  $m \leq 4$ . Furthermore, exploiting canonical module duality and Proposition 3.2 as in our proof of Theorem 1.1, this implies that  $\text{Segre}(m, n, r)$  has a characteristic-free resolution whenever  $m \leq 4$ . Therefore the integral homology groups for  $\Delta_{\gamma, \delta}$  are torsion-free whenever  $\gamma$  or  $\delta$  has at most 4 parts. Note that this bound of at most 4 parts is tight, since as previously discussed [Ha1, BLVZ] we know that  $\Delta_{5,5} = \Delta_{(1,1,1,1,1), (1,1,1,1,1)}$  has 3-torsion in its homology. This suggests the following problems:

**Problem 4.1.** *Are the complexes  $\Delta_{\gamma, \delta}$  homotopy equivalent to wedges of spheres whenever  $\gamma$  or  $\delta$  has at most 4 parts?*

**Problem 4.2.** *Does Veronese( $n, 2, r$ ) have a characteristic-free resolution for  $n \leq 6$ , i.e. do the complexes  $\Delta_\gamma$  have torsion-free integral homology whenever  $\gamma$  has at most 6 parts? Are these complexes homotopy equivalent to wedges of spheres?*

Another interesting question deals with vanishing theorems and the connectivity of the associated simplicial complexes. In [BLVZ], it is proven that  $\Delta_{m,n}$  is topologically  $(\nu - 2)$ -connected, where

$$\nu = \min \left\{ m, n, \left\lfloor \frac{1}{3}(m + n + 1) \right\rfloor \right\}$$

and it is conjectured that this bound is tight, i.e. that  $\Delta_{m,n}$  has some nontrivial  $(\nu - 1)$ -homology. It is also proven there that  $\Delta_n$  is  $(\lfloor \frac{n+1}{3} \rfloor - 2)$ -connected.

**Problem 4.3.** *Generalize these connectivity results to arbitrary chessboard complexes with multiplicity  $\Delta_{\gamma,\delta}$  and the multi-matching complexes  $\Delta_\gamma$ .*

Such results would give characteristic-free constraints on the resolutions of Segre( $m, n, r$ ) and Veronese( $n, 2, r$ ).

**Acknowledgments.** The authors would like to thank Mark Shimozono and William Doran for the content of Remark 5.4, and Naoki Terai for the references [Sta1, Theorem 7.9] and [Ha2].

#### REFERENCES

- [ABW1] K. Akin, D .A. Buchsbaum, and J. Weyman, *Schur functors and Schur complexes*, Adv. Math. **44** (1982), 207-278.
- [ABW2] K. Akin, D .A. Buchsbaum, and J. Weyman, *Resolutions of determinantal ideals: the submaximal case*, Adv. Math. **39** (1981), 1-30.
- [An] J. Anderson, *Determinantal rings associated with symmetric matrices: a counterexample*, Ph.D. thesis, Univ. of Minnesota (1992).
- [BBLSW] E. Babson, A. Björner, S. Linusson, J. Shareshian, V. Welker, *Complexes of not  $i$ -connected graphs*, preprint (1996).
- [BH] W. Bruns and J. Herzog, *Semigroup rings and simplicial complexes*, Preprint 171 (Osnabrücker Math. Schriften), Universität Osnabrück 1995.
- [BLVZ] A. Björner, L. Lovasz, S. Vrećica, and R. Živaljević, *Chessboard complexes and matching complexes*, J. Lond. Math. Soc. **49** (1994), 25-39.
- [CGR] Y. M. Chen, A. M. Garsia, J. Remmel, *Algorithms for plethysm*, Combinatorics and algebra, (Boulder, Colo. 1983), Amer. Math. Soc., Providence, 1984, pp. 109-153.
- [CM] A. Campillo and C. Marijuan, *Higher order relations for a numerical semigroup*, Séminaire de Théorie des Nombres **3** (1991), 249-260.
- [Ha1] M. Hashimoto, *Determinantal ideals without minimal free resolutions*, Nagoya J. Math. **118** (1990), 203-216.
- [Ha2] ———, *Resolutions of determinantal ideals:  $t$ -minors of  $(t-2) \times n$  matrices*, J. Algebra **142** (1991), 456-491.
- R. Hartshorne, *Algebraic Geometry*, Graduate Texts in Mathematics 52, Springer-Verlag, Berlin, 1977.
- [HE] M. Hochster and J. A. Eagon, *Cohen-Macaulay rings, invariant theory, and the generic perfection of determinantal loci*, Amer. J. Math. **93** (1971), 1020-1058.
- [EN] J. Eagon and D. Northcott, *Generically acyclic complexes and generically perfect ideals*, Prod. Roy. Soc. Ser. A **299** (1967), 147-172.
- [FrH] J. Friedman and P. Hanlon, *On the Betti numbers of chessboard complexes*, preprint (1996).

- [FuH] W. Fulton and J. Harris, *Representation Theory: A first course*, Graduate Texts in Mathematics: Readings in Mathematics, vol. 129, 1991.
- [Ga] D. Gay, *Characters of the Weyl Group of  $SU(n)$  on Zero Weight Spaces and Centralizers of Permutation Representations*, Rocky Moun. J. Math. **6** (1976), 449-455.
- [JPW] T. Józefiak, P. Pragacz, and J. Weyman, *Resolutions of determinantal varieties and tensor complexes associated with symmetric and antisymmetric matrices*, Asterisque **87-88** (1981), 109-189.
- [Ka] D. Karagueusian, *Homology of complexes of degree one graphs*, Ph.D. thesis, Stanford Univ. (1994).
- [La] A. Lascoux, *Syzygies des variétés déterminantales*, Adv. Math. **30** (1978), 202-237.
- [PW] P. Pragacz and J. Weyman, *Complexes associated with trace and evaluation. Another approach to Lascoux' resolution*, Adv. Math. (1985), 163-207.
- [Sta1] R. P. Stanley, *Combinatorics and commutative algebra, 2nd edition*, Progress in Mathematics Vol. 41, Birkhäuser, Boston, 1996.
- [Sta2] R. P. Stanley, *Invariants of finite groups and their applications to combinatorics*, Bull. Amer. Math. Soc. **1** (1979), 475-511.
- [Stu] B. Sturmfels, *Gröbner bases and convex polytopes*, University Lecture Series., vol. 8, American Mathematical Society, Providence, 1996.

SCHOOL OF MATHEMATICS, UNIVERSITY OF MINNESOTA, MINNEAPOLIS, MN 55455

*E-mail address:* roberts@math.umn.edu, reiner@math.umn.edu