



Integration and Homogeneous Functions

Author(s): Jean B. Lasserre

Source: *Proceedings of the American Mathematical Society*, Vol. 127, No. 3, (Mar., 1999), pp. 813-818

Published by: American Mathematical Society

Stable URL: <http://www.jstor.org/stable/119015>

Accessed: 25/04/2008 16:06

Your use of the JSTOR archive indicates your acceptance of JSTOR's Terms and Conditions of Use, available at <http://www.jstor.org/page/info/about/policies/terms.jsp>. JSTOR's Terms and Conditions of Use provides, in part, that unless you have obtained prior permission, you may not download an entire issue of a journal or multiple copies of articles, and you may use content in the JSTOR archive only for your personal, non-commercial use.

Please contact the publisher regarding any further use of this work. Publisher contact information may be obtained at <http://www.jstor.org/action/showPublisher?publisherCode=ams>.

Each copy of any part of a JSTOR transmission must contain the same copyright notice that appears on the screen or printed page of such transmission.

JSTOR is a not-for-profit organization founded in 1995 to build trusted digital archives for scholarship. We enable the scholarly community to preserve their work and the materials they rely upon, and to build a common research platform that promotes the discovery and use of these resources. For more information about JSTOR, please contact support@jstor.org.

INTEGRATION AND HOMOGENEOUS FUNCTIONS

JEAN B. LASSERRE

(Communicated by David H. Sharp)

ABSTRACT. We show that integrating a (positively) homogeneous function f on a compact domain $\Omega \subset R^n$ reduces to integrating a related function on the boundary $\partial\Omega$. The formula simplifies when the boundary $\partial\Omega$ is determined by homogeneous functions. Similar results are also presented for integration of exponentials and logarithms of homogeneous functions.

1. INTRODUCTION

We consider the integration of a continuous (positively) homogeneous function $f : R^n \rightarrow R$ on a compact domain Ω with boundary $\partial\Omega$. Using Euler's identity for homogeneous functions, Green's formula simplifies so that integrating f on Ω reduces to integrating a (simply related) function on the boundary $\partial\Omega$. In the particular case where $\Omega := \{x \in R^n \mid g_i(x) \leq a_i, i = 1, \dots, m\}$ and where the functions g_i are each (positively) homogeneous of degree p_i , the formula is even simpler. Actually, an alternative proof that uses only Euler's formula is also outlined.

We thus extend to more general domains a result in [4] for integrating a homogeneous function on a convex polytope.

A potential application is the integration of arbitrary continuous functions on a compact set Ω . Indeed, as the polynomials are sums of homogeneous polynomials, and are dense in $C(\Omega)$, the space of continuous functions on Ω with the sup norm, or even in $L_1(\Omega)$, one could approximate $\int_{\Omega} f dx$ by $\sum_i \int_{\Omega} P_i dx$, where the P_i 's are homogeneous polynomials, and therefore use the previous result. This result could also be used in finite element methods for the integration of a polynomial on each individual volume element.

Finally, we also provide simple formulas for the integration of exponentials and logarithms of homogeneous functions. For instance, integrating $\log f$ on a convex polytope Ω reduces to integrating $\log f$ on $\partial\Omega$ and to computing the volume of Ω .

2. INTEGRATION OF A HOMOGENEOUS FUNCTION

Let $f : R^n \rightarrow R$ be a real (positively) *homogeneous* function of degree p (or in short, f is p -homogeneous), i.e. $f(\lambda x) = \lambda^p f(x)$ for all $\lambda > 0$, $x \in R^n$. For a (positively) p -homogeneous function that is continuously differentiable, Euler's

Received by the editors July 8, 1997.

1991 *Mathematics Subject Classification.* Primary 65D30.

Key words and phrases. Numerical integration in R^n , homogeneous functions.

formula states that

$$(2.1) \quad pf(x) = \langle \nabla f(x), x \rangle \quad \text{for all } x.$$

2.1. On the Riemann-Green formula. We first treat the 2-dimensional case. Let Ω be a compact domain in R^2 with boundary $\partial\Omega$. Let $P(x, y)$ and $Q(x, y)$ be continuously differentiable on Ω . Then, under some regularity conditions, the well-known Riemann-Green formula (cf. [5]) states that

$$(2.2) \quad \iint_{\Omega} \left(\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} \right) dx dy = \int_{\partial\Omega} P dx + Q dy.$$

This yields:

Lemma 2.1. *Let $f(x, y)$ be a continuously differentiable p -homogeneous function on Ω . Then*

$$(2.3) \quad (p+2) \iint_{\Omega} f dx dy = \int_{\partial\Omega} f(x, y)(x dy - y dx).$$

Proof. As f is p -homogeneous, from (2.1)

$$pf(x, y) = x \frac{\partial f}{\partial x} + y \frac{\partial f}{\partial y} \quad \forall (x, y) \in \Omega.$$

Let $P(x, y) := -yf(x, y)$ and $Q(x, y) := xf(x, y)$ for every $(x, y) \in \Omega$.

$$\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} = 2f(x, y) + x \frac{\partial f}{\partial x} + y \frac{\partial f}{\partial y} = (p+2)f(x, y) \quad \forall (x, y) \in \Omega.$$

Hence, applying the Riemann-Green formula (2.2) to P and Q yields

$$(p+2) \iint_{\Omega} f dx dy = \int_{\partial\Omega} f(x, y)(x dy - y dx),$$

the desired result. □

When $f \equiv 1$ (i.e. $p = 0$), one immediately retrieves the well-known formula

$$\text{area}(\Omega) = \frac{1}{2} \int_{\partial\Omega} x dy - y dx.$$

2.2. The general case. The previous result generalizes to R^n as follows:

Lemma 2.2. *Let $f : R^n \rightarrow R$ be a continuously differentiable p -homogeneous function and Ω a compact domain R^n with boundary $\partial\Omega$. Then*

$$(2.4) \quad (n+p) \int_{\Omega} f d\omega = \int_{\partial\Omega} \langle \vec{M}, \vec{n} \rangle f(M) d\sigma,$$

where \vec{n} is the unit outward-pointing normal to $\partial\Omega$.

Proof. With notation as in [6], let X be the vector field $X := \sum_j x_j \frac{\partial}{\partial x_j}$. From Proposition 2.3 in [6], we have

$$(2.5) \quad \int_{\Omega} \text{div}(X) f d\omega + \int_{\Omega} X f d\omega = \int_{\partial\Omega} \langle X, \vec{n} \rangle f d\sigma.$$

Now, $\text{div}(X) = n$ and from (2.1), $Xf = pf$ so that the result follows. □

Thus, integrating f on Ω reduces to integrating $\langle \vec{M}, \vec{n} \rangle f$ on the boundary $\partial\Omega$. We next show that Lemma 2.2 further simplifies in the case where the boundary $\partial\Omega$ is determined by homogeneous functions.

3. BOUNDARY DETERMINED BY HOMOGENEOUS FUNCTIONS

We now consider the special case where

$$(3.1) \quad \Omega(\subset R^n) := \{x \in R^n \mid g_i(x) \leq a_i, i = 1, \dots, m\},$$

and the functions g_i are each continuously differentiable (positively) p_i -homogeneous functions, $i = 1, \dots, m$. We still assume that Ω is compact. Finally, let $\Omega_i := \{x \in \Omega \mid g_i(x) = a_i\}$ so that $\partial\Omega = \bigcup_{i=1}^m \Omega_i$.

Theorem 3.1. *Assume that f is a continuously differentiable q -homogeneous function and the functions g_i are each continuously differentiable and p_i -homogeneous, $i = 1, \dots, m$. Then*

$$(3.2) \quad (n+q) \int_{\Omega} f d\omega = \sum_{i=1}^m p_i a_i \int_{\Omega_i} \frac{f}{\|\nabla g_i\|} d\sigma.$$

In particular, with $f \equiv 1$, one obtains

$$(3.3) \quad n \times \text{vol}(\Omega) = \sum_{i=1}^m p_i a_i \int_{\Omega_i} \|\nabla g_i\|^{-1} d\sigma.$$

Proof. Lemma 2.2 applies so that

$$(3.4) \quad (n+q) \int_{\Omega} f d\omega = \sum_{i=1}^m \int_{\Omega_i} \langle X, \vec{n} \rangle f d\sigma.$$

Now, $\vec{n} = \|\nabla g_i\|^{-1} \nabla g_i$ on Ω_i . Therefore, using (2.1), we have on Ω_i

$$\langle X, \nabla g_i \rangle = \langle x, \nabla g_i(x) \rangle = p_i g_i(x) = p_i a_i,$$

which yields the desired result. \square

Remark. When Ω is a convex polytope, i.e. $g_i(x) = \langle A_i, x \rangle$ and $p_i = 1$, $i = 1, \dots, m$, then $\|\nabla g_i\| = \|A_i\|$ and thus (3.2) simplifies to

$$(n+q) \int_{\Omega} f d\omega = \sum_{i=1}^m \frac{a_i}{\|A_i\|} \int_{\Omega_i} f d\sigma.$$

In addition, one may iterate the process for $\int_{\Omega_i} f d\sigma$ (cf. [4] for more details).

An alternative proof. Interestingly enough, there is a direct proof that does not use Green's formula. It was used in [4] for the case where Ω is a convex polytope.

Write $a_i = b_i^{p_i}$, $i = 1, \dots, m$, and let

$$(3.5) \quad h(b) := \int_{\Omega} f d\omega = \int_{\{g_i(x) \leq b_i^{p_i}, i=1, \dots, m\}} f dx.$$

It is immediate that $h(\lambda b) = \lambda^{n+q} h(b)$, i.e. h is $(n+q)$ -homogeneous and continuously differentiable. In addition, one may show that

$$\frac{\partial h}{\partial b_i} = p_i b_i^{p_i-1} \int_{\Omega_i} \frac{f}{\|\nabla g_i\|} d\sigma.$$

Hence, applying Euler's formula to h yields

$$(3.6) \quad (n+q)h(b) = \langle \nabla h(b), b \rangle = \sum_{i=1}^m p_i a_i \int_{\Omega_i} \frac{f}{\|\nabla g_i\|} d\sigma,$$

the desired result.

Example. In R^2 , consider the domain

$$\Omega := \{(x, y) \in R^2 \mid xy \leq 1; x^2 + y^2 \leq R^2; x, y \geq 0\}$$

and its volume $\int_{\Omega} d\omega$ (i.e. $f \equiv 1$ is 0-homogeneous).

Note that in (3.3), the coordinate axis boundaries of Ω do not contribute since the boundary values a_i are zero in this case.

Both $g_i(x, y) := xy$ and $g_2(x, y) := x^2 + y^2$ are 2-homogeneous. $\Omega_1 := \{(x, y) \in \Omega \mid xy = 1\}$ and $\Omega_2 := \{(x, y) \in \Omega \mid x^2 + y^2 = R^2\}$. $\|\nabla g_1(M)\| = \sqrt{x^2 + y^2}$ so that on Ω_1 we have $\|\nabla g_1(M)\| = \sqrt{(x^4 + 1)/x^2}$ and $d\sigma = \sqrt{(x^4 + 1)/x^4} dx$, which yields

$$(3.7) \quad \int_{\Omega_1} \|\nabla g_1\|^{-1} d\sigma = \int_{a^{-1}}^a x^{-1} dx = 2 \log(a),$$

where $x = a$ is the solution to $x^2 + y^2 = R^2$, $xy = 1$.

Similarly, $\|\nabla g_2\| = 2R$ on Ω_2 , so that

$$\int_{\Omega_2} \|\nabla g_2\|^{-1} d\sigma = 2 \times (1/2R) \times (R\alpha) = \alpha,$$

where $\alpha = \arctan(a^{-2})$. Hence, Theorem 3.1 yields

$$2 \operatorname{vol}(\Omega) = 4 \log(a) + R^2 \arctan(a^{-2}),$$

i.e. $\operatorname{vol}(\Omega) = 2 \log(a) + R^2 \arctan(a^{-2})/2$. This can be retrieved directly.

If we now take $\Omega := \{(x, y) \mid xy \leq 1; b \leq y \leq a; x \geq 0\}$ with $a \geq b > 0$, we get $\Omega_2 := \{(x, y) \in \Omega \mid y = a\}$ and $\Omega_3 := \{(x, y) \in \Omega \mid -y = -b\}$. Therefore,

$$\int_{\Omega_1} \|\nabla g_1\|^{-1} d\sigma = \log\left(\frac{a}{b}\right); \quad \int_{\Omega_2} \|\nabla g_2\|^{-1} d\sigma = \frac{1}{a}; \quad \int_{\Omega_3} \|\nabla g_3\|^{-1} d\sigma = \frac{1}{b},$$

so that Theorem 3.1 yields

$$2 \operatorname{vol}(\Omega) = 2 \log\left(\frac{a}{b}\right) + a \frac{1}{a} - b \frac{1}{b} = 2 \log\left(\frac{a}{b}\right),$$

i.e. $\operatorname{vol}(\Omega) = \log(a/b)$, which is immediate to check.

4. INTEGRATING EXPONENTIALS AND LOGARITHMS

In this section, we consider exponentials and logarithms of homogeneous functions. Indeed, we show that integrating such functions on Ω reduces to integrating a related function on the boundary $\partial\Omega$.

4.1. Exponentials. We first consider the case of an exponential $e^{h(x)}$ where $h : R^n \rightarrow R$ is q -homogeneous. The particular case where $h = \langle c, x \rangle$ was considered in [1] and [2] and was used to compute the number of integral points in a convex polytope. We have the following result.

Theorem 4.1. *Let Ω be the convex polytope $\{x \in R^n \mid \langle A_i, x \rangle \leq a_i, i = 1, \dots, m\}$.*

(a) Let $h : R^n \rightarrow R$ be q -homogeneous and continuously differentiable. Then, for every $k = 0, 1, \dots$,

$$(4.1) \quad (n + qk) \int_{\Omega} h^k e^h d\omega + q \int_{\Omega} h^{k+1} e^h d\omega = \sum_{i=1}^m \frac{a_i}{\|A_i\|} \int_{\Omega_i} h^k e^h d\sigma.$$

(b) Let $h : R^n \rightarrow R$ be q -homogeneous, continuously differentiable and strictly positive on Ω . Then, for every $k = 1, \dots$,

$$(4.2) \quad (n - qk) \int_{\Omega} h^{-k} e^h d\omega + q \int_{\Omega} h^{-k+1} e^h d\omega = \sum_{i=1}^m \frac{a_i}{\|A_i\|} \int_{\Omega_i} h^{-k} e^h d\sigma.$$

(c) If $p := n/q$ is an integer, then

$$(4.3) \quad q \int_{\Omega} h^{-p+1} e^h d\omega = \sum_{i=1}^m \frac{a_i}{\|A_i\|} \int_{\Omega_i} h^{-p} e^h d\sigma$$

and $\int_{\Omega} h^k e^h d\omega$, $k = -p + 1, \dots, -1, 0, 1, \dots$, can all be expressed as integrals on $\partial\Omega$. In particular,

$$(4.4) \quad q \int_{\Omega} e^h d\omega = \sum_{i=1}^m \frac{a_i}{\|A_i\|} \int_{\Omega_i} e^h [h^{-1} - (p-1)h^{-2} + (p-1)(p-2)h^{-3} + \dots + (-1)^{p+1}(p-1)!h^{-p}] d\sigma.$$

(d) With $h := \langle c, x \rangle$, and for every $\xi \in R^n$, we get

$$(4.5) \quad \langle c, \xi \rangle \int_{\Omega} e^{\langle c, x \rangle} d\omega = \sum_{i=1}^m \frac{\langle A_i, \xi \rangle}{\|A_i\|} \int_{\Omega_i} e^{\langle c, x \rangle} d\sigma.$$

Proof. Again, with $f := h^k e^h$ and with the vector field $X := \sum_i x_i \frac{\partial}{\partial x_i}$, Green's formula yields

$$(4.6) \quad n \int_{\Omega} f d\omega + \int_{\Omega} Xf d\omega = \int_{\partial\Omega} \langle X, \vec{n} \rangle f d\sigma.$$

Using

$$Xf = (kh^{k-1} + h^k) e^h \sum_{i=1}^n x_i \frac{\partial h}{\partial x_i} = q(kh^k + h^{k+1}) e^h, \text{ and } \langle X, \vec{n} \rangle = \frac{a_i}{\|A_i\|} \text{ on } \Omega_i,$$

we obtain

$$(4.7) \quad (n + qk) \int_{\Omega} h^k e^h d\omega + q \int_{\Omega} h^{k+1} e^h d\omega = \sum_{i=1}^m \frac{a_i}{\|A_i\|} \int_{\Omega_i} h^k e^h d\sigma,$$

i.e. (4.1). Similarly, with $f := h^{-k} e^h$ and similar arguments we obtain (4.2).

To get (c), just notice from (4.2) that $\int_{\Omega} h^{-n/q+1} e^h d\omega$ is expressed directly as an integral over $\partial\Omega$, which yields (4.3). Using (4.2) with $k := n/q - 1$ and (4.3), one obtains $\int_{\Omega} h^{-n/q+2} e^h d\omega$ as an integral on $\partial\Omega$. Therefore, iterating and using (4.2) and (4.1), all the $\int_{\Omega} h^k e^h d\omega$, for $k := -n/q + 1, \dots, -1, 0, 1, \dots$, are expressed as integrals on $\partial\Omega$.

To get (d), apply Green's formula (4.6), but now with $X = \sum_i \xi_i \partial/\partial x_i$, so that $\operatorname{div}(X) = 0$. \square

The formula (4.1) remains valid even with $h \equiv 0$ and $k = 0$. With the convention $0^0 = 1$, (4.1) reduces to the volume formula (3.3) of Ω .

Note that (4.5) was obtained in [1] using Stokes' formula, and a similar argument was used to show that it suffices to consider the vertices of the polyhedron. The same can be done using the above argument.

Indeed, let \mathcal{H}_i be the $(n-1)$ -dimensional affine variety that contains Ω_i , and $\{u_1, \dots, u_{n-1}\}$ an orthonormal basis of the associated vector space. $x \in \Omega_i$ can be

written $x_0 + \sum_{i=1}^{n-1} y_i u_i$ with $x_0 \in \Omega_i$, arbitrary. On Ω_i , consider the vector field $\sum_1^{n-1} \xi_i \partial / \partial y_i$. Green's formula yields

$$\left\langle c, \sum_{i=1}^{n-1} \xi_i u_i \right\rangle \int_{\Omega_i} e^{\langle c, x \rangle} d\sigma = \sum_{j \neq i} \langle \xi, \vec{n}_i \rangle \int_{\Omega_{ij}} e^{\langle c, x \rangle} d\nu,$$

where $\Omega_{ij} := \Omega_i \cap \Omega_j$ and \vec{n}_i (in the basis $\{u_1, \dots, u_{n-1}\}$) is the unit outward-pointing normal to Ω_{ij} . Obviously, the process can be repeated up to the 0-dimensional faces, i.e. the vertices of Ω .

4.2. Logarithms. Consider now the function $\log f$ where $f : R^n \rightarrow R$ is continuously differentiable, q -homogeneous and strictly positive on Ω .

Lemma 4.2. *Let $\Omega \subset R^n$ be a compact domain with boundary $\partial\Omega$, and let $f : R^n \rightarrow R$ be continuously differentiable, q -homogeneous and strictly positive on Ω . Then*

$$(4.8) \quad n \int_{\Omega} \log f \, d\omega + q \times \text{vol}(\Omega) = \int_{\partial\Omega} \langle \vec{M}, \vec{n} \rangle \log f(M) \, d\sigma.$$

In addition, if $\Omega := \{x \in R^n \mid \langle A_i, x \rangle \leq a_i, i = 1, \dots, m\}$, and $\Omega_i := \{x \in \Omega \mid \langle A_i, x \rangle = a_i\}$, then

$$(4.9) \quad n \int_{\Omega} \log f \, d\omega + q \times \text{vol}(\Omega) = \sum_{i=1}^m \frac{a_i}{\|A_i\|} \int_{\Omega_i} \log f \, d\sigma.$$

Proof. The proof is again the same as for the exponential. Note that with the vector field $X := \sum_j x_j \partial / \partial x_j$, we have $X \log f = q$ and $\text{div}(X) = n$, so that (2.5) reduces to (4.9). \square

Hence, integrating $\log f$ on a convex polytope Ω reduces to integrating $\log f$ on the boundary $\partial\Omega$ and to computing the volume of the polyhedron.

REFERENCES

1. A. Barvinok, *Computing the volume, counting integral points, and exponential sums*, Discrete & Computational Geometry 10 (1993), pp. 123-141. MR 94d:52005
2. M. Brion, *Points entiers dans les polyèdres convexes*, Ann. Sci. Ec. Norm. Sup., Srie IV, 21 (1988), pp. 653-663. MR 90d:52020
3. J.B. Lasserre, *An analytical expression and an algorithm for the volume of a convex polyhedron in R^n* , J. Optim. Theor. Appl. 39 (1983), pp. 363-377. MR 84m:52018
4. J.B. Lasserre, *Integration on a convex polytope*, Proc. Amer. Math. Soc. 126 (1998), 2433-2441. CMP 97:15
5. G. Strang, *Introduction to Applied Mathematics*, Wellesley-Cambridge Press, 1986. MR 88a:00006
6. M.E. Taylor, *Partial Differential Equations: Basic Theory*, Springer, New York, 1996. MR 98b:35002b

LAAS-CNRS, 7 AVENUE DU COLONEL ROCHE, 31077 TOULOUSE CÉDEX 4, FRANCE
E-mail address: lasserre@laas.fr