

VOLUMES  
AND  
EHRHART POLYNOMIALS  
OF  
CONVEX POLYTOPES

dedicated to the memory of  
Gian-Carlo Rota

Richard P. Stanley  
Department of Mathematics  
M.I.T. 2-375  
Cambridge, MA 02139  
rstan@math.mit.edu  
<http://www-math.mit.edu/~rstan>

Transparencies available at:

<http://www-math.mit.edu/~rstan/trans.html>

$\mathcal{P}$  = convex polytope in  $\mathbb{R}^n$


**integer polytope**: vertices  $\in \mathbb{Z}^n$

$V(\mathcal{P})$  = volume of  $\mathcal{P}$

If  $\mathcal{P}$  is an integer polytope, let

$$\tilde{V}(\mathcal{P}) = n! V(\mathcal{P}) \in \mathbb{Z},$$

the **normalized volume** of  $\mathcal{P}$ .



$$V(P) = 3/2$$

$$\tilde{V}(P) = 3$$

## A Refinement of Volume

Let  $\mathcal{P}$  be an integer polytope and let  $r \geq 1$ . Define

$$r\mathcal{P} = \{rv : v \in \mathcal{P}\}$$

$$i(\mathcal{P}, r) = \#(r\mathcal{P} \cap \mathbb{Z}^n),$$

the **Ehrhart polynomial** of  $\mathcal{P}$ .

- $i(\mathcal{P}, r)$  is a polynomial in  $r$
- $i(\mathcal{P}, 0) = 1$
- If  $r > 0$ , then

$$i(\mathcal{P}, -r) = (-1)^{\dim \mathcal{P}} \#(\text{int}(r\mathcal{P}) \cap \mathbb{Z}^n)$$

- $i(\mathcal{P}, r) = V(\mathcal{P})r^n + O(r^{n-1})$ .

- Let  $\dim \mathcal{P} = n$  and


$$\sum_{r \geq 0} i(\mathcal{P}, r)x^r = \frac{h_0 + h_1x + \cdots + h_nx^n}{(1-x)^{n+1}}.$$

Then  $h_j \in \mathbb{Z}$ ,  $h_j \geq 0$ , and

$$\sum_j h_j = \tilde{V}(\mathcal{P}).$$

**Example.**  $\mathcal{P}$  = unit square:

$$0 \leq x \leq 1, \quad 0 \leq y \leq 1$$



$$\tilde{V}(\mathcal{P}) = 2! \cdot 1 = 2$$

$$i(\mathcal{P}, r) = (r + 1)^2$$

$$i(\mathcal{P}, -r) = (r - 1)^2$$

$$\sum_{r \geq 0} i(\mathcal{P}, r) x^r = \frac{1 + x}{(1 - x)^3}$$

**Example.**  $\mathcal{P}_n =$  unit  $n$ -cube

$$i(\mathcal{P}_n, r) = (r + 1)^n, \quad \tilde{V}(\mathcal{P}_n) = n!$$

If  $w = w_1 w_2 \cdots w_n \in \mathfrak{S}_n$ , then define

$$d(w) = \#\{i : w_i > w_{i+1}\},$$

the number of **descents** of  $w$ . Let

$$A_n(x) = \sum_{w \in \mathfrak{S}_n} x^{1+d(w)},$$

the  $n$ th **Eulerian polynomial**. E.g.,

$$A_3(x) = x + 4x^2 + x^3.$$

Then


$$\sum_{r \geq 0} (r + 1)^n x^n = \frac{A_n(x)/x}{(1 - x)^{n+1}}.$$

## Zonotopes

The **Minkowski sum**  $\mathcal{P} + \mathcal{Q}$  of convex polytopes  $\mathcal{P}, \mathcal{Q} \subseteq \mathbb{R}^n$  is defined by

$$\mathcal{P} + \mathcal{Q} = \{\alpha + \beta : \alpha \in \mathcal{P}, \beta \in \mathcal{Q}\}.$$

A **zonotope** is a Minkowski sum of line segments.



Let  $v_1, \dots, v_k \in \mathbb{R}^n$  and

$$Z(v_1, \dots, v_k) = [0, v_1] + \dots + [0, v_k].$$


**Theorem.** *If  $v_1, \dots, v_k \in \mathbb{Z}^n$ , then*

$$i(Z(v_1, \dots, v_n), r) = \sum_{\substack{S \subseteq \{v_1, \dots, v_k\} \\ (|S|=j) \\ \text{lin. indep.}}} \gcd \left( \begin{array}{l} j \times j \text{ minors of matrix} \\ \text{with rows } v \in S \end{array} \right) r^j.$$

**Corollary.**  $V(Z(v_1, \dots, v_n)) =$

$$\sum_{\substack{S \subseteq \{v_1, \dots, v_k\} \\ S = \text{basis for } \mathbb{R}^n}} |\det(\text{matrix with rows } v \in S)|.$$





$$7r^2 + 4r + 1$$

<u>matrix</u>	<u>gcd</u>	<u>matrix</u>	<u>gcd</u>
$\emptyset$	1	$\begin{bmatrix} 2 & 0 \\ 0 & 1 \end{bmatrix}$	2
$[2 \ 0]$	2	$\begin{bmatrix} 2 & 0 \\ 1 & 2 \end{bmatrix}$	4
$[0 \ 1]$	1	$\begin{bmatrix} 0 & 1 \\ 1 & 2 \end{bmatrix}$	1

**Example.** Let  $e_i = (0, \dots, 0, \overbrace{1}^i, 0, \dots, 0)$ .

$$Z_n = Z(e_i + e_j + e_{n+1} : 1 \leq i < j \leq n) \subset \mathbb{R}^{n+1}$$


Erdős-Gallai:

$$(d_1, \dots, d_{n+1}) \in Z_n \cap \mathbb{Z}^{n+1}$$

$\Leftrightarrow \exists$  (simple) graph on  $1, 2, \dots, n$  with


$$d_j = \deg(j), \quad 1 \leq j \leq n$$

$$d_{n+1} = \frac{1}{2} (d_1 + \dots + d_n).$$



$(2, 2, 3, 0, 1, 4)$

Let  $f(n) = \#$  **distinct** degree sequences  $(d_1, \dots, d_n)$ , so  $f(n) = i(Z_n, 1)$ .




**Theorem.** 
$$\sum_{n \geq 1} f(n) \frac{x^n}{n!} = \frac{1}{2} \left[ \left( 1 + 2 \sum_{n \geq 1} n^n \frac{x^n}{n!} \right)^{1/2} \times \left( 1 - \sum_{n \geq 1} (n-1)^{n-1} \frac{x^n}{n!} \right) + 1 \right] \times \exp \sum_{n \geq 1} n^{n-2} \frac{x^n}{n!}$$

# Catalan Numbers

$$C_n = \frac{1}{n+1} \binom{2n}{n}$$

- triangulations of a convex  $(n+2)$ -gon into  $n$  triangles by  $n - 1$  diagonals that do not intersect in their interiors



- binary trees with  $n$  vertices



- lattice paths from  $(0, 0)$  to  $(n, n)$  with steps  $(0, 1)$  or  $(1, 0)$ , never rising above the line  $y = x$



- sequences of  $n$  1's and  $n$   $-1$ 's such that every partial sum is nonnegative (with  $-1$  denoted simply as  $-$  below)


$111 - - -$        $11 - 1 - -$        $11 - - 1 -$   
 $1 - 11 - -$        $1 - 1 - 1 -$

For 62 additional combinatorial interpretations of  $C_n$ , see Exercise 6.19 of R. Stanley, *Enumerative Combinatorics*, volume 2, Cambridge University Press, 1999

## The Catalanotope

$$A_n^+ = \{e_i - e_j \in \mathbb{R}^{n+1} : 1 \leq i < j \leq n+1\}$$


$$\mathcal{C}_n = \text{conv}(A_n^+ \cup \{0\}) \subset \mathbb{R}^{n+1}.$$



$$\dim \mathcal{C}_n = n$$

Let  $T$  be a tree with vertex set  $\{1, \dots, n+1\}$  and edge set  $E$ . Let  $e_{ij} = e_i - e_j$ . Define the simplex

$$\sigma_T = \text{conv} \left( \{e_{ij} : ij \in E, i < j\} \cup \{0\} \right).$$




$$\sigma_T = \text{conv}\{e_{16}, e_{25}, e_{36}, e_{46}, e_{56}, e_{57}, 0\}$$




$T$  is **alternating** if either every neighbor of vertex  $i$  is less than  $i$  or every neighbor is greater than  $i$ .

$T$  is **noncrossing** if there are not edges  $ik$  and  $jl$  where  $i < j < k < l$ .




**Theorem** (A. Postnikov). *The simplices  $\sigma_T$ , where  $T$  ranges over all noncrossing alternating trees with vertex set  $\{1, \dots, n+1\}$ , are the maximal faces of a triangulation of  $\mathcal{C}_n$ .*



**Easy:**  $\tilde{V}(\sigma_T) = 1$ .

**Lemma.** *The number of noncrossing alternating trees with vertex set  $\{1, \dots, n + 1\}$  is the Catalan number  $C_n$ .*



**Corollary.**  $\tilde{V}(\mathcal{C}_n) = C_n$ .


## Theorem.

$$\sum_{r \geq 0} i(\mathcal{C}_n, r) x^r = \frac{\sum_{j=0}^{n-1} \frac{1}{n} \binom{n}{j} \binom{n}{j+1} x^j}{(1-x)^{n+1}}$$

Here  $\frac{1}{n} \binom{n}{j} \binom{n}{j+1}$  is a **Narayana number**.

# The Parking Function Polytope

(with J. Pitman)



Car  $C_i$  prefers space  $a_i$ . If  $a_i$  is occupied, then  $C_i$  takes the next available space. We call  $(a_1, \dots, a_n)$  a **parking function** (of length  $n$ ) if all cars can park.

$n = 2$  : 11 12 21

$n = 3$  : 111 112 121 211 113 131 311 122  
212 221 123 132 213 231 312 321

**Theorem.** *A sequence  $(a_1, \dots, a_n)$  of positive integers is a parking function if and only if its increasing rearrangement  $b_1 \leq \dots \leq b_n$  satisfies  $b_i \leq i$ .*

111

112 121 211


122 212 221

113 131 311

123 132 213 231 312 321

**Theorem.** (Pyke, 1959; Konheim & Weiss, 1966) The number of parking functions of length  $n$  is  $(n + 1)^{n-1}$ .

**Proof** (Pollak, c. 1974). Add an additional space  $n + 1$ , and arrange the spaces in a circle. Allow  $n + 1$  also as a preferred space.



Now all cars can park, and there will be one empty space.  $A$  is a parking function if and only if the empty space is  $n + 1$ . If  $A = (a_1, \dots, a_n)$  leads to car  $C_i$  parking at space  $p_i$ , then  $(a_1 + j, \dots, a_n + j)$  (modulo  $n + 1$ ) will lead to car  $C_i$  parking at space  $p_i + j$ . Hence exactly one of the vectors

$$(a_1 + i, a_2 + i, \dots, a_n + i) \pmod{n + 1}$$

is a parking function, so

$$f(n) = \frac{(n + 1)^n}{n + 1} = (n + 1)^{n-1}.$$



More generally, Let

$$\alpha = (\alpha_1, \dots, \alpha_n), \quad 0 < \alpha_1 \leq \dots \leq \alpha_n.$$

An  **$\alpha$ -parking function** is a sequence  $(a_1, \dots, a_n) \in \mathbb{P}^n$  whose increasing rearrangement  $b_1 \leq \dots \leq b_n$  satisfies  $b_i \leq \alpha_i$ .

E.g.,  $\alpha = (1, 3) : 11 \ 12 \ 21 \ 13 \ 31$

Ordinary parking functions:

$$\alpha = (1, 2, \dots, n)$$

Let  $\mathbf{L}(\alpha)$  = number of  $\alpha$ -parking functions

**Theorem** (Steck 1968, Gessel 1996):

$$\mathbf{L}(\alpha) = n! \det \left[ \frac{\alpha_i^{j-i+1}}{(j-i+1)!} \right]_{i,j=1}^n$$

**Theorem.** Let  $x_j = \alpha_j - \alpha_{j-1}$  (with  $\alpha_0 = 0$ ). Then

$$L(\alpha) = \sum_{\substack{\text{parking functions} \\ (j_1, \dots, j_n)}} x_{j_1} \cdots x_{j_n}.$$

E.g.,  $\alpha = (1, 3)$ ,  $(x_1, x_2) = (1, 2)$ :

$$x_1^2 + x_1x_2 + x_2x_1 = 5.$$

Given  $x_1, \dots, x_n \in \mathbb{R}_{\geq 0}$ , define  $\mathcal{P} = \mathcal{P}(\mathbf{x}_1, \dots, \mathbf{x}_n) \subset \mathbb{R}^n$  by:  $(y_1, \dots, y_n) \in \mathcal{P}_n$  if

$$y_j \geq 0, \quad y_1 + \dots + y_j \leq x_1 + \dots + x_j$$

for  $1 \leq j \leq n$ .

**Theorem.** (a) Let  $x_1, \dots, x_n \in \mathbb{N}$ .  
Then

$$n! V(\mathcal{P}_n) = L(\alpha),$$

where  $\alpha_{n-j+1} = x_1 + \dots + x_j$ .


(b) Let  $x_1 = a, x_2 = \dots = x_n = b$ .  
Then

$$i(\mathcal{P}_n, 1) = \frac{1}{n!} (a+1)(a+nb+2)(a+nb+3) \cdots (a+nb+n).$$

# Flow Polytopes

(with A. Postnikov)

Let  $G$  a directed graph with vertices  $0, 1, \dots, m$  and edge set  $E$  such that if  $i \rightarrow j$  is an edge, then  $i < j$ . Call  $G$  a **flow graph**. For simplicity we assume that  $(0, i)$  is an edge for  $1 \leq i \leq m$  and  $(j, m)$  is an edge for  $0 \leq j \leq m - 1$ .




Define the **flow polytope**  $\mathcal{F}_G$  to be the set of all  $f \in \mathbb{R}_{\geq 0}^E$  satisfying

$$\sum_{(0,j) \in E} f(0, j) = 1$$

$$\sum_{(j,m) \in E} f(j, m) = 1$$

$$\sum_{i: (i,j) \in E} f(i, j) = \sum_{k: (j,k) \in E} f(j, k),$$

for  $1 \leq j \leq m - 1$ .



total flow out of 0 and into  $m$  is 1

flow into an internal vertex = flow out

**Triangulating  $\mathcal{F}_G$ .** Let

$$E^* = \{(i, j) \in E : 1 \leq i < j \leq m-1\}.$$

$$\text{Let } x^G = \prod_{(i,j) \in E^*} x_{ij}.$$

Continually apply the relations

$$x_{ij}x_{jk} \rightarrow x_{ik}(x_{ij} + x_{jk})$$

until unable to do so.

## Example.

$$x_{12}x_{13}x_{14}x_{23}x_{34}$$

$$\rightarrow x_{13}(x_{12} + x_{23})x_{14}(x_{13} + x_{34})x_{14}$$

$$= 2 \text{ terms} + (x_{12} + x_{23})x_{14}^2x_{13}x_{34}$$

$$\rightarrow 2 \text{ terms} + (x_{12} + x_{23})x_{14}^3(x_{13} + x_{34})$$

$$= 5 \text{ terms} + x_{14}^3x_{23}x_{34}$$

$$\rightarrow 5 \text{ terms} + x_{14}^3x_{24}(x_{23} + x_{34})$$

$$= 7 \text{ terms.}$$

Each monomial  $u$  corresponds to a simplex  $\sigma_u$  with  $\tilde{V}(\sigma_u) = 1$  in a triangulation of  $\mathcal{F}_G$ . Hence  $\tilde{V}(\mathcal{F}_G) = \text{number of monomials}$ .



More generally, let

$$x_{ij}x_{jk} \rightarrow x_{ik}(x_{ij} + x_{jk} - 1)$$

until unable to do so. Then let  $x_{ij} = 1/(1-x)$  and divide by  $(1-x)^m$  to get  $\sum_{r \geq 0} i(\mathcal{F}_G, r)x^r$ .

**Example.**  $x_{12}x_{23} \rightarrow x_{13}(x_{12} + x_{23} - 1)$

$$\begin{aligned} &\rightarrow \frac{1}{(1-x)^4} \frac{1}{1-x} \left( \frac{2}{1-x} - 1 \right) \\ &= \frac{1+x}{(1-x)^6}. \end{aligned}$$

Let  $d_i(G)$  be the outdegree of vertex  $i$  of  $G$ .

**Theorem.**  $\tilde{V}(\mathcal{F}_G)$  is the number of ways to write the vector


$$(d_1 + \cdots + d_{m-2} - m + 2, 1 - d_1, \dots, 1 - d_{m-2})$$

as a sum of vectors  $e_i - e_j$ , where

$$(i - 1, j - 1) \in E(G) \text{ and } j < m$$

(without regard to order).

## Example.



$$\begin{aligned} & (d_1 + \cdots + d_3 - 3, 1 - d_1, 1 - d_2, 1 - d_3) \\ &= (3, -1, -1, -1) \\ &= 3e_{12} + 2e_{23} + e_{34} \\ &= e_{12} + 2e_{13} + e_{34} \\ &= 2e_{12} + e_{13} + e_{23} + e_{34} \\ &= 2e_{12} + e_{14} + e_{23} \\ &= e_{12} + e_{13} + e_{14}. \end{aligned}$$

**Example.**  $G_m$  = **complete flow graph** on  $0, 1, \dots, m$ :  $E = \{(i, j) : 0 \leq i < j \leq m\}$ .

**Chan-Robbins conjecture:**  $\tilde{V}(\mathcal{F}_{G_m}) = C_1 C_2 \cdots C_{m-2}$ .

**Equivalently:** The number of ways to write the vector

$$\left( \binom{m+1}{2}, -m, -m+1, \dots, -1 \right)$$

as a sum of vectors

$$e_i - e_j, \quad 1 \leq i < j \leq m+1.$$

is  $C_1 C_2 \cdots C_m$  (**Kostant's partition function**).

**Corollary.** *Let  $CT$  denote the constant term of a Laurent series. Then*

$$\begin{aligned} & \tilde{V}(\mathcal{F}_{G_m}) = \\ & CT \prod_{i=1}^m (1-x_i)^{-2} \prod_{1 \leq i < j \leq m} (x_j - x_i)^{-1}. \end{aligned}$$


**Theorem** (Morris).

$$\begin{aligned} & CT \prod_{i=1}^m (1-x_i)^{-a} \prod_{i=1}^m x_i^{-b} \prod_{1 \leq i < j \leq m} (x_j - x_i)^{-2c} \\ &= \frac{1}{m!} \prod_{j=0}^{m-1} \frac{\Gamma(a+b+(m-1+j)c)\Gamma(c)}{\Gamma(a+jc)\Gamma(c+jc)\Gamma(b+jc+1)}. \end{aligned}$$

**Corollary** (Zeilberger)  $\tilde{V}(\mathcal{F}_{G_m}) = C_1 \cdots C_{m-2}$ .

Simpler proof?

## One Further Flow Graph



$$\begin{aligned} E(G) = & \{(i, i+1) : 0 \leq i \leq m-1\} \\ & \cup \{(0, i) : 1 \leq i \leq m-1\} \\ & \cup \{(i, m) : 1 \leq i \leq m-1\}. \end{aligned}$$

**Theorem.**  $\tilde{V}(\mathcal{F}_G) = C_{m-1}$ .

$$\begin{aligned}
x_{12}x_{23}x_{34} &\rightarrow x_{13}(x_{12} + x_{23})x_{34} \\
&\rightarrow x_{14}(x_{13} + x_{34})(x_{12} + x_{23}) \\
&\rightarrow x_{14}x_{13}x_{12} + x_{14}x_{13}x_{23} \\
&\quad + x_{14}x_{34}x_{12} + x_{14}x_{24}(x_{23} + x_{34})
\end{aligned}$$

(five terms)