

Pareto Optima of Multicriteria Integer Linear Programs

Jesús A. De Loera

Department of Mathematics, University of California, Davis, Davis, California 95616,
 deloera@math.ucdavis.edu

Raymond Hemmecke, Matthias Köppe

Department of Mathematics, Institute for Mathematical Optimization (IMO),
 Otto-von-Guericke-Universität Magdeburg, 39106 Magdeburg, Germany
 {hemmecke@imo.math.uni-magdeburg.de, mkoepp@imo.math.uni-magdeburg.de}

We settle the computational complexity of fundamental questions related to multicriteria integer linear programs, when the dimensions of the strategy space and of the outcome space are considered fixed constants. In particular we construct: (1) polynomial-time algorithms to determine exactly the number of Pareto optima and Pareto strategies; (2) a polynomial-space polynomial-delay prescribed-order enumeration algorithm for arbitrary projections of the Pareto set; (3) a polynomial-time algorithm to minimize the distance of a Pareto optimum from a prescribed comparison point with respect to arbitrary polyhedral norms; and (4) a fully polynomial-time approximation scheme for the problem of minimizing the distance of a Pareto optimum from a prescribed comparison point with respect to the Euclidean norm.

Key words: integer programming; multicriteria programming; analysis of algorithms, computational complexity; mathematics: combinatorics

History: Accepted by Karen Aardal, Area Editor for Design and Analysis of Algorithms; received July 2007; revised October 2007; accepted January 2008. Published online in *Articles in Advance* September 5, 2008.

1. Introduction

Let $A = (a_{ij})$ be an integral $m \times n$ -matrix and $\mathbf{b} \in \mathbb{Z}^m$ such that the convex polyhedron $P = \{\mathbf{u} \in \mathbb{R}^n: \mathbf{A}\mathbf{u} \leq \mathbf{b}\}$ is bounded. Given k linear functionals $f_1, f_2, \dots, f_k \in \mathbb{Z}^n$, we consider the *multicriterion integer linear programming problem*

$$\begin{aligned} \text{vmin} \quad & (f_1(\mathbf{u}), f_2(\mathbf{u}), \dots, f_k(\mathbf{u})) \\ \text{subject to} \quad & \mathbf{A}\mathbf{u} \leq \mathbf{b}, \\ & \mathbf{u} \in \mathbb{Z}^n, \end{aligned} \tag{1}$$

where vmin is defined as the problem of finding all Pareto optima and a corresponding Pareto strategy. For a lattice point \mathbf{u} , the vector $\mathbf{f}(\mathbf{u}) = (f_1(\mathbf{u}), \dots, f_k(\mathbf{u}))$ is called an *outcome vector*. Such an outcome vector is a *Pareto optimum* for the above problem if and only if there is no other point $\tilde{\mathbf{u}}$ in the feasible set such that $f_i(\tilde{\mathbf{u}}) \leq f_i(\mathbf{u})$ for all i and $f_j(\tilde{\mathbf{u}}) < f_j(\mathbf{u})$ for at least one index j . The corresponding feasible point \mathbf{u} is called a *Pareto strategy*. Thus, a feasible vector is a Pareto strategy if no feasible vector can decrease some criterion without causing a simultaneous increase in at least one other criterion. For general information about the multicriteria problems, see, e.g., Figueira et al. (2005) and Sawaragi et al. (1985).

In general multiobjective problems, the number of Pareto-optimal solutions may be infinite, but throughout the paper we assume that the polyhedron $P = \{\mathbf{u} \in \mathbb{R}^n: \mathbf{A}\mathbf{u} \leq \mathbf{b}\}$ is bounded. This implies that the number of Pareto optima and strategies is finite. There are several well-known techniques to generate Pareto optima. Some popular methods used to solve such problems include, e.g., weighting the objectives or using a so-called global criterion approach; see Ehrgott and Gandibleux (2000). In abnormally nice situations, such as *multicriteria linear programs* (Isermann 1974), one knows a way to generate all Pareto optima, but most techniques reach only some of the Pareto optima.

The purpose of this article is to study the sets of all Pareto optima and strategies of a multicriterion integer linear program using the algebraic structures of generating functions. The set V^{Pareto} of Pareto points can be represented as the multivariate polynomial that is the sum of monomials

$$g(V^{\text{Pareto}}; \mathbf{z}) = \sum \{ \mathbf{z}^{\mathbf{v}}: \mathbf{u} \in P \cap \mathbb{Z}^n, \mathbf{v} = \mathbf{f}(\mathbf{u}) \in \mathbb{Z}^k \text{ is a Pareto optimum} \}, \tag{2}$$

the so-called *generating function* of V^{Pareto} . (Here, we are using the *multiexponent notation* $\mathbf{z}^{\mathbf{v}} = z_1^{v_1} z_2^{v_2} \dots z_k^{v_k}$.)

We remark that the number of monomials in expression (2) usually is exponentially large. Our main theoretical result states that, under the assumption that the number n of variables and the number k of criteria is fixed, we can compute in polynomial time a so-called *short rational generating function* of V^{Pareto} ; i.e., a polynomial-size sum of rational functions of a special kind,

$$\sum_{i \in I} \gamma_i \frac{\mathbf{z}^{\mathbf{c}_i}}{(1 - \mathbf{z}^{\mathbf{d}_{i1}})(1 - \mathbf{z}^{\mathbf{d}_{i2}}) \dots (1 - \mathbf{z}^{\mathbf{d}_{is}})}, \quad (3)$$

that after removing its removable singularities, describes the same polynomial function $g(V^{\text{Pareto}}; \mathbf{z})$. The same can be done for the corresponding Pareto strategies when written in the form

$$g(U^{\text{Pareto}}; \mathbf{x}) = \sum \{ \mathbf{x}^{\mathbf{u}} : \mathbf{u} \in P \cap \mathbb{Z}^n, \mathbf{f}(\mathbf{u}) \text{ is a Pareto optimum} \}. \quad (4)$$

THEOREM 1. *Let the dimension n and the number k of objective functions be fixed.*

(a) *There exists a constant $s = s(k, n)$ and a polynomial-time algorithm for the following problem. Given as input, in binary encoding,*

(I₁) *an integer m , a matrix $A \in \mathbb{Z}^{m \times n}$, a vector $\mathbf{b} \in \mathbb{Z}^m$, and*

(I₂) *linear functionals $f_1, \dots, f_k \in \mathbb{Z}^n$, output, in binary encoding,*

(O₁) *for every i in an index set I an integer γ_i , an integer vector \mathbf{c}_i , and integer vectors \mathbf{d}_{ij} for $j = 1, \dots, s$, such that the generating function $g(V^{\text{Pareto}}; \mathbf{z})$ of the Pareto set V^{Pareto} of the multicriterion integer programming problem (1) has a representation as the rational function*

$$g(V^{\text{Pareto}}; \mathbf{z}) = \sum_{i \in I} \gamma_i \frac{\mathbf{x}^{\mathbf{c}_i}}{(1 - \mathbf{x}^{\mathbf{d}_{i1}})(1 - \mathbf{x}^{\mathbf{d}_{i2}}) \dots (1 - \mathbf{x}^{\mathbf{d}_{is}})}. \quad (5)$$

(b) *The same holds for U^{Pareto} instead of V^{Pareto} .*

(c) *There exists a polynomial-time algorithm for the following problem. Given as input, in binary encoding,*

(I₁) *an integer m , a matrix $A \in \mathbb{Z}^{m \times n}$, a vector $\mathbf{b} \in \mathbb{Z}^m$ and*

(I₂) *linear functionals $f_1, \dots, f_k \in \mathbb{Z}^n$, output, in binary encoding,*

(O₁) *the number $\#V^{\text{Pareto}}$ of Pareto optima and*

(O₂) *the number $\#U^{\text{Pareto}}$ of Pareto strategies of the multicriterion integer programming problem (1).*

(d) *There exist output-sensitive polynomial-time algorithms (more strongly, polynomial-space polynomial-delay prescribed-order enumeration algorithms) for the following two problems. Given as input, in binary encoding,*

(I₁) *an integer m , a matrix $A \in \mathbb{Z}^{m \times n}$, a vector $\mathbf{b} \in \mathbb{Z}^m$, and*

(I₂) *linear functionals $f_1, \dots, f_k \in \mathbb{Z}^n$,*

output, in binary encoding,

(O₁) *in lexicographic order all vectors in the set V^{Pareto} (or U^{Pareto} , respectively),*

where V^{Pareto} and U^{Pareto} are the sets of Pareto optima and strategies of the multicriterion integer programming problem (1).

In contrast, it is known that for a nonfixed dimension it is #P-hard to enumerate Pareto optima and NP-hard to find them (Emelichev and Perepelitsa 1992, Sergienko and Perepelitsa 1991). The proof of Theorem 1, parts (a) and (b), will be given in §2. It is based on the theory of short rational generating functions as developed in Barvinok (1994), Barvinok and Pommersheim (1999), Barvinok and Woods (2003), and Woods (2004). Part (d) of Theorem 1 will be proved in §3; we actually prove the result for arbitrary term orders rather than just the lexicographic order.

For a user that knows some or all of the Pareto optima or strategies, a goal is to select the “best” member of the family. One is interested in selecting one Pareto optimum that realizes the best compromise between the individual objective functions. The quality of the compromise is often measured by the distance of a Pareto optimum \mathbf{v} from a user-defined comparison point $\hat{\mathbf{v}}$. For example, often users take as a good comparison point the so-called *ideal point* $\mathbf{v}^{\text{ideal}} \in \mathbb{Z}^k$ of the multicriterion problem, which is defined as

$$v_i^{\text{ideal}} = \min \{ f_i(\mathbf{u}) : \mathbf{u} \in P \cap \mathbb{Z}^n \}.$$

The criteria of comparison with the point $\hat{\mathbf{v}}$ are quite diverse, but some popular ones include computing the minimum over the possible sums of absolute differences of the individual objective functions, evaluated at the different Pareto strategies, from the comparison point $\hat{\mathbf{v}}$; i.e.,

$$f(\mathbf{u}) = |f_1(\mathbf{u}) - \hat{v}_1| + \dots + |f_k(\mathbf{u}) - \hat{v}_k|, \quad (6a)$$

or the maximum of the absolute differences,

$$f(\mathbf{u}) = \max \{ |f_1(\mathbf{u}) - \hat{v}_1|, \dots, |f_k(\mathbf{u}) - \hat{v}_k| \}, \quad (6b)$$

over all Pareto optima $(f_1(\mathbf{u}), \dots, f_k(\mathbf{u}))$. Another popular criterion, sometimes called the *global criterion*, is to minimize the sum of relative distances of the individual objectives from their known minimal values; i.e.,

$$f(\mathbf{u}) = \frac{f_1(\mathbf{u}) - v_1^{\text{ideal}}}{|v_1^{\text{ideal}}|} + \dots + \frac{f_k(\mathbf{u}) - v_k^{\text{ideal}}}{|v_k^{\text{ideal}}|}. \quad (6c)$$

We stress that if we take any one of these functions as an objective function of an integer program, the optimal solution will be, in general, a non-Pareto solution of the multicriterion problem (1). In contrast,

we show in this paper that by encoding Pareto optima and strategies as a rational function, we avoid this problem because we evaluate the objective functions directly on the space of Pareto optima.

All of the above criteria (6) measure the distance from a prescribed point with respect to a *polyhedral norm*. In §4, we prove:

THEOREM 2. *Let the dimension n and the number k of objective functions be fixed.*

(a) *There exists a polynomial-time algorithm for the following problem. Given as input, in binary encoding,*

(I₁) *an integer m , a matrix $A \in \mathbb{Z}^{m \times n}$, a vector $\mathbf{b} \in \mathbb{Z}^m$,*

(I₂) *linear functionals $f_1, \dots, f_k \in \mathbb{Z}^n$,*

(I₃) *a prescribed point $\hat{\mathbf{v}} \in \mathbb{Z}^k$, and*

(I₄) *a vertex or inequality description of a rational polytope $Q \subseteq \mathbb{R}^k$,*

output, in binary encoding,

(O₁) *a Pareto optimum \mathbf{v} of the multicriterion integer programming problem (1) that minimizes the distance $\|\mathbf{v} - \hat{\mathbf{v}}\|_Q$ from the prescribed point $\hat{\mathbf{v}}$, where $\|\cdot\|_Q$ is the polyhedral norm whose unit ball is Q .*

(b) *There exists a polynomial-space polynomial-delay enumeration algorithm for the following problem. Given as input the same data as in (a), output, in binary encoding,*

(O₁) *all Pareto optima \mathbf{v} of the multicriterion integer programming problem (1) in the order of increasing distances $\|\mathbf{v} - \hat{\mathbf{v}}\|_Q$ from the prescribed point $\hat{\mathbf{v}}$, where $\|\cdot\|_Q$ is the polyhedral norm whose unit ball is Q .*

Often users are actually interested in finding a Pareto optimum that minimizes the Euclidean distance from a prescribed comparison point $\hat{\mathbf{v}}$,

$$f(\mathbf{u}) = \sqrt{|f_1(\mathbf{u}) - \hat{v}_1|^2 + \dots + |f_k(\mathbf{u}) - \hat{v}_k|^2}, \quad (7)$$

but to our knowledge no method of the literature gives a satisfactory solution to that problem. In §4, however, we prove the following theorem, which gives a very strong approximation result—a fully polynomial-time approximation scheme (FPTAS).

THEOREM 3. *Let the dimension n and the number k of objective functions be fixed. Then, there exists a polynomial-time algorithm for the following problem. Given as input*

(I₁) *an integer m , a matrix $A \in \mathbb{Z}^{m \times n}$, a vector $\mathbf{b} \in \mathbb{Z}^m$, linear functionals $f_1, \dots, f_k \in \mathbb{Z}^n$, all in binary encoding,*

(I₂) *a prescribed comparison point $\hat{\mathbf{v}} \in \mathbb{Z}^k$ in binary encoding, and*

(I₃) *a positive rational number $1/\epsilon$ in unary encoding,*

output, in binary encoding,

(O₁) *a Pareto optimum \mathbf{v}_ϵ of the multicriterion integer programming problem (1) such that*

$$\|\mathbf{v}_\epsilon - \hat{\mathbf{v}}\| \leq (1 + \epsilon) \min\{\|\mathbf{v} - \hat{\mathbf{v}}\| : \mathbf{v} \in V^{\text{Pareto}}\},$$

where $\|\cdot\|$ denotes the Euclidean norm and V^{Pareto} is the set of Pareto optima of (1).

We actually prove this theorem in a somewhat more general setting, using an arbitrary norm whose unit ball is representable by a homogeneous polynomial inequality.

2. The Rational Function Encoding of All Pareto Optima

We give a very brief overview of the theory of rational generating functions necessary to establish Theorem 1. For full details, we recommend Barvinok (1994), Barvinok and Pommersheim (1999), Barvinok and Woods (2003), De Loera et al. (2006), and the references therein. Barvinok (1994) gave an algorithm for counting the lattice points in $P = \{\mathbf{u} \in \mathbb{R}^n : A\mathbf{u} \leq \mathbf{b}\}$ in polynomial time when the dimension n of the feasible polyhedron is a constant. The input for Barvinok's algorithm is the binary encoding of the integers a_{ij} and b_i , and the output is a formula for the multivariate generating function

$$g(P; \mathbf{x}) = \sum_{\mathbf{u} \in P \cap \mathbb{Z}^n} \mathbf{x}^{\mathbf{u}},$$

where $\mathbf{x}^{\mathbf{u}}$ is an abbreviation of $x_1^{u_1} x_2^{u_2} \dots x_n^{u_n}$. This long polynomial with exponentially many monomials is encoded as a much shorter sum of rational functions of the form

$$g(P; \mathbf{x}) = \sum_{i \in I} \gamma_i \frac{\mathbf{x}^{\mathbf{c}_i}}{(1 - \mathbf{x}^{\mathbf{d}_{i1}})(1 - \mathbf{x}^{\mathbf{d}_{i2}}) \dots (1 - \mathbf{x}^{\mathbf{d}_{im}})}. \quad (8)$$

Barvinok and Woods (2003) further developed a set of powerful manipulation rules for using these short rational functions in Boolean constructions on various sets of lattice points.

THEOREM 4 (BOOLEAN OPERATIONS LEMMA; COROLLARY 3.7 IN BARVINOK AND WOODS 2003). *Let m and l be fixed integers, and let $\phi : \{0, 1\}^m \rightarrow \{0, 1\}$ be any Boolean function such that $\phi(\mathbf{0}) = 0$. Then, there exists a constant $s = s(l, m)$ and a polynomial-time algorithm for the following problem. Given as input, in binary encoding,*

(I₁) *the dimension n and*

(I₂) *rational generating functions*

$$g(S_p; \mathbf{x}) = \sum_{i \in I_p} \gamma_{pi} \frac{\mathbf{x}^{\mathbf{c}_{pi}}}{(1 - \mathbf{x}^{\mathbf{d}_{pi1}}) \dots (1 - \mathbf{x}^{\mathbf{d}_{pis}})},$$

of m finite sets $S_p \subseteq \mathbb{Z}^n$, represented by the rational numbers γ_{pi} , integer vectors \mathbf{c}_{pi} , and \mathbf{d}_{pij} for $p = 1, \dots, m$, $i \in I_p$, $j = 1, \dots, l_{mp}$ such that the numbers l_{mp} of terms in the denominators are at most l ,

output, in binary encoding,

(O₁) *rational numbers γ_i , integer vectors \mathbf{c}_i , \mathbf{d}_{ij} for $i \in I$, $j = 1, \dots, s_i$, where $s_i \leq s$ such that*

$$g(S; \mathbf{x}) = \sum_{i \in I} \gamma_i \frac{\mathbf{x}^{\mathbf{c}_i}}{(1 - \mathbf{x}^{\mathbf{d}_{i1}}) \dots (1 - \mathbf{x}^{\mathbf{d}_{is_i}})}$$

is a rational generating function of the finite set S that is the Boolean combination of S_1, \dots, S_p corresponding to the function ϕ .

We will use the Boolean operations lemma to extract special monomials present in the expansion of a generating function. The essential step in the algorithm of Theorem 4 is the use of the *Hadamard product* (Barvinok and Woods 2003, Definition 3.2) and a special monomial substitution. The Hadamard product is a bilinear operation on rational functions (we denote it by $*$). The computation is carried out for pairs of summands as in (8). Note that the Hadamard product $m_1 * m_2$ of two monomials m_1, m_2 is zero unless $m_1 = m_2$.

Another key subroutine introduced by Barvinok and Woods (2003) is the following projection theorem.

THEOREM 5 (PROJECTION THEOREM; THEOREM 1.7 IN BARVINOK AND WOODS 2003). *Let the dimension n be a fixed constant. Then, there exists a constant $s(n)$ and a polynomial-time algorithm for the following problem. Given as input, in binary encoding,*

(I₁) *an inequality description of a rational polytope $P \subset \mathbb{R}^n$,*

(I₂) *a positive integer k , and*

(I₃) *an integral matrix $T \in \mathbb{Z}^{k \times n}$,*

output, in binary encoding,

(O₁) *rational numbers γ_i , integer vectors $\mathbf{c}_i, \mathbf{d}_{ij}$ for $i \in I, j = 1, \dots, s_i$, where $s_i \leq s$ such that*

$$g(S; \mathbf{x}) = \sum_{i \in I} \gamma_i \frac{\mathbf{x}^{\mathbf{c}_i}}{(1 - \mathbf{x}^{\mathbf{d}_{i1}}) \dots (1 - \mathbf{x}^{\mathbf{d}_{is_i}})}$$

is a rational generating function of the set $S = T(P \cap \mathbb{Z}^n)$.

One has to be careful when using the lemmas above that the sets in question are finite. The proof of Theorem 1 will require us to project and intersect sets of lattice points represented by rational functions. We cannot, in principle, do those operations for infinite sets of lattice points. Fortunately, in our setting, it is possible to restrict our attention to finite sets.

Finally, one important comment. If we want to count the points of a lattice point set S , such as the set of Pareto optima, it would apparently suffice to substitute 1 for all the variables x_i of the generating function

$$g(S; \mathbf{x}) = \sum_{\mathbf{u} \in S} \mathbf{x}^{\mathbf{u}} = \sum_{i \in I} \gamma_i \frac{\mathbf{x}^{\mathbf{c}_i}}{(1 - \mathbf{x}^{\mathbf{d}_{i1}})(1 - \mathbf{x}^{\mathbf{d}_{i2}}) \dots (1 - \mathbf{x}^{\mathbf{d}_{im}})}$$

to get the specialization $|S| = g(S; \mathbf{x} = \mathbf{1})$. But this cannot be done directly due to the singularities in the rational function representation. Instead, choose a generic vector $\boldsymbol{\lambda} = (\lambda_1, \dots, \lambda_n)$ and substitute each of the variables x_i by $e^{t\lambda_i}$. Then, we get

$$g(S, \mathbf{e}^{t\boldsymbol{\lambda}}) = \sum_{i \in I} \gamma_i \frac{e^{t\langle \boldsymbol{\lambda}, \mathbf{c}_i \rangle}}{(1 - e^{t\langle \boldsymbol{\lambda}, \mathbf{d}_{i1} \rangle})(1 - e^{t\langle \boldsymbol{\lambda}, \mathbf{d}_{i2} \rangle}) \dots (1 - e^{t\langle \boldsymbol{\lambda}, \mathbf{d}_{im} \rangle})}$$

Counting the number of lattice points is the same as computing the constant terms of the Laurent series for each summand and adding them up. This can be done efficiently using elementary complex residue techniques:

THEOREM 6 (SPECIALIZATION THEOREM; BARVINOK 1994, BARVINOK AND POMMERSHEIM 1999). *Let n and l be fixed integers. Then, there exists a polynomial-time algorithm for the following problem. Given as input, in binary encoding,*

(I₁) *rational numbers γ_i , integer vectors $\mathbf{c}_i, \mathbf{d}_{ij}$ for $i \in I, j = 1, \dots, l_i$, where $l_i \leq l$ such that $\mathbf{1}$ is a removable singularity of the rational function*

$$g(P; \mathbf{x}) = \sum_{i \in I} \gamma_i \frac{\mathbf{x}^{\mathbf{c}_i}}{(1 - \mathbf{x}^{\mathbf{d}_{i1}}) \dots (1 - \mathbf{x}^{\mathbf{d}_{il_i}})},$$

output, in binary encoding,

(O₁) *the number*

$$g(P; \mathbf{x} = \mathbf{1}) = \lim_{\mathbf{x} \rightarrow \mathbf{1}} \sum_{i \in I} \gamma_i \frac{\mathbf{x}^{\mathbf{c}_i}}{(1 - \mathbf{x}^{\mathbf{d}_{i1}}) \dots (1 - \mathbf{x}^{\mathbf{d}_{il_i}})}.$$

With these powerful results at hand, we can prove our first theorem.

PROOF OF THEOREM 1 (PARTS (a)–(c)). The proof of part (a) has two steps:

Step 1. For $i = 1, \dots, k$, let $\bar{v}_i \in \mathbb{Z}$ be an upper bound of polynomial encoding size for the value of f_i over P . Such a bound exists because of the boundedness of P , and it can be computed in polynomial time by linear programming. We will denote the vector of upper bounds by $\bar{\mathbf{v}} \in \mathbb{Z}^k$. We consider the *truncated multi-epigraph* of the objective functions f_1, \dots, f_k over the linear relaxation of the feasible region P ,

$$P_{f_1, \dots, f_k}^{\geq} = \{(\mathbf{u}, \mathbf{v}) \in \mathbb{R}^n \times \mathbb{R}^k : \mathbf{u} \in P, \bar{v}_i \geq v_i \geq f_i(\mathbf{u}) \text{ for } i = 1, \dots, k\}, \quad (9)$$

which is a rational convex polytope in $\mathbb{R}^n \times \mathbb{R}^k$. Let $V^{\geq} \subseteq \mathbb{Z}^k$ denote the integer projection of $P_{f_1, \dots, f_k}^{\geq}$ on the \mathbf{v} variables, i.e., the set

$$V^{\geq} = \{\mathbf{v} \in \mathbb{Z}^k : \exists \mathbf{u} \in \mathbb{Z}^n \text{ with } (\mathbf{u}, \mathbf{v}) \in P_{f_1, \dots, f_k}^{\geq} \cap (\mathbb{Z}^n \times \mathbb{Z}^k)\}. \quad (10)$$

Clearly, the vectors in V^{\geq} are all integer vectors in the outcome space that are weakly dominated by some outcome vector $(f_1(\mathbf{u}), f_2(\mathbf{u}), \dots, f_k(\mathbf{u}))$ for a feasible solution \mathbf{u} in $P \cap \mathbb{Z}^n$; however, we have truncated away all outcome vectors that weakly dominate the computed bound $\bar{\mathbf{v}}$. Let us consider the generating function of V^{\geq} , the multivariate polynomial

$$g(V^{\geq}; \mathbf{z}) = \sum \{\mathbf{z}^{\mathbf{v}} : \mathbf{v} \in V^{\geq}\}.$$

In the terminology of polynomial ideals, the monomials in $g(V^{\geq}; \mathbf{z})$ form a truncated ideal generated by the Pareto optima. By the projection theorem (our Theorem 5), we can compute $g(V^{\geq}; \mathbf{z})$ in the form of a polynomial-size rational function in polynomial time.

Step 2. Let $V^{\text{Pareto}} \subseteq \mathbb{Z}^k$ denote the set of Pareto optima. Clearly, we have

$$V^{\text{Pareto}} = (V^{\geq} \setminus (\mathbf{e}_1 + V^{\geq})) \cap \dots \cap (V^{\geq} \setminus (\mathbf{e}_k + V^{\geq})),$$

where $\mathbf{e}_i \in \mathbb{Z}^k$ denotes the i th unit vector and

$$\mathbf{e}_i + V^{\geq} = \{\mathbf{e}_i + \mathbf{v} : \mathbf{v} \in V^{\geq}\}.$$

The generating function $g(V^{\text{Pareto}}; \mathbf{z})$ can be computed by the Boolean operations lemma (Theorem 4) in polynomial time from $g(V^{\geq}; \mathbf{z})$ as

$$\begin{aligned} g(V^{\text{Pareto}}; \mathbf{z}) &= (g(V^{\geq}; \mathbf{z}) - g(V^{\geq}; \mathbf{z}) * z_1 g(V^{\geq}; \mathbf{z})) \\ &\quad * \dots * (g(V^{\geq}; \mathbf{z}) - g(V^{\geq}; \mathbf{z}) * z_k g(V^{\geq}; \mathbf{z})), \end{aligned} \quad (11)$$

where $*$ denotes taking the Hadamard product of the rational functions.

PROOF OF PART (b). Now we recover the Pareto strategies that gave rise to the Pareto optima; i.e., we compute a generating function for the set

$$U^{\text{Pareto}} = \{\mathbf{u} \in \mathbb{Z}^n : \mathbf{u} \in P \cap \mathbb{Z}^n \text{ and } f(\mathbf{u}) \text{ is a Pareto optimum}\}.$$

To this end, we first compute the generating function for the set

$$S^{\text{Pareto}} = \{(\mathbf{u}, \mathbf{v}) \in \mathbb{Z}^n \times \mathbb{Z}^k : \mathbf{v} \text{ is a Pareto point with Pareto strategy } \mathbf{u}\}.$$

For this purpose, we consider the multigraph of the objective functions f_1, \dots, f_k over P ,

$$\begin{aligned} P_{f_1, \dots, f_k}^{\text{=}} &= \{(\mathbf{u}, \mathbf{v}) \in \mathbb{R}^n \times \mathbb{R}^k : \\ &\quad \mathbf{u} \in P, v_i = f_i(\mathbf{u}) \text{ for } i = 1, \dots, k\}. \end{aligned} \quad (12)$$

Using Barvinok's theorem, we can compute in polynomial time the generating function for the integer points in P ,

$$g(P; \mathbf{x}) = \sum \{\mathbf{x}^{\mathbf{u}} : \mathbf{u} \in P \cap \mathbb{Z}^n\},$$

and also, using the monomial substitution $x_j \rightarrow x_j z_1^{f_1(\mathbf{e}_j)} \dots z_k^{f_k(\mathbf{e}_j)}$ for all j , the generating function is transformed into

$$g(P_{f_1, \dots, f_k}^{\text{=}}; \mathbf{x}, \mathbf{z}) = \sum \{\mathbf{x}^{\mathbf{u}} \mathbf{z}^{\mathbf{v}} : (\mathbf{u}, \mathbf{v}) \in P_{f_1, \dots, f_k}^{\text{=}} \cap (\mathbb{Z}^n \times \mathbb{Z}^k)\},$$

where the variables \mathbf{x} carry on the monomial exponents the information of the \mathbf{u} -coordinates of $P_{f_1, \dots, f_k}^{\text{=}}$

and the \mathbf{z} variables of the generating function carry the \mathbf{v} -coordinates of lattice points in $P_{f_1, \dots, f_k}^{\text{=}}$. Now,

$$\begin{aligned} g(S^{\text{Pareto}}; \mathbf{x}, \mathbf{z}) &= (g(P; \mathbf{x}) g(V^{\text{Pareto}}; \mathbf{z})) \\ &\quad * g(P_{f_1, \dots, f_k}^{\text{=}}; \mathbf{x}, \mathbf{z}), \end{aligned} \quad (13)$$

which can be computed in polynomial time for a fixed dimension by the theorems outlined earlier in this section. Finally, to obtain the generating function $g(U^{\text{Pareto}}; \mathbf{x})$ of the Pareto strategies, we need to compute the projection of S^{Pareto} into the space of the strategy variables \mathbf{u} . Because the projection is one-to-one, it suffices to compute the specialization

$$g(U^{\text{Pareto}}; \mathbf{x}) = g(S^{\text{Pareto}}; \mathbf{x}, \mathbf{z} = \mathbf{1}),$$

which can be done in polynomial time.

PROOF OF PART (c). To obtain the number of Pareto optima or Pareto strategies, we first compute the respective rational generating function by the algorithms above and then use Theorem 6 for computing the specialization $g(V^{\text{Pareto}}; \mathbf{z} = \mathbf{1})$ in polynomial time. \square

3. Efficiently Listing All Pareto Optima

The Pareto optimum that corresponds to the best compromise between the individual objective functions is often chosen in an *interactive mode*, where a visualization of the Pareto optima is presented to the user, who then chooses a Pareto optimum. Because the outcome space frequently is of a too-large dimension for visualization, an important task is to list (explicitly enumerate) the elements of the *projection* of the Pareto set into some lower-dimensional linear space.

It is clear that the set of Pareto optima (and thus also any projection) is of exponential size in general, ruling out the existence of a polynomial-time enumeration algorithm. To analyze the running time of an enumeration algorithm, we must turn to *output-sensitive complexity analysis*.

Various notions of output-sensitive efficiency have appeared in the literature; we follow the discussion of Johnson et al. (1988). Let $W \subseteq \mathbb{Z}^p$ be a finite set to be enumerated. An enumeration algorithm is said to run in *polynomial total time* if its running time is bounded by a polynomial in the encoding size of the input and the output. A stronger notion is that of *incremental polynomial time*: Such an algorithm receives a list of solutions $\mathbf{w}_1, \dots, \mathbf{w}_N \in W$ as an additional input. In polynomial time, it outputs one solution $\mathbf{w} \in W \setminus \{\mathbf{w}_1, \dots, \mathbf{w}_N\}$ or asserts that there are no more solutions. An even stronger notion is that of a *polynomial-delay* algorithm, which takes only polynomial time (in the encoding size of the input) before

the first solution is output, between successive outputs of solutions, and after the last solution is output to the termination of the algorithm. Because the algorithm could take exponential time to output all solutions, it could also build exponential-size data structures in the course of the enumeration. This observation gives rise to an even stronger notion of efficiency, a *polynomial-space polynomial-delay* enumeration algorithm.

We also wish to prescribe an *order*, like the lexicographic order, in which the elements are to be enumerated. We consider term orders \prec_R on monomials $\mathbf{y}^{\mathbf{w}}$ that are defined as in Mora and Robbiano (1988) by a nonnegative integral $p \times p$ -matrix R of full rank. Two monomials satisfy $\mathbf{y}^{\mathbf{w}_1} \prec_R \mathbf{y}^{\mathbf{w}_2}$ if and only if $R\mathbf{w}_1$ is lexicographically smaller than $R\mathbf{w}_2$. In other words, if $\mathbf{r}_1, \dots, \mathbf{r}_n$ denote the rows of R , there is some $j \in \{1, \dots, n\}$ such that $\langle \mathbf{r}_j, \mathbf{w}_1 \rangle = \langle \mathbf{r}_j, \mathbf{w}_2 \rangle$ for $i < j$, and $\langle \mathbf{r}_j, \mathbf{w}_1 \rangle < \langle \mathbf{r}_j, \mathbf{w}_2 \rangle$. For example, the unit matrix $R = I_n$ describes the lexicographic term ordering.

We prove the existence of a polynomial-space polynomial-delay prescribed-order enumeration algorithm in a general setting, where the set W to be enumerated is given as the projection of a set presented by a rational generating function.

THEOREM 7. *Let the dimension k and the maximum number l of binomials in the denominator be fixed. Then, there exists a polynomial-space polynomial-delay enumeration algorithm for the following enumeration problem. Given as input, in binary encoding,*

- (I₁) a number $M \in \mathbb{Z}_+$,
- (I₂) rational numbers γ_i , integer vectors $\mathbf{c}_i, \mathbf{d}_{ij}$ for $i \in I, j = 1, \dots, l_i$, where $l_i \leq l$ such that

$$\sum_{i \in I} \gamma_i \frac{\mathbf{z}^{\mathbf{c}_i}}{(1 - \mathbf{z}^{\mathbf{d}_{i1}})(1 - \mathbf{z}^{\mathbf{d}_{i2}}) \dots (1 - \mathbf{z}^{\mathbf{d}_{il_i}})}$$

is a rational generating function of a set $V \subseteq \mathbb{Z}^k$ of lattice points with $V \subseteq [-M, M]^k$,

- (I₃) an integer p with $1 \leq p \leq k$,
- (I₄) a matrix $R \in \mathbb{Z}_+^{p \times p}$,

output, in binary encoding,

(O₁) all points in the projection of V onto the last p components,

$$W = \{\mathbf{w} \in \mathbb{Z}^p : \exists \mathbf{t} \in \mathbb{Z}^{k-p} \text{ such that } (\mathbf{t}, \mathbf{w}) \in V\},$$

in the order \prec_R , which is the term order on monomials in y_1, \dots, y_p induced by the matrix R .

The algorithm can be implemented without using the projection lemma.

We remark that Theorem 7 is a stronger result than what can be obtained by the repeated application of the monomial-extraction technique of Lemma 7 from De Loera et al. (2004), which would only give an incremental polynomial-time enumeration algorithm.

PROOF. We give a simple recursive algorithm that is based on the iterative bisection of intervals.

Input: Lower and upper bound vectors $\mathbf{l}, \mathbf{u} \in \mathbb{Z}^p$.

Output: All vectors \mathbf{w} in W with $\mathbf{l} \leq R\mathbf{w} \leq \mathbf{u}$, sorted in the order \leq_R .

(1) If the set $W \cap \{\mathbf{w} : \mathbf{l} \leq R\mathbf{w} \leq \mathbf{u}\}$ is empty, do nothing.

(2) Otherwise, if $\mathbf{l} = \mathbf{u}$, compute the unique point $\mathbf{w} \in \mathbb{Z}^k$ with $R\mathbf{w} = \mathbf{l} = \mathbf{u}$ and output \mathbf{w} .

(3) Otherwise, let j be the smallest index with $l_j \neq u_j$. We bisect the integer interval $\{l_j, \dots, u_j\}$ evenly into $\{l_j, \dots, m_j\}$ and $\{m_j + 1, \dots, u_j\}$, where $m_j = \lfloor (l_j + u_j)/2 \rfloor$. We invoke the algorithm recursively on the first part, then on the second part, using the corresponding lower and upper bound vectors.

We first need to compute appropriate lower and upper bound vectors \mathbf{l}, \mathbf{u} to start the algorithm. To this end, let N be the largest number in the matrix R , and let $\mathbf{l} = -pMN\mathbf{1}$ and $\mathbf{u} = pMN\mathbf{1}$. Then, $\mathbf{l} \leq R\mathbf{w} \leq \mathbf{u}$ holds for all $\mathbf{w} \in W$. Clearly, the encoding length of \mathbf{l} and \mathbf{u} is bounded polynomially in the input data.

In Step 1 of the algorithm, to determine whether

$$W \cap \{\mathbf{w} : \mathbf{l} \leq R\mathbf{w} \leq \mathbf{u}\} = \emptyset, \tag{14}$$

we consider the polytope

$$Q_{\mathbf{l}, \mathbf{u}} = [-M, M]^{k-p} \times \{\mathbf{w} \in \mathbb{R}^p : \mathbf{l} \leq R\mathbf{w} \leq \mathbf{u}\} \subseteq \mathbb{R}^k, \tag{15}$$

a parallelepiped in \mathbb{R}^k . Because W is the projection of V and because $V \subseteq [-M, M]^k$, we have (14) if and only if $V \cap Q_{\mathbf{l}, \mathbf{u}} = \emptyset$. The rational generating function $g(Q_{\mathbf{l}, \mathbf{u}}; \mathbf{z})$ can be computed in polynomial time. By using the Boolean operations lemma, we can compute the rational generating function $g(V \cap Q_{\mathbf{l}, \mathbf{u}}; \mathbf{z})$ in polynomial time. The specialization $g(V \cap Q_{\mathbf{l}, \mathbf{u}}; \mathbf{z} = \mathbf{1})$ can also be computed in polynomial time by Theorem 6. It gives the number of lattice points in $V \cap Q_{\mathbf{l}, \mathbf{u}}$; in particular, we can decide whether $V \cap Q_{\mathbf{l}, \mathbf{u}} = \emptyset$.

It is clear that the algorithm outputs the elements of W in the order given by \prec_R . We next show that the algorithm is a polynomial-space polynomial-delay enumeration algorithm. The subproblem in Step 1 only depends on the input data as stated in the theorem and on the vectors \mathbf{l} and \mathbf{u} , whose encoding length only decreases in recursive invocations. Therefore, each of the subproblems can be solved in polynomial time (thus also in polynomial space).

The recursion of the algorithm corresponds to a binary tree whose nodes are labeled by the bound vectors \mathbf{l} and \mathbf{u} . There are two types of leaves in the tree, one corresponding to the “empty-box” situation (14) in Step 1, and one corresponding to the “solution-output” situation in Step 2. Inner nodes of the tree correspond to the recursive invocation of the algorithm in Step 3. It is clear that the depth of the recursion is $O(p \log(pMN))$ because the integer intervals are bisected evenly. Thus, the stack space of the

algorithm is polynomially bounded. Because the algorithm does not maintain any global data structures, the whole algorithm uses polynomial space only.

Let $\mathbf{w}_i \in W$ be an arbitrary solution and let \mathbf{w}_{i+1} be its direct successor in the order \prec_R . We shall show that the algorithm only spends polynomial time between the output of \mathbf{w}_i and the output of \mathbf{w}_{i+1} . The key property of the recursion tree of the algorithm is the following:

Every inner node is the root of a subtree that contains at least one solution-output leaf. (16)

The reason for that property is the test for situation (14) in Step 1 of the algorithm. Therefore, the algorithm can visit only $O(p \log(pMN))$ inner nodes and empty-box leaves between the solution-output leaves for \mathbf{w}_i and \mathbf{w}_{i+1} . For the same reason, also the time before the first solution is output and the time after the last solution is output are polynomially bounded. \square

The following corollary, which is a stronger formulation of Theorem 1(d), is immediate. It asserts the existence of an output-sensitive polynomial-time algorithm for the enumeration of arbitrary projections of the set of Pareto optima and strategies.

COROLLARY 8. *Let n and k be fixed integers. Then, there exists a polynomial-space polynomial-delay prescribed-order enumeration algorithm for the following problem. Given as input, in binary encoding,*

(I₁) *an integer m , a matrix $A \in \mathbb{Z}^{m \times n}$, a vector $\mathbf{b} \in \mathbb{Z}^m$, linear functionals $f_1, \dots, f_k \in \mathbb{Z}^n$ (representing a multicriterion integer programming problem),*

(I₂) *an integral matrix $T \in \mathbb{Z}^{p \times (k+n)}$ (representing a projection),*

(I₃) *a matrix $R \in \mathbb{Z}_+^{p \times p}$ (representing a term order),*
 output, in binary encoding,

(O₁) *all vectors in the projection $TS^{\text{Pareto}} \subseteq \mathbb{Z}^p$ of the set S^{Pareto} of Pareto optima and strategies of the multicriterion integer programming problem (1), in the order \prec_R , which is the term order on monomials in y_1, \dots, y_p induced by the matrix R .*

REMARK 9. We remark that Theorem 7 is of general interest. For example, it also implies the existence of a polynomial-space polynomial-delay prescribed-order enumeration algorithm for Hilbert bases of rational polyhedral cones in fixed dimension.

Indeed, fix the dimension d , and let $C = \text{cone}\{\mathbf{b}_1, \dots, \mathbf{b}_n\} \subseteq \mathbb{R}^d$ be a pointed rational polyhedral cone. The Hilbert basis of C is defined as the inclusion-minimal set $H \subseteq C \cap \mathbb{Z}^d$, which generates $C \cap \mathbb{Z}^d$ as a monoid. For simplicial cones C (where $\mathbf{b}_1, \dots, \mathbf{b}_n$ are linearly independent), Barvinok and Woods (2003) proved that one can compute the rational generating function $g(H; \mathbf{z})$ (having a constant number of binomials in the denominators) of the Hilbert basis of $C \cap \mathbb{Z}^d$

using the projection theorem. The same technique works for nonsimplicial pointed cones. Now, Theorem 7 gives a polynomial-space polynomial-delay prescribed-order enumeration algorithm.

4. Selecting a Pareto Optimum Using Global Criteria

Now that we know that all Pareto optima of multicriteria integer linear programs can be encoded in a rational generating function, and that they can be listed efficiently on the output size, we can aim to apply selection criteria stated by a user. The advantage of our setup is that when we optimize a global objective function, it guarantees to return a Pareto optimum because we evaluate the global criterion only on the Pareto optima. Let us start with the simplest global criterion that generalizes the use of the l_1 norm distance function:

THEOREM 10. *Let the dimension k and the maximum number l of binomials in the denominator be fixed.*

(a) *There exists a polynomial-time algorithm for the following problem. Given as input, in binary encoding,*

(I₁) *a bound $M \in \mathbb{Z}_+$,*

(I₂) *rational numbers γ_i , integer vectors \mathbf{c}_i , \mathbf{d}_{ij} for $i \in I$, $j = 1, \dots, l_i$, where $l_i \leq l$ such that*

$$g(V; \mathbf{z}) = \sum_{i \in I} \gamma_i \frac{\mathbf{z}^{\mathbf{c}_i}}{(1 - \mathbf{z}^{\mathbf{d}_{i1}})(1 - \mathbf{z}^{\mathbf{d}_{i2}}) \dots (1 - \mathbf{z}^{\mathbf{d}_{il_i}})}$$

is a rational generating function of a bounded set $V \subseteq \mathbb{Z}^k$ of lattice points with $V \subseteq [-M, M]^k$,

(I₃) *a prescribed point $\hat{\mathbf{v}} \in \mathbb{Z}^k$,*

(I₄) *a vertex description or inequality description of a rational convex central-symmetric polytope $Q \subseteq \mathbb{R}^k$,*
 output, in binary encoding,

(O₁) *a point $\mathbf{v} \in V$ that minimizes the distance $d_Q(\mathbf{v}, \hat{\mathbf{v}}) = \|\mathbf{v} - \hat{\mathbf{v}}\|_Q$ from the prescribed point $\hat{\mathbf{v}}$, where the polyhedral norm $\|\cdot\|_Q$ be defined using the Minkowski functional*

$$\|\mathbf{y}\|_Q = \inf\{\lambda \geq 0: \mathbf{y} \in \lambda Q\}. \quad (17)$$

(b) *There exists a polynomial-space polynomial-delay enumeration algorithm for the following enumeration problem. Given as input, in binary encoding, the same data as above and in addition*

(I₅) *a matrix $R \in \mathbb{Z}_+^{p \times p}$ (representing a term order),*
 output, in binary encoding,

(O₁) *the points of V in the order of increasing distances d_Q from the prescribed point $\hat{\mathbf{v}}$, refined by the term order \prec_R defined by the matrix R .*

Theorem 2, as stated in the introduction, is an immediate corollary of this theorem.

PROOF. Because the dimension k is fixed, we can compute an inequality description

$$Q = \{\mathbf{y} \in \mathbb{R}^k: A\mathbf{y} \leq \mathbf{b}\}$$

of Q with $A \in \mathbb{Z}^{m \times k}$ and $\mathbf{b} \in \mathbb{Z}^k$ in polynomial time, if Q is not already given by an inequality description. Let $\mathbf{v} \in V$ be arbitrary; then

$$\begin{aligned} d_Q(\widehat{\mathbf{v}}, \mathbf{v}) &= \|\mathbf{v} - \widehat{\mathbf{v}}\|_Q \\ &= \inf\{\lambda \geq 0: \mathbf{v} - \widehat{\mathbf{v}} \in \lambda Q\} \\ &= \min\{\lambda \geq 0: \lambda \mathbf{b} \geq A(\mathbf{v} - \widehat{\mathbf{v}})\}. \end{aligned}$$

Thus, there exists an index $i \in \{1, \dots, m\}$ such that

$$d_Q(\widehat{\mathbf{v}}, \mathbf{v}) = \frac{(A\mathbf{v})_i - (A\widehat{\mathbf{v}})_i}{b_i},$$

so $d_Q(\widehat{\mathbf{v}}, \mathbf{v})$ is an integer multiple of $1/b_i$. Hence, for every $\mathbf{v} \in V$, we have that

$$d_Q(\widehat{\mathbf{v}}, \mathbf{v}) \in \frac{1}{\text{lcm}(b_1, \dots, b_m)} \mathbb{Z}_+, \quad (18)$$

where $\text{lcm}(b_1, \dots, b_m)$ clearly is a number of polynomial encoding size. On the other hand, every $\mathbf{v} \in V$ certainly satisfies

$$d_Q(\widehat{\mathbf{v}}, \mathbf{v}) \leq ka(M + \max\{|\widehat{v}_1|, \dots, |\widehat{v}_d|\}), \quad (19)$$

where a is the largest number in A , which is also a bound of polynomial encoding size.

Using Barvinok’s algorithm, we can compute the rational generating function $g(\widehat{\mathbf{v}} + \lambda Q; \mathbf{z})$ for any rational λ of polynomial encoding size in polynomial time. We can also compute the rational generating function $g(V \cap (\widehat{\mathbf{v}} + \lambda Q); \mathbf{z})$ using the Boolean operations lemma. By computing the specialization $g(V \cap (\widehat{\mathbf{v}} + \lambda Q); \mathbf{z} = \mathbf{1})$, we can compute the number of points in $V \cap (\widehat{\mathbf{v}} + \lambda Q)$; thus we can decide whether this set is empty or not.

Hence, we can use binary search for the smallest $\lambda \geq 0$ such that $V \cap (\widehat{\mathbf{v}} + \lambda Q)$ is nonempty. Because of (18) and (19), it runs in polynomial time. By using the recursive bisection algorithm of Theorem 7, it is then possible to construct one Pareto optimum in $V \cap (\widehat{\mathbf{v}} + \lambda Q)$ for part (a), or to construct a sequence of Pareto optima in the desired order for part (b). \square

Now we consider a global criterion using a distance function corresponding to a nonpolyhedral norm like the Euclidean norm $\|\cdot\|_2$ (or any other l_p -norm for $1 < p < \infty$). We are able to prove a very strong type of approximation result, a so-called fully polynomial-time approximation scheme (FPTAS), in a somewhat more general setting.

DEFINITION 11 (FPTAS). Consider the optimization problems

$$\max\{f(\mathbf{v}): \mathbf{v} \in V\}, \quad (20a)$$

$$\min\{f(\mathbf{v}): \mathbf{v} \in V\}. \quad (20b)$$

An FPTAS for the maximization problem (20a) or the minimization problem (20b), respectively, is a family $\{\mathcal{A}_\epsilon: \epsilon \in \mathbb{Q}, \epsilon > 0\}$ of approximation algorithms \mathcal{A}_ϵ , each of which returns an ϵ -approximation; i.e., a solution $\mathbf{v}_\epsilon \in V$ with

$$f(\mathbf{v}_\epsilon) \geq (1 - \epsilon)f^*, \quad \text{where } f^* = \max_{\mathbf{v} \in V} f(\mathbf{v}), \quad (21a)$$

or, respectively,

$$f(\mathbf{v}_\epsilon) \leq (1 + \epsilon)f^*, \quad \text{where } f^* = \min_{\mathbf{v} \in V} f(\mathbf{v}), \quad (21b)$$

such that the algorithms \mathcal{A}_ϵ run in polynomial time in the input size and $1/\epsilon$.

REMARK 12. An FPTAS is based on the notion of ϵ -approximation (21), which gives an approximation guarantee relative to the value f^* of an optimal solution. It is clear that this notion is most useful for objective functions f that are nonnegative on the feasible region V . Because the approximation quality of a solution changes when the objective function is changed by an additive constant, it is not possible in general to convert an FPTAS for a maximization problem to an FPTAS for a minimization problem.

We shall present an FPTAS for the problem of minimizing the distance of a Pareto optimum from a prescribed outcome vector $\widehat{\mathbf{v}} \in \mathbb{Z}^k$. We consider distances $d(\widehat{\mathbf{v}}, \cdot)$ induced by a pseudonorm $\|\cdot\|_Q$ via

$$d(\widehat{\mathbf{v}}, \mathbf{v}) = \|\mathbf{v} - \widehat{\mathbf{v}}\|_Q. \quad (22a)$$

To this end, let $Q \subseteq \mathbb{R}^k$ be a compact basic semialgebraic set with $\mathbf{0} \in \text{int } Q$, which is described by one polynomial inequality,

$$Q = \{\mathbf{y} \in \mathbb{R}^k: q(\mathbf{y}) \leq 1\}, \quad (22b)$$

where $q \in \mathbb{Q}[y_1, \dots, y_k]$ is a homogeneous polynomial of (even) degree D . The pseudonorm $\|\cdot\|_Q$ is now defined using the Minkowski functional:

$$\|\mathbf{y}\|_Q = \inf\{\lambda \geq 0: \mathbf{y} \in \lambda Q\}. \quad (22c)$$

Note that we do not make any assumptions of the convexity of Q , which would make $\|\cdot\|_Q$ a norm. Because Q is compact and $\mathbf{0} \in \text{int } Q$, there exist positive rational numbers (norm equivalence constants) α, β with

$$\alpha B_\infty \subseteq Q \subseteq \beta B_\infty, \quad \text{where } B_\infty = \{\mathbf{y} \in \mathbb{R}^k: \|\mathbf{y}\|_\infty \leq 1\}; \quad (23)$$

see Figure 1.

Now we can formulate our main theorem, which has Theorem 3, which we stated in the introduction, as an immediate corollary.

THEOREM 13. *Let the dimension n and the number k of objective functions be fixed. Moreover, let a degree D and two rational numbers $0 < \alpha \leq \beta$ be fixed. Then, there exists a polynomial-time algorithm for the following problem.*

INFORMS holds copyright to this article and distributed this copy as a courtesy to the author(s). Additional information, including rights and permission policies, is available at <http://journals.informs.org/>.




Figure 1 A Set Defining a Pseudonorm with the Inscribed and Circumscribed Cubes αB_∞ and βB_∞ (Dashed)

Given as input

(I₁) an integer m , a matrix $A \in \mathbb{Z}^{m \times n}$, a vector $\mathbf{b} \in \mathbb{Z}^m$, linear functionals $f_1, \dots, f_k \in \mathbb{Z}^n$, in binary encoding,

(I₂) a prescribed comparison point $\hat{\mathbf{v}} \in \mathbb{Z}^k$, in binary encoding,

(I₃) a list of coefficients $q_i \in \mathbb{Q}$, encoded in binary encoding, and exponent vectors $\alpha_i \in \mathbb{Z}_+$, encoded in unary encoding, representing a homogeneous polynomial

$$q = \sum_i q_i \mathbf{y}^{\alpha_i} \in \mathbb{Q}[y_1, \dots, y_k]$$

of degree D ,

(I₄) a positive rational number $1/\epsilon$ encoded in unary encoding,

output, in binary encoding,

(O₁) a Pareto optimum \mathbf{v}_ϵ of the multicriterion integer programming problem (1) such that

$$\|\mathbf{v}_\epsilon - \hat{\mathbf{v}}\| \leq (1 + \epsilon) \min\{\|\mathbf{v} - \hat{\mathbf{v}}\| : \mathbf{v} \in V^{\text{Pareto}}\},$$

where $\|\cdot\|$ denotes the Euclidean norm and V^{Pareto} is the set of Pareto optima of (1).

The proof is based on the following result, which is a more general formulation of Theorem 1.1 from De Loera et al. (2006).

THEOREM 14 (FPTAS FOR MAXIMIZING NONNEGATIVE POLYNOMIALS OVER FINITE LATTICE POINT SETS). For all fixed integers k (dimension) and l (maximum number of binomials in the denominator), there exists a polynomial-time algorithm for the following problem.

Given as input

(I₁) two vectors $\mathbf{v}_L, \mathbf{v}_U \in \mathbb{Z}^k$,

(I₂) rational numbers γ_i , integer vectors $\mathbf{c}_i, \mathbf{d}_{ij}$ for $i \in I$, $j = 1, \dots, l_i$, where $l_i \leq l$ such that

$$g(V; \mathbf{z}) = \sum_{i \in I} \gamma_i \frac{\mathbf{z}^{\mathbf{c}_i}}{(1 - \mathbf{z}^{\mathbf{d}_{i1}})(1 - \mathbf{z}^{\mathbf{d}_{i2}}) \dots (1 - \mathbf{z}^{\mathbf{d}_{il_i}})}$$

is a rational generating function of a finite set $V \subseteq \mathbb{Z}^k$ of lattice points that is contained in the box $\{\mathbf{v} : \mathbf{v}_L \leq \mathbf{v} \leq \mathbf{v}_U\}$,

(I₃) a list of coefficients $f_i \in \mathbb{Q}$, encoded in binary encoding, and exponent vectors $\alpha_i \in \mathbb{Z}_+$, encoded in unary encoding, representing a polynomial

$$f = \sum_i f_i \mathbf{v}^{\alpha_i} \in \mathbb{Q}[v_1, \dots, v_k]$$

that is nonnegative on V ,

(I₄) a positive rational number $1/\epsilon$ encoded in unary encoding,

output, in binary encoding,

(O₁) a point $\mathbf{v}_\epsilon \in V$ that satisfies

$$f(\mathbf{v}_\epsilon) \geq (1 - \epsilon)f^*, \quad \text{where } f^* = \max_{\mathbf{v} \in V} f(\mathbf{v}).$$

In the paper by De Loera et al. (2006), the above result was stated and proved only for sets V that consist of the lattice points of a rational polytope; however, the same proof yields the result above.

PROOF OF THEOREM 13. Using Theorem 1, we first compute the rational generating function $g(V^{\text{Pareto}}; \mathbf{z})$ of the Pareto optima. With binary search using the Boolean operations lemma with generating functions of cubes as in §3, we can find the smallest nonnegative integer γ such that

$$(\hat{\mathbf{v}} + \gamma B_\infty) \cap V^{\text{Pareto}} \neq \emptyset. \quad (24)$$

If $\gamma = 0$, then the prescribed outcome vector $\hat{\mathbf{v}}$ itself is a Pareto optimum, so it is the optimal solution to the problem.

Otherwise, let \mathbf{v}_0 be an arbitrary outcome vector in $(\hat{\mathbf{v}} + \gamma B_\infty) \cap V^{\text{Pareto}}$. Then,

$$\begin{aligned} \gamma &\geq \|\mathbf{v}_0 - \hat{\mathbf{v}}\|_\infty = \inf\{\lambda : \mathbf{v}_0 - \hat{\mathbf{v}} \in \lambda B_\infty\} \\ &\geq \inf\left\{\lambda : \mathbf{v}_0 - \hat{\mathbf{v}} \in \lambda \frac{1}{\alpha} Q\right\} = \alpha \|\mathbf{v}_0 - \hat{\mathbf{v}}\|_Q, \end{aligned}$$

thus $\|\mathbf{v}_0 - \hat{\mathbf{v}}\|_Q \leq \gamma/\alpha$. Let $\delta = \beta\gamma/\alpha$. Then, for every $\mathbf{v}_1 \in \mathbb{R}^k$ with $\|\mathbf{v}_1 - \hat{\mathbf{v}}\|_\infty \geq \delta$, we have

$$\begin{aligned} \delta &\leq \|\mathbf{v}_1 - \hat{\mathbf{v}}\|_\infty = \inf\{\lambda : \mathbf{v}_1 - \hat{\mathbf{v}} \in \lambda B_\infty\} \\ &\leq \inf\left\{\lambda : \mathbf{v}_1 - \hat{\mathbf{v}} \in \lambda \frac{1}{\beta} Q\right\} = \beta \|\mathbf{v}_1 - \hat{\mathbf{v}}\|_Q; \end{aligned}$$

thus

$$\|\mathbf{v}_1 - \hat{\mathbf{v}}\|_Q \geq \delta/\beta = \gamma/\alpha \geq \|\mathbf{v}_0 - \hat{\mathbf{v}}\|_Q.$$

Therefore, a Pareto optimum $\mathbf{v}^* \in V^{\text{Pareto}}$ minimizing the distance d_Q from the prescribed outcome vector $\hat{\mathbf{v}}$ is contained in the cube $\hat{\mathbf{v}} + \delta B_\infty$. Moreover, for all points $\mathbf{v} \in \hat{\mathbf{v}} + \delta B_\infty$, we have

$$\|\mathbf{v}_0 - \hat{\mathbf{v}}\|_Q \leq \delta/\alpha = \beta\gamma/\alpha^2.$$

We define a function f by

$$f(\mathbf{v}) = (\beta\gamma/\alpha^2)^D - \|\mathbf{v} - \hat{\mathbf{v}}\|_Q^D, \quad (25)$$

which is nonnegative over the cube $\hat{\mathbf{v}} + \delta B_\infty$. Because q is a homogeneous polynomial of degree D , we obtain

$$f(\mathbf{v}) = (\beta\gamma/\alpha^2)^D - q(\mathbf{v} - \hat{\mathbf{v}}), \quad (26)$$

so f is a polynomial.

We next compute the rational generating function

$$g(V^{\text{Pareto}} \cap (\hat{\mathbf{v}} + \delta B_\infty); \mathbf{z})$$

from $g(V^{\text{Pareto}}; \mathbf{z})$ using the Boolean operations lemma. Let $\epsilon' > 0$ be a rational number, which we will determine later. By Theorem 14, we compute a solution $\mathbf{v}_{\epsilon'} \in V^{\text{Pareto}}$ with

$$f(\mathbf{v}_{\epsilon'}) \geq (1 - \epsilon')f(\mathbf{v}^*),$$

or, equivalently,

$$f(\mathbf{v}^*) - f(\mathbf{v}_{\epsilon'}) \leq \epsilon' f(\mathbf{v}^*).$$

Thus,

$$\begin{aligned} [d_Q(\hat{\mathbf{v}}, \mathbf{v}_{\epsilon'})]^D - [d_Q(\hat{\mathbf{v}}, \mathbf{v}^*)]^D &= \|\mathbf{v}_{\epsilon'} - \hat{\mathbf{v}}\|_Q^D - \|\mathbf{v}^* - \hat{\mathbf{v}}\|_Q^D \\ &= f(\mathbf{v}^*) - f(\mathbf{v}_{\epsilon'}) \\ &\leq \epsilon' f(\mathbf{v}^*) \\ &= \epsilon' ((\beta\gamma/\alpha^2)^D - \|\mathbf{v}^* - \hat{\mathbf{v}}\|_Q^D). \end{aligned}$$

Because γ is the smallest integer with (24) and also $\|\mathbf{v}^* - \hat{\mathbf{v}}\|_\infty$ is an integer, we have

$$\gamma \leq \|\mathbf{v}^* - \hat{\mathbf{v}}\|_\infty \leq \beta \|\mathbf{v}^* - \hat{\mathbf{v}}\|_Q.$$

Thus,

$$[d_Q(\hat{\mathbf{v}}, \mathbf{v}_{\epsilon'})]^D - [d_Q(\hat{\mathbf{v}}, \mathbf{v}^*)]^D \leq \epsilon' \left[\left(\frac{\beta}{\alpha} \right)^{2D} - 1 \right] \|\mathbf{v}^* - \hat{\mathbf{v}}\|_Q^D.$$

An elementary calculation yields

$$d_Q(\hat{\mathbf{v}}, \mathbf{v}_{\epsilon'}) - d_Q(\hat{\mathbf{v}}, \mathbf{v}^*) \leq \frac{\epsilon'}{D} \left[\left(\frac{\beta}{\alpha} \right)^{2D} - 1 \right] d_Q(\hat{\mathbf{v}}, \mathbf{v}^*).$$

Thus, we can choose

$$\epsilon' = \epsilon D \left[\left(\frac{\beta}{\alpha} \right)^{2D} - 1 \right]^{-1} \tag{27}$$

to get the desired estimate. Because α , β , and D are fixed constants, we have $\epsilon' = \Theta(\epsilon)$. Thus, the computation of $\mathbf{v}_{\epsilon'} \in V^{\text{Pareto}}$ by Theorem 14 runs in polynomial time in the input encoding size and $1/\epsilon$. \square

REMARK 15. It is straightforward to extend this result to also include the l_p norms for odd integers p , by solving the approximation problem separately for all of the $2^k = O(1)$ shifted orthants $\hat{\mathbf{v}} + O_\sigma = \{\mathbf{v}: \sigma_i(v_i - \hat{v}_i) \geq 0\}$, where $\sigma \in \{\pm 1\}^k$. On each of the orthants, the l_p -norm has a representation by a polynomial as required by Theorem 13.

Acknowledgments

The first author gratefully acknowledges support from National Science Foundation Grant DMS-0608785. The third author was supported by a Feodor Lynen Research Fellowship from the Alexander von Humboldt Foundation.

Note Added in Proof

It has come to our attention that Blanco (2007) has independently investigated rational generating function techniques for multicriteria integer linear programs for his Ph.D. work.

References

Barvinok, A. I. 1994. Polynomial time algorithm for counting integral points in polyhedra when the dimension is fixed. *Math. Oper. Res.* **19** 769–779.

Barvinok, A. I., J. E. Pommersheim. 1999. An algorithmic theory of lattice points in polyhedra. L. J. Billera, A. Björner, C. Greene, R. E. Simion, R. P. Stanley, eds. *New Perspectives in Algebraic Combinatorics, Mathematical Sciences Research Institute Publications*, Vol. 38. Cambridge University Press, Cambridge, UK, 91–147.

Barvinok, A. I., K. Woods. 2003. Short rational generating functions for lattice point problems. *J. Amer. Math. Soc.* **16** 957–979.

Blanco, V. 2007. Personal communication, September 3.

De Loera, J. A., R. Hemmecke, M. Köppe, R. Weismantel. 2006. Integer polynomial optimization in fixed dimension. *Math. Oper. Res.* **31** 147–153.

De Loera, J. A., D. Haws, R. Hemmecke, P. Huggins, B. Sturmfels, R. Yoshida. 2004. Short rational functions for toric algebra and applications. *J. Symbolic Comput.* **38** 959–973.

Ehrgott, M., X. Gandibleux. 2000. A survey and annotated bibliography of multiobjective combinatorial optimization. *OR Spektrum* **22** 425–460.

Emelichev, V. A., V. A. Perepelitsa. 1992. On the cardinality of the set of alternatives in discrete many-criterion problems. *Discrete Math. Appl.* **2** 461–471.

Figueira, J., S. Greco, M. Ehrgott, eds. 2005. *Multiple Criteria Decision Analysis. State of the Art Surveys*. Springer, New York.

Isermann, H. 1974. Proper efficiency and the linear vector maximum problem. *Oper. Res.* **22** 189–191.

Johnson, D. S., M. Yannakakis, C. H. Papadimitriou. 1988. On generating all maximal independent sets. *Inform. Processing Lett.* **27** 119–123.

Mora, T., L. Robbiano. 1988. The Gröbner fan of an ideal. *J. Symbolic Comput.* **6** 183–208.

Sawaragi, Y., H. Nakayama, T. Tanino, eds. 1985. *Theory of Multiobjective Optimization*. Academic Press, Orlando, FL.

Sergienko, I. V., V. A. Perepelitsa. 1991. Finding the set of alternatives in discrete multi-criterion problems. *Cybernetics* **3** 673–683.

Woods, K. 2004. Rational generating functions and lattice point sets. Ph.D. thesis, University of Michigan, Ann Arbor.

INFORMS holds copyright to this article and distributed this copy as a courtesy to the author(s). Additional information, including rights and permission policies, is available at http://journals.informs.org/.