

## MAT 22B Problem Set 6 (Due 9/2)

1. Find the solution of the initial value problem

$$u'' + u = F(t), \quad u(0) = 0, \quad u'(0) = 0,$$

where

$$F(t) = \begin{cases} F_0 t, & 0 \leq t \leq \pi \\ F_0(2\pi - t), & \pi < t \leq 2\pi, \\ 0, & 2\pi < t. \end{cases}$$

2. Compute the Laplace transform of the following functions

$$(a) \quad f(t) = \begin{cases} 1 & 0 \leq t < \pi \\ 0 & \pi \leq t < \infty \end{cases}$$

$$(b) \quad f(t) = \begin{cases} t & 0 \leq t < 1 \\ 2-t & 1 \leq t < 2 \\ 0 & 2 \leq t < \infty \end{cases}$$

$$(c) \quad f(t) = t^n e^{at}$$

3. Compute the inverse Laplace transform of the following functions

$$(a) \quad F(s) = \frac{4}{(s-1)^3}$$

$$(b) \quad F(s) = \frac{2s+2}{s^2+2s+5}$$

$$(c) \quad F(s) = \frac{8s^2-4s+12}{s(s^2+4)}$$

4. Use the Laplace Transform to solve the following initial value problems

$$(a) \quad y'' - y' - 6y = 0, \quad y(0) = 1, \quad y'(0) = -1$$

$$(b) \quad y'' - 2y' + 2y = 0, \quad y(0) = 0, \quad y'(0) = 1$$

$$(c) \quad y^{(4)} - y = 0, \quad y(0) = 1, \quad y'(0) = 0, \quad y''(0) = 1, \quad y'''(0) = 0$$

$$(d) \quad y'' - 2y' + 2y = e^{-t}, \quad y(0) = 0, \quad y'(0) = 1$$

5. Transform the following differential equations into a system of first-order differential equations. If initial values are given, also transform the initial values of the original equation to appropriate initial conditions for the system of equations.

$$(a) \quad t^2 u'' + tu' + (t^2 - 0.25)u = 0$$

$$(b) \quad u^{(4)} - u = 0$$

$$(c) \quad u'' + 0.25u' + 4u = 2\cos(3t), \quad u(0) = 1, \quad u'(0) = -2$$

$$(d) \quad u'' + p(t)u' + q(t)u = g(t), \quad u(0) = u_0, \quad u'(0) = u'_0$$