

CHAPTER 2

Lebesgue Measure on \mathbb{R}^n

Our goal is to construct a notion of the volume, or Lebesgue measure, of rather general subsets of \mathbb{R}^n that reduces to the usual volume of elementary geometrical sets such as cubes or rectangles.

If $\mathcal{L}(\mathbb{R}^n)$ denotes the collection of Lebesgue measurable sets and

$$\mu : \mathcal{L}(\mathbb{R}^n) \rightarrow [0, \infty]$$

denotes Lebesgue measure, then we want $\mathcal{L}(\mathbb{R}^n)$ to contain all n -dimensional rectangles and $\mu(R)$ should be the usual volume of a rectangle R . Moreover, we want μ to be countably additive. That is, if

$$\{A_i \in \mathcal{L}(\mathbb{R}^n) : i \in \mathbb{N}\}$$

is a countable collection of disjoint measurable sets, then their union should be measurable and

$$\mu\left(\bigcup_{i=1}^{\infty} A_i\right) = \sum_{i=1}^{\infty} \mu(A_i).$$

The reason for requiring countable additivity is that finite additivity is too weak a property to allow the justification of any limiting processes, while uncountable additivity is too strong; for example, it would imply that if the measure of a set consisting of a single point is zero, then the measure of every subset of \mathbb{R}^n would be zero.

It is not possible to define the Lebesgue measure of all subsets of \mathbb{R}^n in a geometrically reasonable way. For example, Banach and Tarski (1924) showed that if $n \geq 3$, one can cut up a ball into a finite number of pieces and use isometries (translations and rotations) to reassemble the pieces into a ball of any desired volume *e.g.* reassemble a pea into the sun. This process requires the axiom of choice,¹ and the moral of the Banach-Tarski ‘paradox’ is that some subsets of \mathbb{R}^n are too irregular to define their Lebesgue measure in a way that preserves additivity together with the invariance of the measure under isometries.²

There are many ways to construct Lebesgue measure, all of which lead to the same result. We will follow an approach due to Carathéodory (which generalizes to other measures): We first construct an outer measure on all subsets of \mathbb{R}^n by approximating them from the outside by countable unions of rectangles; we then

¹Solovay (1970) proved that one has to use the axiom of choice to obtain non-Lebesgue measurable sets.

²Banach showed that if $n = 1$ or $n = 2$ there are finitely additive, isometrically invariant extensions of Lebesgue measure on \mathbb{R}^n that are defined on all subsets of \mathbb{R}^n ; but these extensions are not countably additive. We will not consider any possible extensions of $\mathcal{L}(\mathbb{R}^n)$ since the σ -algebra of Lebesgue measurable sets is already large enough to include all set of ‘practical’ importance in analysis.

restrict this outer measure to a σ -algebra of measurable subsets on which it is countably additive. This approach is somewhat asymmetrical in that we approximate sets (and their complements) from the outside by elementary sets, but we do not approximate them directly from the inside.

See Jones [5], Stein and Shakarchi [6], and Wheeler and Zygmund [8] for detailed introductions to Lebesgue measure on \mathbb{R}^n . Cohn [2] gives a similar development to the one here, and Evans and Gariepy [3] discusses many more advanced topics.

2.1. Lebesgue outer measure

We use rectangles as our elementary sets, defined as follows.

Definition 2.1. An n -dimensional, closed rectangle with sides oriented parallel to the coordinate axes, or rectangle for short, is a subset $R \subset \mathbb{R}^n$ of the form

$$R = [a_1, b_1] \times [a_2, b_2] \times \cdots \times [a_n, b_n]$$

where $-\infty < a_i \leq b_i < \infty$ for $i = 1, \dots, n$. The volume $\mu(R)$ of R is

$$\mu(R) = (b_1 - a_1)(b_2 - a_2) \cdots (b_n - a_n).$$

If $n = 1$ or $n = 2$, the volume of a rectangle is its length or area, respectively. We also consider the empty set to be a rectangle with $\mu(\emptyset) = 0$. We denote the collection of all n -dimensional rectangles by $\mathcal{R}(\mathbb{R}^n)$, or \mathcal{R} when n is understood, and then $R \mapsto \mu(R)$ defines a map

$$\mu : \mathcal{R}(\mathbb{R}^n) \rightarrow [0, \infty).$$

The use of this particular class of elementary sets is for convenience. We could equally well use open or half-open rectangles, cubes, balls, or other suitable elementary sets; the result would be the same.

Definition 2.2. The outer Lebesgue measure $\mu^*(E)$ of a subset $E \subset \mathbb{R}^n$, or outer measure for short, is

$$(2.1) \quad \mu^*(E) = \inf \left\{ \sum_{i=1}^{\infty} \mu(R_i) : E \subset \bigcup_{i=1}^{\infty} R_i, R_i \in \mathcal{R}(\mathbb{R}^n) \right\}$$

where the infimum is taken over all countable collections of rectangles whose union contains E . The map

$$\mu^* : \mathcal{P}(\mathbb{R}^n) \rightarrow [0, \infty], \quad \mu^* : E \mapsto \mu^*(E)$$

is called outer Lebesgue measure.

In this definition, a sum $\sum_{i=1}^{\infty} \mu(R_i)$ and $\mu^*(E)$ may take the value ∞ . We do not require that the rectangles R_i are disjoint, so the same volume may contribute to multiple terms in the sum on the right-hand side of (2.1); this does not affect the value of the infimum.

Example 2.3. Let $E = \mathbb{Q} \cap [0, 1]$ be the set of rational numbers between 0 and 1. Then E has outer measure zero. To prove this, let $\{q_i : i \in \mathbb{N}\}$ be an enumeration of the points in E . Given $\epsilon > 0$, let R_i be an interval of length $\epsilon/2^i$ which contains q_i . Then $E \subset \bigcup_{i=1}^{\infty} R_i$ so

$$0 \leq \mu^*(E) \leq \sum_{i=1}^{\infty} \mu(R_i) = \epsilon.$$

Hence $\mu^*(E) = 0$ since $\epsilon > 0$ is arbitrary. The same argument shows that any countable set has outer measure zero. Note that if we cover E by a *finite* collection of intervals, then the union of the intervals would have to contain $[0, 1]$ since E is dense in $[0, 1]$ so their lengths sum to at least one.

The previous example illustrates why we need to use countably infinite collections of rectangles, not just finite collections, to define the outer measure.³ The ‘countable ϵ -trick’ used in the example appears in various forms throughout measure theory.

Next, we prove that μ^* is an outer measure in the sense of Definition 1.1.

Theorem 2.4. *Lebesgue outer measure μ^* has the following properties.*

- (a) $\mu^*(\emptyset) = 0$;
- (b) if $E \subset F$, then $\mu^*(E) \leq \mu^*(F)$;
- (c) if $\{E_i \subset \mathbb{R}^n : i \in \mathbb{N}\}$ is a countable collection of subsets of \mathbb{R}^n , then

$$\mu^*\left(\bigcup_{i=1}^{\infty} E_i\right) \leq \sum_{i=1}^{\infty} \mu^*(E_i).$$

PROOF. It follows immediately from Definition 2.2 that $\mu^*(\emptyset) = 0$, since every collection of rectangles covers \emptyset , and that $\mu^*(E) \leq \mu^*(F)$ if $E \subset F$, since any cover of F covers E .

The main property to prove is the countable subadditivity of μ^* . If $\mu^*(E_i) = \infty$ for some $i \in \mathbb{N}$, there is nothing to prove, so we may assume that $\mu^*(E_i)$ is finite for every $i \in \mathbb{N}$. If $\epsilon > 0$, there is a countable covering $\{R_{ij} : j \in \mathbb{N}\}$ of E_i by rectangles R_{ij} such that

$$\sum_{j=1}^{\infty} \mu(R_{ij}) \leq \mu^*(E_i) + \frac{\epsilon}{2^i}, \quad E_i \subset \bigcup_{j=1}^{\infty} R_{ij}.$$

Then $\{R_{ij} : i, j \in \mathbb{N}\}$ is a countable covering of

$$E = \bigcup_{i=1}^{\infty} E_i$$

and therefore

$$\mu^*(E) \leq \sum_{i,j=1}^{\infty} \mu(R_{ij}) \leq \sum_{i=1}^{\infty} \left\{ \mu^*(E_i) + \frac{\epsilon}{2^i} \right\} = \sum_{i=1}^{\infty} \mu^*(E_i) + \epsilon.$$

Since $\epsilon > 0$ is arbitrary, it follows that

$$\mu^*(E) \leq \sum_{i=1}^{\infty} \mu^*(E_i)$$

which proves the result. □

³The use of finitely many intervals leads to the notion of the Jordan content of a set, introduced by Peano (1887) and Jordan (1892), which is closely related to the Riemann integral; Borel (1898) and Lebesgue (1902) generalized Jordan’s approach to allow for countably many intervals, leading to Lebesgue measure and the Lebesgue integral.

2.2. The outer measure of a rectangle

In this section, we prove the geometrically obvious, but not entirely trivial, fact that the outer measure of a rectangle is equal to its volume. The main point is to show that the volumes of a countable collection of rectangles that cover a rectangle R cannot sum to less than the volume of R .⁴

We begin with some combinatorial facts about finite covers of rectangles [6]. We denote the interior of a rectangle R by R° , and we say that rectangles R, S are almost disjoint if $R^\circ \cap S^\circ = \emptyset$, meaning that they intersect at most along their boundaries. The proofs of the following results are cumbersome to write out in detail (it's easier to draw a picture) but we briefly explain the argument.

Lemma 2.5. *Suppose that*

$$R = I_1 \times I_2 \times \cdots \times I_n$$

is an n -dimensional rectangle, and each closed, bounded interval $I_i \subset \mathbb{R}$ is an almost disjoint union of closed, bounded intervals $\{I_{i,j} \subset \mathbb{R} : j = 1, \dots, N_i\}$,

$$I_i = \bigcup_{j=1}^{N_i} I_{i,j}.$$

Define the rectangles

$$(2.2) \quad S_{j_1 j_2 \dots j_n} = I_{1,j_1} \times I_{1,j_2} \times \cdots \times I_{j_n}.$$

Then

$$\mu(R) = \sum_{j_1=1}^{N_1} \cdots \sum_{j_n=1}^{N_n} \mu(S_{j_1 j_2 \dots j_n}).$$

PROOF. Denoting the length of an interval I by $|I|$, using the fact that

$$|I_i| = \sum_{j=1}^{N_i} |I_{i,j}|,$$

and expanding the resulting product, we get that

$$\begin{aligned} \mu(R) &= |I_1| |I_2| \cdots |I_n| \\ &= \left(\sum_{j_1=1}^{N_1} |I_{1,j_1}| \right) \left(\sum_{j_2=1}^{N_2} |I_{2,j_2}| \right) \cdots \left(\sum_{j_n=1}^{N_n} |I_{n,j_n}| \right) \\ &= \sum_{j_1=1}^{N_1} \sum_{j_2=1}^{N_2} \cdots \sum_{j_n=1}^{N_n} |I_{1,j_1}| |I_{2,j_2}| \cdots |I_{n,j_n}| \\ &= \sum_{j_1=1}^{N_1} \sum_{j_2=1}^{N_2} \cdots \sum_{j_n=1}^{N_n} \mu(S_{j_1 j_2 \dots j_n}). \end{aligned}$$

□

⁴As a partial justification of the need to prove this fact, note that it would not be true if we allowed uncountable covers, since we could cover any rectangle by an uncountable collection of points all of whose volumes are zero.

Proposition 2.6. *If a rectangle R is an almost disjoint, finite union of rectangles $\{R_1, R_2, \dots, R_N\}$, then*

$$(2.3) \quad \mu(R) = \sum_{i=1}^N \mu(R_i).$$

If R is covered by rectangles $\{R_1, R_2, \dots, R_N\}$, which need not be disjoint, then

$$(2.4) \quad \mu(R) \leq \sum_{i=1}^N \mu(R_i).$$

PROOF. Suppose that

$$R = [a_1, b_1] \times [a_2, b_2] \times \cdots \times [a_n, b_n]$$

is an almost disjoint union of the rectangles $\{R_1, R_2, \dots, R_N\}$. Then by ‘extending the sides’ of the R_i , we may decompose R into an almost disjoint collection of rectangles

$$\{S_{j_1 j_2 \dots j_n} : 1 \leq j_i \leq N_i \text{ for } 1 \leq i \leq n\}$$

that is obtained by taking products of subintervals of partitions of the coordinate intervals $[a_i, b_i]$ into unions of almost disjoint, closed subintervals. Explicitly, we partition $[a_i, b_i]$ into

$$a_i = c_{i,0} \leq c_{i,1} \leq \cdots \leq c_{i,N_i} = b_i, \quad I_{i,j} = [c_{i,j-1}, c_{i,j}].$$

where the $c_{i,j}$ are obtained by ordering the left and right i th coordinates of all faces of rectangles in the collection $\{R_1, R_2, \dots, R_N\}$, and define rectangles $S_{j_1 j_2 \dots j_n}$ as in (2.2).

Each rectangle R_i in the collection is an almost disjoint union of rectangles $S_{j_1 j_2 \dots j_n}$, and their union contains all such products exactly once, so by applying Lemma 2.5 to each R_i and summing the results we see that

$$\sum_{i=1}^N \mu(R_i) = \sum_{j_1=1}^{N_1} \cdots \sum_{j_n=1}^{N_n} \mu(S_{j_1 j_2 \dots j_n}).$$

Similarly, R is an almost disjoint union of all the rectangles $S_{j_1 j_2 \dots j_n}$, so Lemma 2.5 implies that

$$\mu(R) = \sum_{j_1=1}^{N_1} \cdots \sum_{j_n=1}^{N_n} \mu(S_{j_1 j_2 \dots j_n}),$$

and (2.3) follows.

If a finite collection of rectangles $\{R_1, R_2, \dots, R_N\}$ covers R , then there is a almost disjoint, finite collection of rectangles $\{S_1, S_2, \dots, S_M\}$ such that

$$R = \bigcup_{i=1}^M S_i, \quad \sum_{i=1}^M \mu(S_i) \leq \sum_{i=1}^N \mu(R_i).$$

To obtain the S_i , we replace R_i by the rectangle $R \cap R_i$, and then decompose these possibly non-disjoint rectangles into an almost disjoint, finite collection of sub-rectangles with the same union; we discard ‘overlaps’ which can only reduce the sum of the volumes. Then, using (2.3), we get

$$\mu(R) = \sum_{i=1}^M \mu(S_i) \leq \sum_{i=1}^N \mu(R_i),$$

which proves (2.4). \square

The outer measure of a rectangle is defined in terms of countable covers. We reduce these to finite covers by using the topological properties of \mathbb{R}^n .

Proposition 2.7. *If R is a rectangle in \mathbb{R}^n , then $\mu^*(R) = \mu(R)$.*

PROOF. Since $\{R\}$ covers R , we have $\mu^*(R) \leq \mu(R)$, so we only need to prove the reverse inequality.

Suppose that $\{R_i : i \in \mathbb{N}\}$ is a countably infinite collection of rectangles that covers R . By enlarging R_i slightly we may obtain a rectangle S_i whose interior S_i° contains R_i such that

$$\mu(S_i) \leq \mu(R_i) + \frac{\epsilon}{2^i}.$$

Then $\{S_i^\circ : i \in \mathbb{N}\}$ is an open cover of the compact set R , so it contains a finite subcover, which we may label as $\{S_1^\circ, S_2^\circ, \dots, S_N^\circ\}$. Then $\{S_1, S_2, \dots, S_N\}$ covers R and, using (2.4), we find that

$$\mu(R) \leq \sum_{i=1}^N \mu(S_i) \leq \sum_{i=1}^N \left\{ \mu(R_i) + \frac{\epsilon}{2^i} \right\} \leq \sum_{i=1}^{\infty} \mu(R_i) + \epsilon.$$

Since $\epsilon > 0$ is arbitrary, we have

$$\mu(R) \leq \sum_{i=1}^{\infty} \mu(R_i)$$

and it follows that $\mu(R) \leq \mu^*(R)$. \square

2.3. Measurable sets

The following is the Carathéodory definition of measurability.

Definition 2.8. A subset $A \subset \mathbb{R}^n$ is Lebesgue measurable if

$$(2.5) \quad \mu^*(E) = \mu^*(E \cap A) + \mu^*(E \cap A^c)$$

for every subset $E \subset \mathbb{R}^n$. We denote the collection of Lebesgue measurable sets in \mathbb{R}^n by $\mathcal{L}(\mathbb{R}^n)$.

We also write $E \cap A^c$ as $E \setminus A$. Thus, a measurable set A splits any set E into disjoint pieces whose outer measures add up to the outer measure of E . Heuristically, this condition means that a set is measurable if it divides other sets in a ‘nice’ way. The regularity of E is not important here; for example, if (2.5) holds whenever E is a rectangle, then it holds for arbitrary sets E .

Since μ^* is subadditive, we always have

$$\mu^*(E) \leq \mu^*(E \cap A) + \mu^*(E \cap A^c).$$

Thus, to prove that $A \subset \mathbb{R}^n$ is measurable, it is sufficient to show that

$$\mu^*(E) \geq \mu^*(E \cap A) + \mu^*(E \cap A^c)$$

for every $E \subset \mathbb{R}^n$, and then we have equality as in (2.5).

Definition 2.8 is perhaps not the most intuitive way to define the measurability of sets, but it leads directly to the following key result. The same result, with the same proof, applies to any outer measure.

Theorem 2.9. *The collection $\mathcal{L}(\mathbb{R}^n)$ of Lebesgue measurable sets is a σ -algebra on \mathbb{R}^n , and the restriction of Lebesgue outer measure μ^* to $\mathcal{L}(\mathbb{R}^n)$ is a measure on $\mathcal{L}(\mathbb{R}^n)$.*

PROOF. It follows immediately from (2.5) that \emptyset is measurable and the complement of a measurable set is measurable, so to prove that $\mathcal{L}(\mathbb{R}^n)$ is a σ -algebra, we only need to show that it is closed under countable unions. We will prove at the same time that μ^* is countably additive on $\mathcal{L}(\mathbb{R}^n)$; since $\mu^*(\emptyset) = 0$, this will prove that μ^* is a measure on $\mathcal{L}(\mathbb{R}^n)$.

First, we prove that the union of measurable sets is measurable. Suppose that $A, B \in \mathcal{L}(\mathbb{R}^n)$ and $E \subset \mathbb{R}^n$. The measurability of A and B implies that

$$\begin{aligned} \mu^*(E) &= \mu^*(E \cap A) + \mu^*(E \cap A^c) \\ (2.6) \quad &= \mu^*(E \cap A \cap B) + \mu^*(E \cap A \cap B^c) \\ &\quad + \mu^*(E \cap A^c \cap B) + \mu^*(E \cap A^c \cap B^c). \end{aligned}$$

Since $A \cup B = (A \cap B) \cup (A \cap B^c) \cup (A^c \cap B)$ and μ^* is subadditive, we have

$$\mu^*(E \cap (A \cup B)) \leq \mu^*(E \cap A \cap B) + \mu^*(E \cap A \cap B^c) + \mu^*(E \cap A^c \cap B).$$

The use of this inequality and the relation $A^c \cap B^c = (A \cup B)^c$ in (2.6) implies that

$$\mu^*(E) \geq \mu^*(E \cap (A \cup B)) + \mu^*(E \cap (A \cup B)^c)$$

so $A \cup B$ is measurable.

Moreover, if A is measurable and $A \cap B = \emptyset$, then by taking $E = A \cup B$ in (2.5), we see that

$$\mu^*(A \cup B) = \mu^*(A) + \mu^*(B).$$

Thus, the outer measure of the union of disjoint, measurable sets is the sum of their outer measures. The repeated application of this result implies that the finite union of measurable sets is measurable and μ^* is finitely additive on $\mathcal{L}(\mathbb{R}^n)$.

Next, we want to show that the countable union of measurable sets is measurable. It is sufficient to consider disjoint unions. To see this, note that if

$$\{A_i \in \mathcal{L}(\mathbb{R}^n) : i \in \mathbb{N}\}.$$

is a countably infinite collection of measurable sets, then

$$B_j = \bigcup_{i=1}^j A_i, \quad \text{for } j \geq 1$$

form an increasing sequence of measurable sets, and

$$C_j = B_j \setminus B_{j-1} \quad \text{for } j \geq 2, \quad C_1 = B_1$$

form a disjoint measurable collection of sets. Moreover

$$\bigcup_{i=1}^{\infty} A_i = \bigcup_{j=1}^{\infty} C_j.$$

Suppose that $\{A_i \in \mathcal{L}(\mathbb{R}^n) : i \in \mathbb{N}\}$ is a countably infinite, disjoint collection of measurable sets, and define

$$B_j = \bigcup_{i=1}^j A_i, \quad B = \bigcup_{i=1}^{\infty} A_i.$$

Let $E \subset \mathbb{R}^n$. Since A_j is measurable and $B_j = A_j \cup B_{j-1}$ is a disjoint union (for $j \geq 2$),

$$\begin{aligned}\mu^*(E \cap B_j) &= \mu^*(E \cap B_j \cap A_j) + \mu^*(E \cap B_j \cap A_j^c), \\ &= \mu^*(E \cap A_j) + \mu^*(E \cap B_{j-1}).\end{aligned}$$

Also $\mu^*(E \cap B_1) = \mu^*(E \cap A_1)$. It follows by induction that

$$\mu^*(E \cap B_j) = \sum_{i=1}^j \mu^*(E \cap A_i).$$

Since B_j is a finite union of measurable sets, it is measurable, so

$$\mu^*(E) = \mu^*(E \cap B_j) + \mu^*(E \cap B_j^c),$$

and since $B_j^c \supset B^c$, we have

$$\mu^*(E \cap B_j^c) \geq \mu^*(E \cap B^c).$$

It follows that

$$\mu^*(E) \geq \sum_{i=1}^j \mu^*(E \cap A_i) + \mu^*(E \cap B^c).$$

Taking the limit of this inequality as $j \rightarrow \infty$ and using the subadditivity of μ^* , we get

$$\begin{aligned}(2.7) \quad \mu^*(E) &\geq \sum_{i=1}^{\infty} \mu^*(E \cap A_i) + \mu^*(E \cap B^c) \\ &\geq \mu^*\left(\bigcup_{i=1}^{\infty} E \cap A_i\right) + \mu^*(E \cap B^c) \\ &\geq \mu^*(E \cap B) + \mu^*(E \cap B^c) \\ &\geq \mu^*(E).\end{aligned}$$

Therefore, we must have equality in (2.7), which shows that $B = \bigcup_{i=1}^{\infty} A_i$ is measurable. Moreover,

$$\mu^*\left(\bigcup_{i=1}^{\infty} E \cap A_i\right) = \sum_{i=1}^{\infty} \mu^*(E \cap A_i),$$

so taking $E = \mathbb{R}^n$, we see that μ^* is countably additive on $\mathcal{L}(\mathbb{R}^n)$. \square

The preceding theorem shows that we get a measure in the sense of Definition 1.5 by restricting an outer measure to measurable sets.

Definition 2.10. Lebesgue measure

$$\mu : \mathcal{L}(\mathbb{R}^n) \rightarrow [0, \infty], \quad \mu = \mu^*|_{\mathcal{L}(\mathbb{R}^n)}$$

is the restriction of Lebesgue outer measure μ^* to the Lebesgue measurable sets $\mathcal{L}(\mathbb{R}^n)$.

From Proposition 2.7, this notation is consistent with our previous use of μ to denote the volume of a rectangle.

Next, we prove that all rectangles are measurable; this implies that $\mathcal{L}(\mathbb{R}^n)$ is a ‘large’ collection of subsets of \mathbb{R}^n . Not all subsets of \mathbb{R}^n are Lebesgue measurable, however; *e.g.* see Example 2.17 below.

Proposition 2.11. *Every rectangle is Lebesgue measurable.*

PROOF. Let R be an n -dimensional rectangle and $E \subset \mathbb{R}^n$. Given $\epsilon > 0$, there is a cover $\{R_i : i \in \mathbb{N}\}$ of E by rectangles R_i such that

$$\mu^*(E) + \epsilon \geq \sum_{i=1}^{\infty} \mu(R_i).$$

We can decompose R_i into an almost disjoint, finite union of rectangles

$$\{\tilde{R}_i, S_{i,1}, \dots, S_{i,N}\}$$

such that

$$R_i = \tilde{R}_i + \bigcup_{j=1}^N S_{i,j}, \quad \tilde{R}_i = R_i \cap R \subset R, \quad S_{i,j} \subset \overline{R^c}.$$

From (2.3),

$$\mu(R_i) = \mu(\tilde{R}_i) + \sum_{j=1}^N \mu(S_{i,j}).$$

Using this result in the previous sum, relabeling the $S_{i,j}$ as S_i , and rearranging the resulting sum, we get that

$$\mu^*(E) + \epsilon \geq \sum_{i=1}^{\infty} \mu(\tilde{R}_i) + \sum_{i=1}^{\infty} \mu(S_i).$$

Since the rectangles $\{\tilde{R}_i : i \in \mathbb{N}\}$ cover $E \cap R$ and the rectangles $\{S_i : i \in \mathbb{N}\}$ cover $E \cap R^c$, we have

$$\mu^*(E \cap R) \leq \sum_{i=1}^{\infty} \mu(\tilde{R}_i), \quad \mu^*(E \cap R^c) \leq \sum_{i=1}^{\infty} \mu(S_i).$$

Hence,

$$\mu^*(E) + \epsilon \geq \mu^*(E \cap R) + \mu^*(E \cap R^c).$$

Since $\epsilon > 0$ is arbitrary, it follows that

$$\mu^*(E) \geq \mu^*(E \cap R) + \mu^*(E \cap R^c),$$

which proves the result. \square

An open rectangle R° is a union of an increasing sequence of closed rectangles whose volumes approach $\mu(R)$; for example

$$\begin{aligned} & (a_1, b_1) \times (a_2, b_2) \times \cdots \times (a_n, b_n) \\ &= \bigcup_{k=1}^{\infty} \left[a_1 + \frac{1}{k}, b_1 - \frac{1}{k} \right] \times \left[a_2 + \frac{1}{k}, b_2 - \frac{1}{k} \right] \times \cdots \times \left[a_n + \frac{1}{k}, b_n - \frac{1}{k} \right]. \end{aligned}$$

Thus, R° is measurable and, from Proposition 1.6,

$$\mu(R^\circ) = \mu(R).$$

Moreover if $\partial R = R \setminus R^\circ$ denotes the boundary of R , then

$$\mu(\partial R) = \mu(R) - \mu(R^\circ) = 0.$$

2.4. Sets of measure zero and completeness

Sets of measure zero play a particularly important role in measure theory and integration. First, we show that all sets with outer Lebesgue measure zero are Lebesgue measurable.

Proposition 2.12. *If $N \subset \mathbb{R}^n$ and $\mu^*(N) = 0$, then N is Lebesgue measurable, and the measure space $(\mathbb{R}^n, \mathcal{L}(\mathbb{R}^n), \mu)$ is complete.*

PROOF. If $N \subset \mathbb{R}^n$ has outer Lebesgue measure zero and $E \subset \mathbb{R}^n$, then

$$0 \leq \mu^*(E \cap N) \leq \mu^*(N) = 0,$$

so $\mu^*(E \cap N) = 0$. Therefore, since $E \supset E \cap N^c$,

$$\mu^*(E) \geq \mu^*(E \cap N^c) = \mu^*(E \cap N) + \mu^*(E \cap N^c),$$

which shows that N is measurable. If N is a measurable set with $\mu(N) = 0$ and $M \subset N$, then $\mu^*(M) = 0$, since $\mu^*(M) \leq \mu(N)$. Therefore M is measurable and $(\mathbb{R}^n, \mathcal{L}(\mathbb{R}^n), \mu)$ is complete. \square

In view of the importance of sets of measure zero, we formulate their definition explicitly.

Definition 2.13. A subset $N \subset \mathbb{R}^n$ has Lebesgue measure zero if for every $\epsilon > 0$ there exists a countable collection of rectangles $\{R_i : i \in \mathbb{N}\}$ such that

$$N \subset \bigcup_{i=1}^{\infty} R_i, \quad \sum_{i=1}^{\infty} \mu(R_i) < \epsilon.$$

The argument in Example 2.3 shows that every countable set has Lebesgue measure zero, but sets of measure zero may be uncountable; in fact the fine structure of sets of measure zero is, in general, very intricate.

Example 2.14. The standard Cantor set, obtained by removing ‘middle thirds’ from $[0, 1]$, is an uncountable set of zero one-dimensional Lebesgue measure.

Example 2.15. The x -axis in \mathbb{R}^2

$$A = \{(x, 0) \in \mathbb{R}^2 : x \in \mathbb{R}\}$$

has zero two-dimensional Lebesgue measure.

2.5. Translational invariance

An important geometric property of Lebesgue measure is its translational invariance. If $A \subset \mathbb{R}^n$ and $h \in \mathbb{R}^n$, let

$$A + h = \{x + h : x \in A\}$$

denote the translation of A by h .

Proposition 2.16. *If $A \subset \mathbb{R}^n$ and $h \in \mathbb{R}^n$, then*

$$\mu^*(A + h) = \mu^*(A),$$

and $A + h$ is measurable if and only if A is measurable.

PROOF. The invariance of outer measure μ^* result is an immediate consequence of the definition, since $\{R_i + h : i \in \mathbb{N}\}$ is a cover of $A + h$ if and only if $\{R_i : i \in \mathbb{N}\}$ is a cover of A , and $\mu(R + h) = \mu(R)$ for every rectangle R . Moreover, the Carathéodory definition of measurability is invariant under translations since

$$(E + h) \cap (A + h) = (E \cap A) + h.$$

□

The space \mathbb{R}^n is a locally compact topological (abelian) group with respect to translation, which is a continuous operation. More generally, there exists a (left or right) translation-invariant measure, called Haar measure, on any locally compact topological group; this measure is unique up to a scalar factor.

The following is the standard example of a non-Lebesgue measurable set.

Example 2.17. Define an equivalence relation \sim on \mathbb{R} by $x \sim y$ if $x - y \in \mathbb{Q}$. This relation has uncountably many equivalence classes, each of which contains a countably infinite number of points and is dense in \mathbb{R} . Let $E \subset [0, 1]$ be a set that contains exactly one element from each equivalence class, so that \mathbb{R} is the disjoint union of the countable collection of rational translates of E . Then we claim that E is not Lebesgue measurable.

To show this, suppose for contradiction that E is measurable. Let $\{q_i : i \in \mathbb{N}\}$ be an enumeration of the rational numbers in the interval $[-1, 1]$ and let $E_i = E + q_i$ denote the translation of E by q_i . Then the sets E_i are disjoint and

$$[0, 1] \subset \bigcup_{i=1}^{\infty} E_i \subset [-1, 2].$$

The translational invariance of Lebesgue measure implies that each E_i is measurable with $\mu(E_i) = \mu(E)$, and the countable additivity of Lebesgue measure implies that

$$1 \leq \sum_{i=1}^{\infty} \mu(E_i) \leq 3.$$

But this is impossible, since $\sum_{i=1}^{\infty} \mu(E_i)$ is either 0 or ∞ , depending on whether $\mu(E) = 0$ or $\mu(E) > 0$.

The above example is geometrically simpler on the circle $\mathbb{T} = \mathbb{R}/\mathbb{Z}$. When reduced modulo one, the sets $\{E_i : i \in \mathbb{N}\}$ partition \mathbb{T} into a countable union of disjoint sets which are translations of each other. Since the sets have equal measures, their measures cannot sum to a finite nonzero number, but the measure of \mathbb{T} is one.

2.6. Borel sets

The relationship between measure and topology is not a simple one. In this section, we show that all open and closed sets in \mathbb{R}^n , and therefore all Borel sets (*i.e.* sets that belong to the σ -algebra generated by the open sets), are Lebesgue measurable.

Let $\mathcal{T}(\mathbb{R}^n) \subset \mathcal{P}(\mathbb{R}^n)$ denote the standard topology on \mathbb{R}^n consisting of all open sets. That is, $G \subset \mathbb{R}^n$ belongs to $\mathcal{T}(\mathbb{R}^n)$ if for every $x \in G$ there exists $r > 0$ such that $B_r(x) \subset G$, where

$$B_r(x) = \{y \in \mathbb{R}^n : |x - y| < r\}$$

is the open ball of radius r centered at $x \in \mathbb{R}^n$ and $|\cdot|$ denotes the Euclidean norm.

Definition 2.18. The Borel σ -algebra $\mathcal{B}(\mathbb{R}^n)$ on \mathbb{R}^n is the σ -algebra generated by the open sets, $\mathcal{B}(\mathbb{R}^n) = \sigma(\mathcal{T}(\mathbb{R}^n))$. A set that belongs to the Borel σ -algebra is called a Borel set.

Since σ -algebras are closed under complementation, the Borel σ -algebra is also generated by the closed sets in \mathbb{R}^n . Moreover, since \mathbb{R}^n is σ -compact (*i.e.* it is a countable union of compact sets) its Borel σ -algebra is generated by the compact sets.

Remark 2.19. This definition is not constructive, since we start with the power set of \mathbb{R}^n and narrow it down until we obtain the smallest σ -algebra that contains the open sets. It is surprisingly complicated to obtain $\mathcal{B}(\mathbb{R}^n)$ by starting from the open or closed sets and taking successive complements, countable unions, and countable intersections. These operations give sequences of collections of sets in \mathbb{R}^n

$$(2.8) \quad G \subset G_\delta \subset G_{\delta\sigma} \subset G_{\delta\sigma\delta} \subset \dots, \quad F \subset F_\sigma \subset F_{\sigma\delta} \subset F_{\delta\sigma\delta} \subset \dots,$$

where G denotes the open sets, F the closed sets, σ the operation of countable unions, and δ the operation of countable intersections. These collections contain each other; for example, $F_\sigma \supset G$ and $G_\delta \supset F$. This process, however, has to be repeated a number of times equal to the first uncountable ordinal before we obtain $\mathcal{B}(\mathbb{R}^n)$. This is because if, for example, $\{A_i : i \in \mathbb{N}\}$ is a countable family of sets such that

$$A_1 \in G_\delta \setminus G, \quad A_2 \in G_{\delta\sigma} \setminus G_\delta, \quad A_3 \in G_{\delta\sigma\delta} \setminus G_{\delta\sigma}, \dots$$

and so on, then there is no guarantee that $\bigcup_{i=1}^{\infty} A_i$ or $\bigcap_{i=1}^{\infty} A_i$ belongs to any of the previously constructed families. In general, one only knows that they belong to the $\omega + 1$ iterates $G_{\delta\sigma\delta\dots\sigma}$ or $G_{\delta\sigma\delta\dots\delta}$, respectively, where ω is the ordinal number of \mathbb{N} . A similar argument shows that in order to obtain a family which is closed under countable intersections or unions, one has to continue this process until one has constructed an uncountable number of families.

To show that open sets are measurable, we will represent them as countable unions of rectangles. Every open set in \mathbb{R} is a countable disjoint union of open intervals (one-dimensional open rectangles). When $n \geq 2$, it is not true that every open set in \mathbb{R}^n is a countable disjoint union of open rectangles, but we have the following substitute.

Proposition 2.20. *Every open set in \mathbb{R}^n is a countable union of almost disjoint rectangles.*

PROOF. Let $G \subset \mathbb{R}^n$ be open. We construct a family of cubes (rectangles of equal sides) as follows. First, we bisect \mathbb{R}^n into almost disjoint cubes $\{Q_i : i \in \mathbb{N}\}$ of side one with integer coordinates. If $Q_i \subset G$, we include Q_i in the family, and if Q_i is disjoint from G , we exclude it. Otherwise, we bisect the sides of Q_i to obtain 2^n almost disjoint cubes of side one-half and repeat the procedure. Iterating this process arbitrarily many times, we obtain a countable family of almost disjoint cubes.

The union of the cubes in this family is contained in G , since we only include cubes that are contained in G . Conversely, if $x \in G$, then since G is open some sufficiently small cube in the bisection procedure that contains x is entirely contained

in G , and the largest such cube is included in the family. Hence the union of the family contains G , and is therefore equal to G . \square

Proposition 2.21. *The Borel algebra $\mathcal{B}(\mathbb{R}^n)$ is generated by the collection of rectangles $\mathcal{R}(\mathbb{R}^n)$. Every Borel set is Lebesgue measurable.*

PROOF. Since \mathcal{R} is a subset of the closed sets, we have $\sigma(\mathcal{R}) \subset \mathcal{B}$. Conversely, by the previous proposition, $\sigma(\mathcal{R}) \supset \mathcal{T}$, so $\sigma(\mathcal{R}) \supset \sigma(\mathcal{T}) = \mathcal{B}$, and therefore $\mathcal{B} = \sigma(\mathcal{R})$. From Proposition 2.11, we have $\mathcal{R} \subset \mathcal{L}$. Since \mathcal{L} is a σ -algebra, it follows that $\sigma(\mathcal{R}) \subset \mathcal{L}$, so $\mathcal{B} \subset \mathcal{L}$. \square

Note that if

$$G = \bigcup_{i=1}^{\infty} R_i$$

is a decomposition of an open set G into an almost disjoint union of closed rectangles, then

$$G \supset \bigcup_{i=1}^{\infty} R_i^{\circ}$$

is a disjoint union, and therefore

$$\sum_{i=1}^{\infty} \mu(R_i^{\circ}) \leq \mu(G) \leq \sum_{i=1}^{\infty} \mu(R_i).$$

Since $\mu(R_i^{\circ}) = \mu(R_i)$, it follows that

$$\mu(G) = \sum_{i=1}^{\infty} \mu(R_i)$$

for any such decomposition and that the sum is independent of the way in which G is decomposed into almost disjoint rectangles.

The Borel σ -algebra \mathcal{B} is not complete and is strictly smaller than the Lebesgue σ -algebra \mathcal{L} . In fact, one can show that the cardinality of \mathcal{B} is equal to the cardinality \mathfrak{c} of the real numbers, whereas the cardinality of \mathcal{L} is equal to $2^{\mathfrak{c}}$. For example, the Cantor set is a set of measure zero with the same cardinality as \mathbb{R} and every subset of the Cantor set is Lebesgue measurable.

We can obtain examples of sets that are Lebesgue measurable but not Borel measurable by considering subsets of sets of measure zero. In the following example of such a set in \mathbb{R} , we use some properties of measurable functions which will be proved later.

Example 2.22. Let $f : [0, 1] \rightarrow [0, 1]$ denote the standard Cantor function and define $g : [0, 1] \rightarrow [0, 1]$ by

$$g(y) = \inf \{x \in [0, 1] : f(x) = y\}.$$

Then g is an increasing, one-to-one function that maps $[0, 1]$ onto the Cantor set C . Since g is increasing it is Borel measurable, and the inverse image of a Borel set under g is Borel. Let $E \subset [0, 1]$ be a non-Lebesgue measurable set. Then $F = g(E) \subset C$ is Lebesgue measurable, since it is a subset of a set of measure zero, but F is not Borel measurable, since if it was $E = g^{-1}(F)$ would be Borel.

Other examples of Lebesgue measurable sets that are not Borel sets arise from the theory of product measures in \mathbb{R}^n for $n \geq 2$. For example, let $N = E \times \{0\} \subset \mathbb{R}^2$ where $E \subset \mathbb{R}$ is a non-Lebesgue measurable set in \mathbb{R} . Then N is a subset of the x -axis, which has two-dimensional Lebesgue measure zero, so N belongs to $\mathcal{L}(\mathbb{R}^2)$ since Lebesgue measure is complete. One can show, however, that if a set belongs to $\mathcal{B}(\mathbb{R}^2)$ then every section with fixed x or y coordinate, belongs to $\mathcal{B}(\mathbb{R})$; thus, N cannot belong to $\mathcal{B}(\mathbb{R}^2)$ since the section $E \notin \mathcal{B}(\mathbb{R})$.

As we show below, $\mathcal{L}(\mathbb{R}^n)$ is the completion of $\mathcal{B}(\mathbb{R}^n)$ with respect to Lebesgue measure, meaning that we get all Lebesgue measurable sets by adjoining all subsets of Borel sets of measure zero to the Borel σ -algebra and taking unions of such sets.

2.7. Borel regularity

Regularity properties of measures refer to the possibility of approximating in measure one class of sets (for example, nonmeasurable sets) by another class of sets (for example, measurable sets). Lebesgue measure is Borel regular in the sense that Lebesgue measurable sets can be approximated in measure from the outside by open sets and from the inside by closed sets, and they can be approximated by Borel sets up to sets of measure zero. Moreover, there is a simple criterion for Lebesgue measurability in terms of open and closed sets.

The following theorem expresses a fundamental approximation property of Lebesgue measurable sets by open and compact sets.

Theorem 2.23. *If $A \subset \mathbb{R}^n$, then*

$$(2.9) \quad \mu^*(A) = \inf \{ \mu(G) : A \subset G, G \text{ open} \},$$

and if A is Lebesgue measurable, then

$$(2.10) \quad \mu(A) = \sup \{ \mu(K) : K \subset A, K \text{ compact} \}.$$

PROOF. First, we prove (2.9). The result is immediate if $\mu^*(A) = \infty$, so we suppose that $\mu^*(A)$ is finite. If $A \subset G$, then $\mu^*(A) \leq \mu(G)$, so

$$\mu^*(A) \leq \inf \{ \mu(G) : A \subset G, G \text{ open} \},$$

and we just need to prove the reverse inequality.

Let $\epsilon > 0$. There is a cover $\{R_i : i \in \mathbb{N}\}$ of A by rectangles R_i such that

$$\sum_{i=1}^{\infty} \mu(R_i) \leq \mu^*(A) + \frac{\epsilon}{2}.$$

Let S_i be an open rectangle whose interior S_i° contains R_i such that

$$\mu(S_i) \leq \mu(R_i) + \frac{\epsilon}{2^{i+1}}.$$

Then the collection of open rectangles $\{S_i^\circ : i \in \mathbb{N}\}$ covers A and

$$G = \bigcup_{i=1}^{\infty} S_i^\circ$$

is an open set that contains A . Moreover, since $\{S_i : i \in \mathbb{N}\}$ covers G ,

$$\mu(G) \leq \sum_{i=1}^{\infty} \mu(S_i) \leq \sum_{i=1}^{\infty} \mu(R_i) + \frac{\epsilon}{2},$$

and therefore

$$(2.11) \quad \mu(G) \leq \mu^*(A) + \epsilon.$$

It follows that

$$\inf \{ \mu(G) : A \subset G, G \text{ open} \} \leq \mu^*(A) + \epsilon,$$

which proves (2.9) since $\epsilon > 0$ is arbitrary.

Next, we prove (2.10). If $K \subset A$, then $\mu(K) \leq \mu(A)$, so

$$\sup \{ \mu(K) : K \subset A, K \text{ compact} \} \leq \mu(A).$$

Therefore, we just need to prove the reverse inequality,

$$(2.12) \quad \mu(A) \leq \sup \{ \mu(K) : K \subset A, K \text{ compact} \}.$$

First, suppose that A is a bounded measurable set, in which case $\mu(A) < \infty$. Let $F \subset \mathbb{R}^n$ be a compact set that contains A . By the preceding result, for any $\epsilon > 0$, there is an open set $G \supset F \setminus A$ such that

$$\mu(G) \leq \mu(F \setminus A) + \epsilon.$$

Then $K = F \setminus G$ is a compact set such that $K \subset A$. Moreover, $F \subset K \cup G$ and $F = A \cup (F \setminus A)$, so

$$\mu(F) \leq \mu(K) + \mu(G), \quad \mu(F) = \mu(A) + \mu(F \setminus A).$$

It follows that

$$\begin{aligned} \mu(A) &= \mu(F) - \mu(F \setminus A) \\ &\leq \mu(F) - \mu(G) + \epsilon \\ &\leq \mu(K) + \epsilon, \end{aligned}$$

which implies (2.12) and proves the result for bounded, measurable sets.

Now suppose that A is an unbounded measurable set, and define

$$(2.13) \quad A_k = \{x \in A : |x| \leq k\}.$$

Then $\{A_k : k \in \mathbb{N}\}$ is an increasing sequence of bounded measurable sets whose union is A , so

$$(2.14) \quad \mu(A_k) \uparrow \mu(A) \quad \text{as } k \rightarrow \infty.$$

If $\mu(A) = \infty$, then $\mu(A_k) \rightarrow \infty$ as $k \rightarrow \infty$. By the previous result, we can find a compact set $K_k \subset A_k \subset A$ such that

$$\mu(K_k) + 1 \geq \mu(A_k)$$

so that $\mu(K_k) \rightarrow \infty$. Therefore

$$\sup \{ \mu(K) : K \subset A, K \text{ compact} \} = \infty,$$

which proves the result in this case.

Finally, suppose that A is unbounded and $\mu(A) < \infty$. From (2.14), for any $\epsilon > 0$ we can choose $k \in \mathbb{N}$ such that

$$\mu(A) \leq \mu(A_k) + \frac{\epsilon}{2}.$$

Moreover, by the previous result, there is a compact set $K \subset A_k$ such that

$$\mu(A_k) \leq \mu(K) + \frac{\epsilon}{2}.$$

Therefore, for every $\epsilon > 0$ there is a compact set $K \subset A$ such that

$$\mu(A) \leq \mu(K) + \epsilon,$$

which gives (2.12), and the theorem follows. \square

It follows that we may determine the Lebesgue measure of a measurable set in terms of the Lebesgue measure of open or compact sets by approximating the set from the outside by open sets or from the inside by compact sets.

The outer approximation in (2.9) does not require that A is measurable. Thus, for any set $A \subset \mathbb{R}^n$, given $\epsilon > 0$, we can find an open set $G \supset A$ such that $\mu(G) - \mu^*(A) < \epsilon$. This condition does *not* imply, in general, that $\mu^*(G \setminus A) < \epsilon$. However, as we show in the next theorem, the latter condition holds if and only if A is measurable.

Theorem 2.24. *A subset $A \subset \mathbb{R}^n$ is Lebesgue measurable if and only if for every $\epsilon > 0$ there is an open set $G \supset A$ such that*

$$(2.15) \quad \mu^*(G \setminus A) < \epsilon.$$

PROOF. First we assume that A is measurable and show that it satisfies the condition given in the theorem.

Suppose that $\mu(A) < \infty$ and let $\epsilon > 0$. From (2.11) there is an open set $G \supset A$ such that $\mu(G) < \mu^*(A) + \epsilon$. Then, since A is measurable,

$$\mu^*(G \setminus A) = \mu^*(G) - \mu^*(G \cap A) = \mu(G) - \mu^*(A) < \epsilon,$$

which proves the result when A has finite measure.

If $\mu(A) = \infty$, define $A_k \subset A$ as in (2.13), and let $\epsilon > 0$. Since A_k is measurable with finite measure, the argument above shows that for each $k \in \mathbb{N}$, there is an open set $G_k \supset A_k$ such that

$$\mu(G_k \setminus A_k) < \frac{\epsilon}{2^k}.$$

Then $G = \bigcup_{k=1}^{\infty} G_k$ is an open set that contains A , and

$$\mu^*(G \setminus A) = \mu^*\left(\bigcup_{k=1}^{\infty} G_k \setminus A\right) \leq \sum_{k=1}^{\infty} \mu^*(G_k \setminus A) \leq \sum_{k=1}^{\infty} \mu^*(G_k \setminus A_k) < \epsilon.$$

Conversely, suppose that $A \subset \mathbb{R}^n$ satisfies the condition in the theorem. Let $\epsilon > 0$, and choose an open set $G \supset A$ such that $\mu^*(G \setminus A) \leq \epsilon$. If $E \subset \mathbb{R}^n$, we have

$$E \cap A^c = (E \cap G^c) \cup (E \cap (G \setminus A)).$$

Hence, by the subadditivity and monotonicity of μ^* and the measurability of G ,

$$\begin{aligned} \mu^*(E \cap A) + \mu^*(E \cap A^c) &\leq \mu^*(E \cap A) + \mu^*(E \cap G^c) + \mu^*(E \cap (G \setminus A)) \\ &\leq \mu^*(E \cap G) + \mu^*(E \cap G^c) + \mu^*(G \setminus A) \\ &\leq \mu^*(E) + \epsilon \end{aligned}$$

Since $\epsilon > 0$ is arbitrary, it follows that

$$\mu^*(E) \geq \mu^*(E \cap A) + \mu^*(E \cap A^c)$$

which proves that A is measurable. \square

This theorem states that a set is Lebesgue measurable if and only if it can be approximated from the outside by an open set in such a way that the difference has arbitrarily small outer Lebesgue measure. This condition can be adopted as the starting point for the definition of Lebesgue measure (rather than the Carathéodory definition which we have used) *c.f.* [5, 6, 8].

The following theorem gives another characterization of Lebesgue measurable sets, as ones that can be ‘squeezed’ between open and closed sets.

Theorem 2.25. *A subset $A \subset \mathbb{R}^n$ is Lebesgue measurable if and only if for every $\epsilon > 0$ there is an open set G and a closed set F such that $G \supset A \supset F$ and*

$$(2.16) \quad \mu(G \setminus F) < \epsilon.$$

If $\mu(A) < \infty$, then F may be chosen to be compact.

PROOF. If A satisfies the condition in the theorem, then it follows from the monotonicity of μ^* that $\mu^*(G \setminus A) \leq \mu(G \setminus F) < \epsilon$, so A is measurable by Theorem 2.24.

Conversely, if A is measurable then A^c is measurable, and by Theorem 2.24 given $\epsilon > 0$, we can choose open sets $G \supset A$ and $H \supset A^c$ such that

$$\mu^*(G \setminus A) < \frac{\epsilon}{2}, \quad \mu^*(H \setminus A^c) < \frac{\epsilon}{2}.$$

Then, defining the closed set $F = H^c$, we have $G \supset A \supset F$ and

$$\mu(G \setminus F) \leq \mu^*(G \setminus A) + \mu^*(A \setminus F) = \mu^*(G \setminus A) + \mu^*(H \setminus A^c) < \epsilon.$$

Finally, suppose that $\mu(A) < \infty$ and let $\epsilon > 0$. From Theorem 2.23, since A is measurable, there is a compact set $K \subset A$ such that $\mu(A) < \mu(K) + \epsilon/2$ and

$$\mu(A \setminus K) = \mu(A) - \mu(K) < \frac{\epsilon}{2}.$$

As before, from Theorem 2.24 there is an open set $G \supset A$ such that

$$\mu(G) < \mu(A) + \epsilon/2.$$

It follows that $G \supset A \supset K$ and

$$\mu(G \setminus K) = \mu(G \setminus A) + \mu(A \setminus K) < \epsilon,$$

which shows that we may take $F = K$ compact when A has finite measure. \square

Using open and closed sets, we can approximate measurable sets up to sets of arbitrarily small but, in general, nonzero measure. By taking countable intersections of open sets or countable unions of closed sets, we can approximate measurable sets up to sets of measure zero

Definition 2.26. The collection of sets in \mathbb{R}^n that are countable intersections of open sets is denoted by $G_\delta(\mathbb{R}^n)$, and the collection of sets in \mathbb{R}^n that are countable unions of closed sets is denoted by $F_\sigma(\mathbb{R}^n)$.

G_δ and F_σ sets are Borel. Thus, it follows from the next result that every Lebesgue measurable set can be approximated up to a set of measure zero by a Borel set. This is the Borel regularity of Lebesgue measure.

Theorem 2.27. *Suppose that $A \subset \mathbb{R}^n$ is Lebesgue measurable. Then there exist sets $G \in G_\delta(\mathbb{R}^n)$ and $F \in F_\sigma(\mathbb{R}^n)$ such that*

$$G \supset A \supset F, \quad \mu(G \setminus A) = \mu(A \setminus F) = 0.$$

PROOF. For each $k \in \mathbb{N}$, choose an open set G_k and a closed set F_k such that $G_k \supset A \supset F_k$ and

$$\mu(G_k \setminus F_k) \leq \frac{1}{k}$$

Then

$$G = \bigcap_{k=1}^{\infty} G_k, \quad F = \bigcup_{k=1}^{\infty} F_k$$

are G_δ and F_σ sets with the required properties. \square

In particular, since any measurable set can be approximated up to a set of measure zero by a G_δ or an F_σ , the complexity of the transfinite construction of general Borel sets illustrated in (2.8) is ‘hidden’ inside sets of Lebesgue measure zero.

As a corollary of this result, we get that the Lebesgue σ -algebra is the completion of the Borel σ -algebra with respect to Lebesgue measure.

Theorem 2.28. *The Lebesgue σ -algebra $\mathcal{L}(\mathbb{R}^n)$ is the completion of the Borel σ -algebra $\mathcal{B}(\mathbb{R}^n)$.*

PROOF. By the previous theorem, if $A \subset \mathbb{R}^n$ is Lebesgue measurable, then there is a F_σ set $F \subset A$ such that $M = A \setminus F$ has Lebesgue measure zero. It follows by the approximation theorem that there is a Borel set $N \in G_\delta$ with $\mu(N) = 0$ and $M \subset N$. Thus, $A = F \cup M$ where $F \in \mathcal{B}$ and $M \subset N \in \mathcal{B}$ with $\mu(N) = 0$, which proves that $\mathcal{L}(\mathbb{R}^n)$ is the completion of $\mathcal{B}(\mathbb{R}^n)$ as given in Theorem 1.8. \square

