CALCULUS: Math 21C, Spring 2018 Sample Questions: Midterm 2

1. Determine the interval of convergence (including the endpoints) for the following power series. State explicitly for what values of x the series converges absolutely, converges conditionally, and diverges. In each case, specify the radius of convergence R and the center of the interval of convergence a.

(a)
$$\sum_{n=1}^{\infty} \frac{(-1)^n 2^n}{n} (x-1)^n$$
; (b) $\sum_{n=0}^{\infty} \frac{1}{3^n+1} x^{2n}$; (c) $\sum_{n=0}^{\infty} \frac{1}{n^2 5^n} (2x+1)^n$.

2. Let

$$\vec{u} = 2\vec{i} - \vec{j} + 3\vec{k}, \quad \vec{v} = -\vec{i} + 2\vec{j} + 2\vec{k}.$$

Compute: (a) $|\vec{u}|$; (b) $|\vec{v}|$; (c) the angle θ between \vec{u} , \vec{v} (you can express it as an inverse trigonometric function); (d) the projection $\operatorname{proj}_{\vec{v}}\vec{u}$ of \vec{u} in the direction of \vec{v} .

3. (a) Find the area of the triangle with vertices P(-2, 2, 0), Q(0, 1, -1) and R(-1, 2, -2).

(b) Find a parametric equation for the line in which the planes 3x-6y-4z = 15 and 6x + y - 2z = 5 intersect.

4. Let

$$f(x,y) = e^{xy}\ln(y).$$

Compute the partial derivatives f_x , f_y , f_{xx} , f_{xy} and f_{yy} . (You do NOT need to simplify your answers.)

5. (a) Find the Taylor polynomial $P_2(x)$ of order 2 centered at x = 0 for the function $f(x) = e^{-x^2}$.

(b) Use Taylor's theorem with remainder $R_2(x)$ to estimate the maximum error $|f(x) - P_2(x)|$ for $0 \le x \le 1$.

(c) Use the results of (a) and (b) to obtain an approximate value for the integral

$$\int_0^1 e^{-x^2} \, dx$$

and estimate the maximum error in your approximate value for the integral.

6. Suppose that the functions f(x), g(x) have the Taylor series expansions at zero, up to second degree terms, given by

$$f(x) = a_0 + a_1 x + a_2 x^2 + \dots, \qquad g(x) = b_0 + b_1 x + b_2 x^2 + \dots$$

(a) According to Taylor's theorem, how are a_0 , a_1 , a_2 given in terms of f and its derivatives and b_0 , b_1 , b_2 in terms of g and its derivative?

(b) Find the Taylor series for h(x) = f(x)g(x) at zero, up to second degree terms, by multiplying the Taylor series for f(x) and g(x).

(c) Use the product rule to compute h'(x), h''(x) in terms of the derivatives of f(x), g(x). Show that the use of these expressions in Taylor's theorem for h(x) gives the same series as the one you found in (b).