

Section 3.1

2.) $f(x) = x + 32x^{-2} \xrightarrow{D} f'(x) = 1 - 64x^{-3}$

at $x = 2 \rightarrow f'(2) = 1 - 64(2)^{-3} = 1 - 64\left(\frac{1}{8}\right) = -7$

at $x = 4 \rightarrow f'(4) = 1 - 64(4)^{-3} = 1 - 64\left(\frac{1}{64}\right) = 0$

at $x = 8 \rightarrow f'(8) = 1 - 64(8)^{-3} = 1 - 64\left(\frac{1}{8}\right)^3 = \frac{7}{8}$

3.) $f(x) = (x+2)^{2/3} \xrightarrow{D} f'(x) = \frac{2}{3}(x+2)^{-1/3}$

at $x = -3 \rightarrow f'(-3) = \frac{2}{3}(-1)^{-1/3} = \frac{2}{3}(-1) = -\frac{2}{3}$

at $x = -2 \rightarrow f'(-2) = \frac{2}{3}(0)^{-1/3} = \frac{2}{3(0)^{1/3}}$ so

$f'(-2)$ DNE (corner)

at $x = 1 \rightarrow f'(1) = \frac{2}{3}(1)^{-1/3} = \frac{2}{3}(1) = \frac{2}{3}$

6.) $f(x) = \frac{x^3}{4} - 3x \xrightarrow{D} f'(x) = \frac{3}{4}x^2 - 3$

$= 3\left(\frac{x^2}{4} - 1\right) = 0 \rightarrow x = 2, x = -2$

	+	0	-	0	+	
	-----					f'
rel.	{	$x = -2$	}	$x = 2$	}	rel.
max.	{	$y = 4$	}	$y = -4$	}	min.

f is \uparrow for $x < -2, x > 2$;

f is \downarrow for $-2 < x < 2$.

8.) $f(x) = \frac{x^2}{x+1} \xrightarrow{D} f'(x) = \frac{(x+1)(2x) - x^2(1)}{(x+1)^2}$

$= \frac{2x^2 + 2x - x^2}{(x+1)^2} = \frac{x^2 + 2x}{(x+1)^2} = \frac{x(x+2)}{(x+1)^2} = 0 \rightarrow$

$x = 0, x = -2$: NO

	+	0	-		-	0	+	
	-----							f'
rel.	{	$x = -2$	}	$x = -1$	}	$x = 0$	}	rel.
max.	{	$y = -4$	}	}	}	$y = 0$	}	min.

f is \uparrow for $x < -2, x > 0$;
 f is \downarrow for $-2 < x < -1, -1 < x < 0$.

14.) $y = -x^2 + 2x \xrightarrow{D} y' = -2x + 2 = 0 \rightarrow x = 1 :$

$$\begin{array}{c} + \quad 0 \quad - \\ | \\ x=1 \\ y=1 \end{array} \quad y' \quad \begin{array}{l} y \text{ is } \uparrow \text{ for } x < 1; \\ y \text{ is } \downarrow \text{ for } x > 1. \end{array}$$

15.) $y = x^3 - 6x^2 \xrightarrow{D} y' = 3x^2 - 12x = 3x(x-4) = 0$
 $\rightarrow x = 0, x = 4 :$

$$\begin{array}{c} + \quad 0 \quad - \quad 0 \quad + \\ | \quad | \\ x=0 \quad x=4 \end{array} \quad y' \quad \begin{array}{l} y \text{ is } \uparrow \text{ for } x < 0, x > 4; \\ y \text{ is } \downarrow \text{ for } 0 < x < 4. \end{array}$$

rel. $\left\{ \begin{array}{l} x=0 \\ y=0 \end{array} \right.$ rel. $\left\{ \begin{array}{l} x=4 \\ y=-32 \end{array} \right.$ } min.

18.) $f(x) = \sqrt{4-x^2} \xrightarrow{D} f'(x) = \frac{1}{2}(4-x^2)^{-1/2} \cdot (-2x)$

$= \frac{-x}{\sqrt{4-x^2}} = 0 \rightarrow x = 0 :$

$$\begin{array}{c} \text{No} \quad \text{No} \\ | \quad | \\ x=-2 \quad x=2 \end{array} \quad \begin{array}{c} + \quad 0 \quad - \\ | \\ x=0 \end{array} \quad f' \quad \begin{array}{l} f \text{ is } \uparrow \text{ for } -2 < x < 0; \\ f \text{ is } \downarrow \text{ for } 0 < x < 2. \end{array}$$

rel. $\left\{ \begin{array}{l} x=-2 \\ y=0 \end{array} \right.$ rel. $\left\{ \begin{array}{l} x=0 \\ y=0 \end{array} \right.$ rel. $\left\{ \begin{array}{l} x=2 \\ y=0 \end{array} \right.$ } min.

22.) $y = x^3 - 3x + 2 \xrightarrow{D} y' = 3x^2 - 3 = 3(x-1)(x+1) = 0 \rightarrow$
 $x = 1, x = -1 :$

$$\begin{array}{c} + \quad 0 \quad - \quad 0 \quad + \\ | \quad | \\ x=-1 \quad x=1 \end{array} \quad y' \quad \begin{array}{l} y \text{ is } \uparrow \text{ for } x < -1, x > 1; \\ y \text{ is } \downarrow \text{ for } -1 < x < 1. \end{array}$$

rel. $\left\{ \begin{array}{l} x=-1 \\ y=4 \end{array} \right.$ rel. $\left\{ \begin{array}{l} x=1 \\ y=0 \end{array} \right.$ } min.

$$23.) f(x) = x\sqrt{x+1} \xrightarrow{D} f'(x) = x \cdot \frac{1}{2}(x+1)^{-\frac{1}{2}} + (1) \cdot \sqrt{x+1}$$

$$= \frac{x}{2\sqrt{x+1}} + \frac{\sqrt{x+1}}{1} = \frac{x+2(x+1)}{2\sqrt{x+1}} = \frac{3x+2}{2\sqrt{x+1}} = 0 \rightarrow$$

$$x = -\frac{2}{3} :$$

//	NO	-	0	+	f'	
x = -1			x = -2/3			} rel. } min.
			y = -2/3√3			

f is ↑ for $x > -2/3$;
f is ↓ for $-1 < x < -2/3$.

$$25.) f(x) = x^4 - 2x^3 \xrightarrow{D} f'(x) = 4x^3 - 6x^2 = 2x^2(2x-3) = 0$$

$$\rightarrow x=0, x=3/2 :$$

-	0	-	0	+	f'	
x = 0			x = 3/2			} rel. } min.
			y = -27/16			

y is ↑ for $x > 3/2$;
y is ↓ for $x < 0$,
 $0 < x < 3/2$.

$$27.) f(x) = \frac{x}{x^2+4} \xrightarrow{D} f'(x) = \frac{(x^2+4)(1) - x \cdot (2x)}{(x^2+4)^2}$$

$$= \frac{x^2+4-2x^2}{(x^2+4)^2} = \frac{4-x^2}{(x^2+4)^2} = \frac{(2-x)(2+x)}{(x^2+4)^2} = 0 \rightarrow$$

$$x=2, x=-2 :$$

-	0	+	0	-	f'	
rel.	x = -2		x = 2	rel.		} rel. } max.
min.	y = -1/4		y = 1/4			

f is ↑ for $-2 < x < 2$;
f is ↓ for $x < -2$,
 $x > 2$.

$$35.) C = \frac{10}{x} + \frac{10x}{x+3} \xrightarrow{D} C' = \frac{-10}{x^2} + \frac{(x+3)(10) - (10x)(1)}{(x+3)^2}$$


$$= \frac{-10}{x^2} + \frac{10x+30-10x}{(x+3)^2} = \frac{-10(x^2+6x+9) + 30x^2}{x^2(x+3)^2}$$

$$= \frac{-10x^2 - 60x - 90 + 30x^2}{x^2(x+3)^2} = \frac{20x^2 - 60x - 90}{x^2(x+3)^2}$$

$$= \frac{10(2x^2 - 6x - 9)}{x^2(x+3)^2} = 0 \rightarrow 2x^2 - 6x - 9 = 0 \rightarrow$$


$$x = \frac{6 \pm \sqrt{36 + 72}}{4} = \frac{6 \pm 6\sqrt{3}}{4} = \frac{3 \pm 3\sqrt{3}}{2} \rightarrow$$

$$x = \frac{3 + 3\sqrt{3}}{2} \approx 4.1 \text{ cars} :$$



$$40.) P = 2.36x - \frac{1}{25,000}x^2 - 3500 \quad \frac{D}{\rightarrow}$$

$$P' = 2.36 - \frac{1}{12,500}x \rightarrow x = 29,500 \text{ bags} :$$



P is \uparrow for $0 < x < 29,500$;

P is \downarrow for $29,500 < x < 50,000$.

Section 3.2

$$5.) g(x) = 6x^3 - 15x^2 + 12x \xrightarrow{D}$$

$$g'(x) = 18x^2 - 30x + 12 = 6(3x^2 - 5x + 2)$$


$$= 6(3x-2)(x-1) = 0 \rightarrow x = \frac{2}{3}, x = 1 :$$

$$\begin{array}{c} + \quad 0 \quad - \quad 0 \quad + \\ \hline \end{array} g'$$

$$\begin{array}{l} \text{rel. } \left\{ \begin{array}{l} x = \frac{2}{3} \\ y = \frac{28}{9} \end{array} \right. \text{ max.} \\ \text{rel. } \left\{ \begin{array}{l} x = 1 \\ y = 3 \end{array} \right. \text{ min.} \end{array}$$

g is \uparrow for $x < \frac{2}{3}, x > 1$;

g is \downarrow for $\frac{2}{3} < x < 1$;



$$x=0 : y=0$$

$$y=0 : x(6x^2 - 15x + 12) = 0 \rightarrow x=0$$

$$8.) h(x) = 2(x-3)^3 \xrightarrow{D} h'(x) = 6(x-3)^2 = 0 \rightarrow$$

$$x=3 :$$

$$\begin{array}{c} + \quad 0 \quad + \\ \hline \end{array} h'$$


$$x=3$$

$$y=0$$

h is \uparrow for $x < 3, x > 3$;

$$x=0 : y = -54$$

$$y=0 : x = 3$$



$$16.) f(x) = x + \frac{1}{x} \xrightarrow{D} f'(x) = 1 - \frac{1}{x^2} = \frac{x^2 - 1}{x^2}$$

$$= \frac{(x-1)(x+1)}{x^2} = 0 \rightarrow x=1, x=-1 :$$

NO

+ 0 - | - 0 + f'

rel. $\left. \begin{matrix} \{x=-1 & x=0 & x=1\} \end{matrix} \right\} \begin{matrix} \text{rel.} \\ \text{max} \{y=-2 & y=2\} \text{min.} \end{matrix}$

f is \uparrow for $x < -1, x > 1$;
 f is \downarrow for $-1 < x < 0$,
 $0 < x < 1$;


$x=0$: (NO)

$y=0$: $0 = x + \frac{1}{x} \rightarrow$

$x = \frac{-1}{x} \rightarrow x^2 = -1$ (NO);

$\lim_{x \rightarrow 0^+} (x + \frac{1}{x}) = 0 + \frac{1}{0^+} = +\infty$

$\lim_{x \rightarrow 0^-} (x + \frac{1}{x}) = 0 + \frac{1}{0^-} = -\infty$



V.A. is $x=0$

18.) $h(x) = \frac{4}{x^2+1} \xrightarrow{D} h'(x) = -4(x^2+1)^{-2} \cdot (2x) = \frac{-8x}{(x^2+1)^2} = 0$

$\rightarrow x=0$:

+ 0 - h'

abs. $\left\{ \begin{matrix} x=0 \\ y=4 \end{matrix} \right.$


h is \uparrow for $x < 0$;
 h is \downarrow for $x > 0$;

$x=0$: $y=4$,

$y=0$: (NO);


$\lim_{x \rightarrow \pm\infty} \frac{4}{x^2+1} = \frac{4}{\infty} = 0$

\rightarrow H.A. is $y=0$




20.) $f(x) = \frac{1}{3}(2x+5)$ on $[0, 5] \xrightarrow{D} y$

$f'(x) = \frac{2}{3}$



abs. $\left. \begin{matrix} x=0 \\ y=5/3 \end{matrix} \right\}$ min. $\left. \begin{matrix} x=5 \\ y=5 \end{matrix} \right\}$ abs. max.

f is \uparrow for $0 < x < 5$



21.) $f(x) = 5 - 2x^2$ on $[-1, 2] \xrightarrow{D}$

$f'(x) = -4x = 0 \rightarrow x = 0$



$x = -1$ $x = 0$ $x = 2$

$y = 3$ $y = 5$ $y = -3$

rel. min. abs. max. abs. min.


f is \uparrow for $-1 < x < 0$;

f is \downarrow for $0 < x < 2$;

$x = 0 : y = 5$

$y = 0 : 0 = 5 - 2x^2 \rightarrow$


$x^2 = \frac{5}{2} \rightarrow x = \pm \sqrt{\frac{5}{2}}$



23.) $f(x) = x^3 - 3x^2$ on $[-1, 4] \xrightarrow{D}$

$f'(x) = 3x^2 - 6x = 3x(x-2) = 0$

$\rightarrow x = 0, x = 2$



$x = -1$ $x = 0$ $x = 2$ $x = 4$

$y = -4$ $y = 0$ $y = -4$ $y = 16$

abs. min. rel. max. abs. min. abs. max.


f is \uparrow for $-1 < x < 0$, $2 < x < 4$;

f is \downarrow for $0 < x < 2$;

$$x=0: y=0$$

$$y=0: x^2(x-3)=0$$

$$\rightarrow x=0, x=3$$



44.) Demand $x = c \cdot \frac{1}{p^3}$, where p : price ;
 if $p = \$10$, $x = 8$ so $8 = c \cdot \frac{1}{1000} \rightarrow c = 8000 \rightarrow$

demand $\boxed{x = \frac{8000}{p^3} \text{ or } p = \frac{20}{x^{1/3}}}$;

cost $C = 25 + 4x$ and revenue

$$R = px = \left(\frac{20}{x^{1/3}}\right)x = 20x^{2/3} \text{ so profit}$$


$$\boxed{P = R - C = 20x^{2/3} - (25 + 4x)}$$

Find maximum profit :

$$P' = \frac{40}{3}x^{-1/3} - 4 = 0 \rightarrow x^{-1/3} = \frac{3}{10} \rightarrow$$

$$x = (x^{-1/3})^{-3} = \left(\frac{3}{10}\right)^{-3} \approx 37;$$

The maximum profit occurs when




$x = 37$ units, $p = \$6$, and profit

$$P = \$49.07.$$

190:45 Let x : demand, p : price and

if $x = 6000$, $p = \$.80$ and $x = 5600$, $p = \$1.00$
 are points on a linear graph, then :



slope is

$$m = \frac{6000 - 5600}{0.8 - 1}$$

$$= \frac{400}{-0.2} = -2000$$

so demand function (line) is

$$X - 5600 = -2000(P - 1) \quad \text{or} \quad \boxed{P = \frac{19}{5} - \frac{X}{2000}} ;$$

cost $C = 5000 + 0.4X$ and revenue

$$R = pX = \frac{19}{5}X - \frac{X^2}{2000} \quad \text{so that profit}$$

$$P = R - C = \frac{19}{5}X - \frac{X^2}{2000} - (5000 + \frac{2}{5}X)$$

$$\text{or} \quad \boxed{P = \frac{-X^2}{2000} + \frac{17}{5}X - 5000} ; \text{ determine}$$

maximum profit \rightarrow

$$P' = \frac{-X}{1000} + \frac{17}{5} = 0 \rightarrow X = 3400$$

so maximum profit occurs when

$$\begin{array}{c} + \quad 0 \quad - \\ \hline X = 3400 \end{array} \quad P'$$

$X = 3400$ case, price $p = \$2.10$, and profit $P = \$780$