

Math 17A  
Kouba  
Functions- Review

DEFINITION : In an equation composed of  $x$ 's and  $y$ 's, variable  $y$  is a function of  $x$  if each admissible  $x$ -value has exactly one  $y$ -value.

NOTE : The graph of a function passes the *vertical line test* . That is, a vertical line passed through the graph will touch the graph in at most one point.

EXAMPLE : Assume that  $xy - 3 = x^2 + 2y$  . Then  $xy - 2y = x^2 + 3 \rightarrow$

$$(x - 2)y = x^2 + 3 \rightarrow$$

$$y = \frac{x^2 + 3}{x - 2} \rightarrow$$

$y$  is a function of  $x$ .

EXAMPLE : Assume that  $xy^2 - 1 = x + y$  . If  $x = 1$ , then

$$y^2 - 1 = 1 + y \rightarrow$$

$$y^2 - y - 2 = 0 \rightarrow$$

$$(y - 2)(y + 1) = 0 \rightarrow$$

$$y = 2 \text{ or } y = -1 \rightarrow$$

$x = 1$  has TWO  $y$ -values  $\rightarrow$

$y$  is NOT a function of  $x$  .

NOTATION : If  $y$  is a function of  $x$ , then we write  $y = f(x)$ .

EXAMPLE : If  $y = x^2 + x$ , then  $y$  is a function of  $x$  and we write  $f(x) = x^2 + x$ ; then

$$\text{a.) } f(-2) = (-2)^2 + (-2) = 4 - 2 = 2.$$

$$\text{b.) } f(2x - 1) = (2x - 1)^2 + (2x - 1) = 4x^2 - 4x + 1 + 2x - 1 = 4x^2 - 2x.$$

DEFINITION : Assume that  $y = f(x)$  is a function. The domain of function  $f$  is the set of all admissible  $x$ -values. The range of function  $f$  is the set of all corresponding  $y$ -values.

EXAMPLE : Consider function  $f(x) = \sqrt{2x - 6}$ . Then  $2x - 6 \geq 0 \rightarrow 2x \geq 6 \rightarrow x \geq 3 \rightarrow$

$$\text{DOMAIN : } x \geq 3.$$

Since  $\sqrt{2x - 6} \geq 0$ ,  $f(3) = 0$ , and  $2x - 6$  gets infinitely large as  $x$  gets infinitely large, it follows that

$$\text{RANGE : } y \geq 0.$$

DEFINITION : A *function*  $y = f(x)$  is one-to-one if each  $y$ -value has exactly one  $x$ -value. More precisely, a one-to-one function has the property that if  $f(x_1) = f(x_2)$  ( $y$ -values are equal), then  $x_1 = x_2$  ( $x$ -values are equal).

NOTE : The graph of a one-to-one function passes the *horizontal line test*. That is, a horizontal line passed through the graph will touch the graph in at most one point.

EXAMPLE : Consider the function (parabola)  $y = x^2 - 5$ . If  $y = 4$ , then

$$4 = x^2 - 5 \rightarrow$$

$$x^2 = 9 \rightarrow$$

$$x = 3 \text{ or } x = -3 \rightarrow$$

$$y = 4 \text{ has TWO } x\text{-values} \rightarrow$$

function  $y$  is NOT one-to-one .

EXAMPLE : Consider the function  $f(x) = \frac{x}{x+3}$  . Prove that  $f$  is one-to-one:

$$\begin{aligned} f(x_1) = f(x_2) &\rightarrow \\ \frac{x_1}{x_1+3} = \frac{x_2}{x_2+3} &\rightarrow \\ x_1(x_2+3) = x_2(x_1+3) &\rightarrow \\ x_1x_2 + 3x_1 = x_1x_2 + 3x_2 &\rightarrow \\ 3x_1 = 3x_2 &\rightarrow \\ x_1 = x_2 &\rightarrow \end{aligned}$$

function  $f$  IS one-to-one .

DEFINITION : Assume that  $y = f(x)$  and  $y = g(x)$  are functions. The composition of functions  $f$  and  $g$  is

$$(f \circ g)(x) = f(g(x)) .$$

EXAMPLE : Consider the functions  $f(x) = \frac{x}{10-x}$  and  $g(x) = \frac{1}{x+8}$ . Then

$$\begin{aligned} (f \circ g)(x) &= f(g(x)) = f\left(\frac{1}{x+8}\right) \\ &= \frac{\frac{1}{x+8}}{10 - \left(\frac{1}{x+8}\right)} \end{aligned}$$

$$\begin{aligned}
&= \frac{1}{10 - \left( \frac{1}{x+8} \right)} \cdot \frac{x+8}{x+8} \\
&= \frac{1}{10(x+8) - 1} \\
&= \frac{1}{10x+79} .
\end{aligned}$$

DEFINITION : The inverse function of function  $y = f(x)$  is the function  $y = f^{-1}(x)$  for which

$$f(f^{-1}(x)) = x \text{ and } f^{-1}(f(x)) = x .$$

FACT : If  $y = f(x)$  is a one-to-one function, then  $f$  has an inverse function.

SEE INVERSE FUNCTION HANDOUT.

EXAMPLE : The function  $f(x) = \frac{x}{x+3}$  is one-to-one. Find its inverse :

$$y = \frac{x}{x+3} \quad \longrightarrow \quad (\text{Switch variables.}) \quad x = \frac{y}{y+3}$$

$$(\text{Solve for } y.) \quad x(y+3) = y \quad \longrightarrow$$

$$xy + 3x = y \quad \longrightarrow$$

$$xy - y = -3x \quad \longrightarrow$$

$$y(x-1) = -3x \quad \longrightarrow$$

$$y = \frac{-3x}{x-1} \quad \longrightarrow \quad \text{inverse function is } f^{-1}(x) = \frac{3x}{1-x} .$$