

Math 21B-B - Homework Set 2

Section 5.3:

1. (a) $\lim_{\|P\| \rightarrow 0} \sum_{k=1}^n (c_k^2 - 3c_k) \Delta x_k$, where P is a partition of $[-7, 5]$.

$$\int_{-7}^5 (x^2 - 3x) dx$$

- (b) $\lim_{\|P\| \rightarrow 0} \sum_{k=1}^n \sqrt{4 - c_k^2} \Delta x_k$, where P is a partition of $[0, 1]$.

$$\int_0^1 \sqrt{4 - x^2} dx$$

- (c) $\lim_{\|P\| \rightarrow 0} \sum_{k=1}^n (\tan c_k) \Delta x_k$, where P is a partition of $[0, \pi/4]$.

$$\int_0^{\pi/4} \tan(x) dx$$

2. Suppose that f and g are integrable and that:

$$\int_1^2 f(x) dx = -4, \quad \int_1^5 f(x) dx = 6, \quad \int_1^5 g(x) dx = 8.$$

(a) $\int_2^2 g(x) dx = 0$

ZERO WIDTH - RULE 2

(B) $\int_5^1 g(x) dx = - \int_1^5 g(x) dx = -8$

ORDER OF INTEGRATION - RULE 1

(C) $\int_1^2 3f(x) dx = 3 \int_1^2 f(x) dx = 3 \cdot -4 = -12.$

CONSTANT MULTIPLE - RULE 3

(D) $\int_2^5 f(x) dx = \int_1^5 f(x) dx - \int_1^2 f(x) dx = 6 - (-4) = 10$

ADDITIVITY - RULE 5

(E) $\int_1^5 [f(x) - g(x)] dx = \int_1^5 f(x) dx - \int_1^5 g(x) dx = 6 - 8 = -2$

SUM AND DIFFERENCE - RULE 4

$$(F) \int_1^5 [4f(x) - g(x)] dx = \int_1^5 4f(x) dx - \int_1^5 g(x) dx = 4 \int_1^5 f(x) dx - \int_1^5 g(x) dx = 4 \cdot 6 - 8 = 24 - 8 = 16$$

CONSTANT MULTIPLE AND SUM/DIFFERENCE - RULES 3 AND 4

3. Suppose that f and h are integrable and that:

$$\int_1^9 f(x) dx = -1, \quad \int_7^9 f(x) dx = 5, \quad \int_7^9 h(x) dx = 4.$$

$$(a) \int_1^9 -2f(x) dx = -2 \int_1^9 f(x) dx = -2 \cdot -1 = 2$$

CONSTANT MULTIPLE - RULE 3

$$(B) \int_7^9 [f(x) + h(x)] dx = \int_7^9 f(x) dx + \int_7^9 h(x) dx = 5 + 4 = 9$$

SUM AND DIFFERENCE - RULE 4

$$(C) \int_7^9 [2f(x) - 3h(x)] dx = \int_7^9 2f(x) dx - \int_7^9 3h(x) dx = 2 \int_7^9 f(x) dx - 3 \int_7^9 h(x) dx = 2 \cdot 5 - 3 \cdot 4 = 10 - 12 = -2$$

CONSTANT MULTIPLE AND SUM/DIFFERENCE - RULES 3 AND 4

$$(D) \int_9^1 f(x) dx = - \int_1^9 f(x) dx = -(-1) = 1$$

ORDER OF INTEGRATION - RULE 1


$$(E) \int_1^7 f(x) dx = \int_1^9 f(x) dx - \int_7^9 f(x) dx = -1 - 5 = -6$$

ADDITIVITY - RULE 5

$$(F) \int_9^7 [h(x) - f(x)] dx = - \int_7^9 [h(x) - f(x)] dx = - \left[\int_7^9 h(x) dx - \int_7^9 f(x) dx \right] = -(4 - 5) = -(-1) = 1$$


ORDER OF INTEGRATION AND SUM/DIFFERENCE - RULES 1 AND 4

$$4. \int_{-2}^1 |x| dx$$




$$\begin{aligned}
 \int_{-2}^1 |x| dx &= \text{Area of left triangle} + \text{Area of right triangle} \\
 &= 1/2 \cdot 2 \cdot 2 + 1/2 \cdot 1 \cdot 1 = 2 + 1/2 \\
 &= 2 + 1/2 \\
 &= 5/2 \text{ square units}
 \end{aligned}$$

5. $\int_{-1}^1 (1 + \sqrt{1 - x^2}) dx$




$$\begin{aligned}
\int_{-1}^1 (1 + \sqrt{1-x^2}) dx &= \text{Area of semicircle} + \text{Area of Rectangle} \\
&= 1/2 \cdot \pi \cdot 1^2 + 1 \cdot 2 \\
&= \frac{\pi}{2} + 2 \text{ square units}
\end{aligned}$$

6. (a) $f(x) = x^2 - 1$ on $[0, \sqrt{3}]$



$$\begin{aligned}
av(f) &= \left(\frac{1}{\sqrt{3}-0} \right) \int_0^{\sqrt{3}} (x^2 - 1) dx \\
&= \left(\frac{1}{\sqrt{3}} \right) \left[\int_0^{\sqrt{3}} x^2 dx - \int_0^{\sqrt{3}} 1 dx \right] \\
&= \left(\frac{1}{\sqrt{3}} \right) \left[\left(\frac{1}{3} \right) x^3 \Big|_0^{\sqrt{3}} - x \Big|_0^{\sqrt{3}} \right] \\
&= \left(\frac{1}{\sqrt{3}} \right) \left[\left(\frac{1}{3} \right) (3\sqrt{3} - 0) - (\sqrt{3} - 0) \right] \\
&= \left(\frac{1}{\sqrt{3}} \right) (\sqrt{3} - \sqrt{3}) \\
&= 0
\end{aligned}$$

(b) $h(x) = -|x|$ on (i.) $[-1, 0]$, (ii.) $[0, 1]$, (iii.) $[-1, 1]$.



i.

$$\begin{aligned}
 av(f) &= \left(\frac{1}{0 - (-1)} \right) \int_{-1}^0 -|x| dx \\
 &= (1) \int_{-1}^0 -(-x) dx \\
 &= \int_{-1}^0 x dx \\
 &= \left. \frac{1}{2}x^2 \right|_{-1}^0 \\
 &= 0 - \frac{1}{2} \\
 &= -\frac{1}{2}
 \end{aligned}$$

ii.

$$\begin{aligned}
 av(f) &= \left(\frac{1}{1 - 0} \right) \int_0^1 -|x| dx \\
 &= (1) \int_0^1 -x dx \\
 &= \left. -\frac{1}{2}x^2 \right|_0^1 \\
 &= -\frac{1}{2}
 \end{aligned}$$

iii.

$$\begin{aligned}av(f) &= \left(\frac{1}{1 - (-1)}\right) \int_{-1}^1 -|x| dx \\&= \left(\frac{1}{2}\right) \left[\int_{-1}^0 -|x| dx + \int_0^1 -|x| dx\right] \\&= \left(\frac{1}{2}\right) \left[\int_{-1}^0 -(-x) dx + \int_0^1 -x dx\right] \\&= \left(\frac{1}{2}\right) \left[\int_{-1}^0 x dx + \int_0^1 -x dx\right] \\&= \left(\frac{1}{2}\right) \left[\frac{1}{2}x^2 \Big|_{-1}^0 - \frac{1}{2}x^2 \Big|_0^1\right] \\&= \left(\frac{1}{2}\right) \left[\left(0 - \left(\frac{1}{2}\right)\right) - \left(\frac{1}{2} - 0\right)\right] \\&= \left(\frac{1}{2}\right) \left[-\frac{1}{2} - \frac{1}{2}\right] \\&= \left(\frac{1}{2}\right) (-1) \\&= -\frac{1}{2}\end{aligned}$$

7. (a) Consider the partition P that subdivides the interval $[a, b]$ into n subintervals of width $\Delta x = \frac{b-a}{n}$ and let c_k be the right endpoint of each subinterval. So the partition is

$$P = \left\{a, a + \frac{b-a}{n}, a + \frac{2(b-a)}{n}, \dots, a + \frac{n(b-a)}{n} = b\right\}$$

and $c_k = a + \frac{k(b-a)}{n}$. We get the Riemann sum

$$\begin{aligned}
\sum_{k=1}^n f(c_k)\Delta x &= \sum_{k=1}^n c_k^2 \left(\frac{b-a}{n} \right) \\
&= \frac{b-a}{n} \sum_{k=1}^n \left(a + \frac{k(b-a)}{n} \right) \\
&= \frac{b-a}{n} \sum_{k=1}^n \left(a^2 + \frac{2ak(b-a)}{n} + \frac{k^2(b-a)^2}{n^2} \right) \\
&= \frac{b-a}{n} \left(\sum_{k=1}^n a^2 + \frac{2a(b-a)}{n} \sum_{k=1}^n k + \frac{(b-a)^2}{n^2} \sum_{k=1}^n k^2 \right) \\
&= \frac{b-a}{n} \cdot na^2 + \frac{2a(b-a)^2}{n^2} \cdot \frac{n(n+1)}{2} + \frac{(b-a)^3}{n^3} \cdot \frac{n(n+1)(2n+1)}{6} \\
&= (b-a)a^2 + a(b-a)^2 \cdot \frac{n+1}{n} + \frac{(b-a)^3(2n^2+3n+1)}{6n^2} \\
&= (b-a)a^2 + a(b-a)^2 \cdot \left(1 + \frac{1}{n} \right) + (b-a)^3 \left(\frac{2}{6} + \frac{3}{6n} + \frac{1}{6n^2} \right).
\end{aligned}$$

As $n \rightarrow \infty$ and $\|P\| \rightarrow 0$, this expression converges to

$$(b-a)a^2 + a(b-a)^2 + (b-a)^3 \cdot \frac{2}{6},$$

which simplifies to $\frac{b^3}{3} - \frac{a^3}{3}$. Thus $\int_a^b x^2 dx = \frac{b^3}{3} - \frac{a^3}{3}$.

- (b) Consider the partition P that subdivides the interval $[-1, 0]$ into n subintervals of width $\Delta x = \frac{1}{n}$ and let c_k be the right endpoint of each subinterval. So the partition is

$$P = \left\{ -1, -1 + \frac{1}{n}, -1 + \frac{2}{n}, \dots, -1 + \frac{n}{n} = 0 \right\}$$

and $c_k = -1 + \frac{k}{n}$. We get the Riemann sum

$$\begin{aligned}
\sum_{k=1}^n f(c_k)\Delta x &= \sum_{k=1}^n \left(\left(-1 + \frac{k}{n} \right) - \left(-1 + \frac{k}{n} \right)^2 \right) \cdot \frac{1}{n} \\
&= \frac{1}{n} \sum_{k=1}^n \left(-1 + \frac{k}{n} - 1 + \frac{2k}{n} - \left(\frac{k}{n} \right)^2 \right) \\
&= \frac{-2}{n} \sum_{k=1}^n 1 + \frac{3}{n^2} \sum_{k=1}^n k - \frac{1}{n^3} \sum_{k=1}^n k^2 \\
&= \frac{-2}{n}n + \frac{3}{n^2} \left(\frac{n(n+1)}{2} \right) - \frac{1}{n^3} \left(\frac{n(n+1)(2n+1)}{6} \right) \\
&= -2 + \frac{3}{2} \left(1 + \frac{1}{n} \right) + \frac{1}{n^3} \left(\frac{2n^3 + 3n^2 + n}{6n^3} \right).
\end{aligned}$$

As $n \rightarrow \infty$ and $\|P\| \rightarrow 0$, this expression converges to $-2 + \frac{3}{2} - \frac{2}{6} = -\frac{5}{6}$. Thus $\int_{-1}^0 (x - x^2) dx = -\frac{5}{6}$.

8. Use the Max-Min Inequality to find upper and lower bounds for the value of

$$\int_0^1 \frac{1}{1+x^2} dx$$

Notice that the integrand $f(x) = \frac{1}{1+x^2}$ is a decreasing function

($f'(x) = \frac{-2x}{(1+x^2)^2} \leq 0$ for $x \in [0, 1]$). Therefore, on the interval f has a maximum at $x = 0$ and has a minimum at $x = 1$.

$$\max f = f(0) = 1$$

$$\min f = f(1) = \frac{1}{2}$$

Thus the Max-Min Inequality gives:

$$\frac{1}{2} \leq \int_0^1 \frac{1}{1+x^2} dx \leq 1$$

9. Use the Max-Min Inequality to find upper and lower bounds for the the value of

$$\int_0^{0.5} \frac{1}{1+x^2} dx \quad \text{and} \quad \int_{0.5}^1 \frac{1}{1+x^2} dx$$

Using the observation (from 65) that $f(x) = \frac{1}{1+x^2}$ is decreasing on $[0, 0.5]$ and $[0.5, 1]$, we get the following two inequalities:

$$0.4 \leq \int_0^{0.5} \frac{1}{1+x^2} dx \leq 0.5$$

$$0.25 \leq \int_{0.5}^1 \frac{1}{1+x^2} dx \leq 0.4.$$

Therefore, by adding the two inequalities we get the new (and improved) estimate:

$$0.65 \leq \int_0^1 \frac{1}{1+x^2} dx \leq 0.9.$$

10. We know that $\sin x \leq x$ for $x \geq 0$. Therefore we can get the upper bound:

$$\begin{aligned}
\int_0^1 \sin x \, dx &\leq \int_0^1 x \, dx \\
&= \frac{1}{2} \cdot 1^2 - \frac{1}{2} \cdot 0^2 \\
&= \frac{1}{2}.
\end{aligned}$$

Section 5.4:

1. (a) $\int_{-2}^0 (2x + 5) \, dx = (x^2 + 5x)|_{-2}^0 = (0 + 0) - (4 - 10) = 6$
- (b) $\int_{-3}^4 \left(5 - \frac{x}{2}\right) \, dx = \left(5x - \frac{x^2}{4}\right)\Big|_{-3}^4 = (20 - 4) - \left(-15 - \frac{9}{4}\right) = 16 + \frac{69}{4} = \frac{133}{4}$
- (c) $\int_0^\pi \sin x \, dx = -\cos x\Big|_0^\pi = -\cos(\pi) + \cos(0) = 1 + 1 = 2$
- (d) $\int_0^{\pi/3} 2 \sec^2 x \, dx = 2 \tan x\Big|_0^{\pi/3} = 2 \tan(\pi/3) - 2 \tan(0) = 2\sqrt{3} - 2 \cdot 0 = 2\sqrt{3}$
- (e) $\int_1^{-1} (r+1)^2 \, dr = -\int_{-1}^1 (r^2 + 2r + 1) \, dr = -\left[\left(\frac{1}{3}r^3 + r^2 + r\right)\Big|_{-1}^1\right] = -\left[\left(\frac{1}{3} + 1 + 1\right) - \left(-\frac{1}{3} + 1 - 1\right)\right] = -\left[\frac{7}{3} + \frac{1}{3}\right] = -\frac{8}{3}$
- (f) $\int_1^2 \left(\frac{1}{x} - e^{-x}\right) \, dx = (\ln x + e^{-x})\Big|_1^2 = (\ln 2 + e^{-2}) - (0 + e^{-1}) = \ln 2 + \frac{1}{e^2} - \frac{1}{e}$
- (g) $\int_0^1 \frac{4}{1+x^2} \, dx = 4 \arctan x\Big|_0^1 = 4 \arctan(1) - 4 \arctan(0) = 4 \cdot \frac{\pi}{4} - 4 \cdot 0 = \pi$
- (h) $\int_2^5 \frac{x}{\sqrt{1+x^2}} \, dx = \int_2^5 x(1+x^2)^{-1/2} \, dx = \frac{1}{2} \cdot 2 \cdot \sqrt{1+x^2}\Big|_2^5 = \sqrt{1+x^2}\Big|_2^5 = \sqrt{26} - \sqrt{5}$

$$(i) \int_0^1 x e^{x^2} dx$$

Notice that $\frac{d}{dx}x^2 = 2x$, and we have an x^2 in the exponent and an x in the product. Thus, keeping in mind the chain rule we get:

$$f(x) = x e^{x^2} \quad \Rightarrow \quad F(x) = \frac{1}{2} e^{x^2}$$

$$\int_0^1 x e^{x^2} dx = F(1) - F(0) = \frac{1}{2} e - \frac{1}{2} = \frac{1}{2}(e - 1).$$

$$(j) \int_1^2 \frac{\ln x}{x} dx$$

Notice that $\frac{d}{dx} \ln x = \frac{1}{x}$, and we have an $\ln x$ and a $\frac{1}{x}$. Thus, keeping in mind the chain rule we get:

$$f(x) = \frac{\ln x}{x} \quad \Rightarrow \quad F(x) = \frac{1}{2} (\ln x)^2$$

$$\int_1^2 \frac{\ln x}{x} dx = F(2) - F(1) = \frac{1}{2} (\ln 2)^2 - 0 = \frac{1}{2} (\ln 2)^2.$$

$$2. (a) \frac{d}{dx} \int_0^{\sqrt{x}} \cos t dt$$

i.

$$\begin{aligned} \int_0^{\sqrt{x}} \cos t dt &= \sin t \Big|_0^{\sqrt{x}} \\ &= \sin(\sqrt{x}) - \sin(0) \\ &= \sin(\sqrt{x}) \end{aligned}$$

Therefore we get:

$$\begin{aligned} \frac{d}{dx} \int_0^{\sqrt{x}} \cos t dt &= \frac{d}{dx} \sin(\sqrt{x}) \\ &= \cos(\sqrt{x}) \cdot \frac{1}{2\sqrt{x}} \end{aligned}$$

ii. Let $u = \sqrt{x}$

$$\begin{aligned} \frac{d}{dx} \int_0^{\sqrt{x}} \cos t dt &= \left(\frac{d}{du} \int_0^u \cos t dt \right) \cdot \frac{du}{dx} \\ &= \cos u \cdot \frac{du}{dx} \\ &= \cos(\sqrt{x}) \cdot \frac{1}{2\sqrt{x}} \end{aligned}$$

$$(b) \frac{d}{d\theta} \int_0^{\tan \theta} \sec^2 y \, dy$$

i.

$$\begin{aligned} \int_0^{\tan \theta} \sec^2 y \, dy &= \tan y \Big|_0^{\tan \theta} \\ &= \tan(\tan \theta) - \tan(0) \\ &= \tan(\tan \theta) \end{aligned}$$

Therefore we get:

$$\begin{aligned} \frac{d}{d\theta} \int_0^{\tan \theta} \sec^2 y \, dy &= \frac{d}{d\theta} \tan(\tan \theta) \\ &= \sec^2(\tan \theta) \cdot \sec^2 \theta \end{aligned}$$

ii. Let $u = \tan \theta$

$$\begin{aligned} \frac{d}{d\theta} \int_0^{\tan \theta} \sec^2 y \, dy &= \left(\frac{d}{du} \int_0^u \sec^2 y \, dy \right) \cdot \frac{du}{d\theta} \\ &= \sec^2 u \cdot \frac{du}{d\theta} \\ &= \sec^2(\tan \theta) \cdot \sec^2 \theta \end{aligned}$$

$$3. \, y = \int_{\sqrt{x}}^0 \sin(t^2) \, dt = - \int_0^{\sqrt{x}} \sin(t^2) \, dt$$

Let $u = \sqrt{x}$. Then we get:

$$\frac{dy}{dx} = \frac{dy}{du} \cdot \frac{du}{dx} = -\sin(u^2) \cdot \frac{du}{dx} = -\sin(x) \cdot \frac{1}{x\sqrt{x}} = -\frac{\sin(x)}{2\sqrt{x}}$$

$$4. \, (a) \, y = -x^2 - 2x \quad \text{where} \quad -3 \leq x \leq 2$$

First, notice that $y = -x^2 - 2x = -x(x+2)$. Therefore in the interval $[-3, 2]$, the function has zeroes at $x = 0$ and $x = -2$.

$$\begin{aligned} \text{Area} &= - \int_{-3}^{-2} (-x^2 - 2x) \, dx + \int_{-2}^0 (-x^2 - 2x) \, dx - \int_0^2 (-x^2 - 2x) \, dx \\ &= - \left(-\frac{1}{3}x^3 - x^2 \right) \Big|_{-3}^{-2} + \left(-\frac{1}{3}x^3 - x^2 \right) \Big|_{-2}^0 - \left(-\frac{1}{3}x^3 - x^2 \right) \Big|_0^2 \\ &= - \left[\left(\frac{8}{3} - 4 \right) - (9 - 9) \right] + \left[(0 - 0) - \left(\frac{8}{3} - 4 \right) \right] - \left[\left(-\frac{8}{3} - 4 \right) - (0 - 0) \right] \\ &= \frac{4}{3} + \frac{4}{3} + \frac{20}{3} \\ &= \frac{28}{3} \end{aligned}$$

(b) $y = 3x^2 - 3$ where $-2 \leq x \leq 2$

First notice that $y = 3(x^2 - 1) = 3(x - 1)(x + 1)$. Therefore in the interval $[-2, 2]$, the function has zeroes at $x = \pm 1$.

$$\begin{aligned} \text{Area} &= \int_{-2}^{-1} (3x^2 - 3) dx - \int_{-1}^1 (3x^2 - 2) dx + \int_1^2 (3x^2 - 3) dx \\ &= (x^3 - 3x)|_{-2}^{-1} - (x^3 - 3x)|_{-1}^1 + (x^3 - 3x)|_1^2 \\ &= [(-1 + 3) - (-8 + 6)] - [(1 - 3) - (-1 + 3)] + [(8 - 6) - (1 - 3)] \\ &= 4 + 4 + 4 \\ &= 12 \end{aligned}$$

(c) $y = x^{1/3} - x$ where $-1 \leq x \leq 8$

First notice that $y = x^{1/3} - x = x^{1/3}(1 - x^{2/3})(1 + x^{1/3})$, so the on $[-1, 8]$ the function has zeroes at $x = 0, \pm 1$.

$$\begin{aligned} \text{Area} &= - \int_{-1}^0 (x^{1/3} - x) dx + \int_0^1 (x^{1/3} - x) dx - \int_1^8 (x^{1/3} - x) dx \\ &= - \left(\frac{3}{4}x^{4/3} - \frac{1}{2}x^2 \right) \Big|_{-1}^0 + \left(\frac{3}{4}x^{4/3} - \frac{1}{2}x^2 \right) \Big|_0^1 - \left(\frac{3}{4}x^{4/3} - \frac{1}{2}x^2 \right) \Big|_1^8 \\ &= - \left[(0 - 0) - \left(\frac{3}{4} - \frac{1}{2} \right) \right] + \left[\left(\frac{3}{4} - \frac{1}{2} \right) - (0 - 0) \right] - \left[(12 - 32) - \left(\frac{3}{4} - \frac{1}{2} \right) \right] \\ &= \frac{1}{4} + \frac{1}{4} + \frac{81}{4} \\ &= \frac{83}{4} \end{aligned}$$

5. (a) $v = \frac{ds}{dt} = \frac{d}{dt} \int_0^t f(x) dx = f(t)$.

Therefore $v(5) = f(5) = 2$ m/s.

(b) Recall that the acceleration is the rate of change of the velocity, so $a = \frac{dv}{dt}$. Since f has a negative slope at $t = 5$, the acceleration is negative at $t = 5$.

(c) Since $s = \int_0^t f(x) dx$, we know that the position at $t = 3$ can be represented as the area under the graph on $[0, 3]$. Notice that on this interval, f is a straight line and so the value of the integral is just the area of a triangle formed by $f(x)$, the x -axis and $x = 3$.

Therefore $s(3) = \frac{1}{2} \cdot 3 \cdot 3 = \frac{9}{2}$

(d) We want to know when s has a maximum and we will determine this by looking at $f(x)$, which is the derivative of the s . We know f is positive on $[0, 6]$ and negative on $[6, 9]$, which means that s goes from increasing to decreasing at $t = 6$. By the first derivative test, we know that s must have relative max at $t = 6$.

(e) At $x \approx 4$ and $x \approx 7$ since $s''(x) = f'(x)$ is zero at these points.

(f) The particle is moving towards the origin when it has negative velocity $\rightarrow f$ is negative. Hence it is moving towards the origin on $[6, 9]$. Similarly the particle is moving away from the origin when f is positive, hence on $[0, 6]$.

(g) Judging by the graph of f , we know that the particle is to the right of the origin (on the positive side) because the $s(9) = \int_0^9 f(x) dx > 0$ (the area above the x -axis is greater than the area below it).

6. Find $\lim_{x \rightarrow 0} \frac{1}{x^3} \int_0^x \frac{t^2}{t^4 + 1} dt$

Notice that $\lim_{x \rightarrow 0} \int_0^x \frac{t^2}{t^4 + 1} dt = 0$ by the rules of integrals and $\lim_{x \rightarrow 0} x^3 = 0$. Thus if you think of the expression inside the limit as a fraction, we have that both the numerator and the denominator go to 0. Hence we can apply l'Hôpital's Rule to the limit and get:

$$\begin{aligned} \lim_{x \rightarrow 0} \frac{1}{x^3} \int_0^x \frac{t^2}{t^4 + 1} dt &= \lim_{x \rightarrow 0} \left[\left(\int_0^x \frac{t^2}{t^4 + 1} dt \right) / x^3 \right] \\ &= \lim_{x \rightarrow 0} \left[\left(\frac{x^2}{x^4 + 1} \right) / (3x^2) \right] \\ &= \lim_{x \rightarrow 0} \frac{1}{3(x^4 + 1)} \\ &= \frac{1}{3} \end{aligned}$$

7. Suppose $f'(x) \geq 0$ for all values of x , and that $f(1) = 0$. Define:

$$g(x) = \int_0^x f(t) dt$$

(a) True. It follows from the FTC, part I.

(b) True. Differentiable \Rightarrow Continuous.

- (c) True. We know that $g'(1) = f(1) = 0$, so the tangent line at $x = 0$ is horizontal.
- (d) False. We know that f crosses the x -axis at $x = 1$ and that f is always increasing. Therefore $g'(x) = f(x)$ is negative for $x < 1$ and $g'(x) = f(x)$ is positive for $x > 1$. By the first derivative test, this is a minimum, not a max.

Note that you could also use the fact that $g'(1) = 0$ and $g''(1) > 0$

- (e) True. See (d.)
- (f) False. $g''(x) = f'(x) > 0$, and hence does not change sign.
- (g) True. $g'(x) = f(x)$ and we know that the $f(1) = 0$ and is increasing.