

SPECTRAL CURVES AND THE SCHRÖDINGER EQUATIONS FOR THE EYNARD-ORANTIN RECURSION

MOTOHICO MULASE AND PIOTR SULKOWSKI

ABSTRACT. It is predicted that the principal specialization of the partition function of a B-model topological string theory, that is mirror dual to an A-model enumerative geometry problem, satisfies a Schrödinger equation, and that the characteristic variety of the Schrödinger operator gives the spectral curve of the B-model theory, when an algebraic K-theory obstruction vanishes. In this paper we present two concrete mathematical A-model examples whose mirror dual partners exhibit these predicted features on the B-model side. The A-model examples we discuss are the generalized Catalan numbers of an arbitrary genus and the single Hurwitz numbers. In each case, we show that the Laplace transform of the counting functions satisfies the Eynard-Orantin topological recursion, that the B-model partition function satisfies the KP equations, and that the principal specialization of the partition function satisfies a Schrödinger equation whose total symbol is exactly the Lagrangian immersion of the spectral curve of the Eynard-Orantin theory.

CONTENTS

1. Introduction and the main results	1
2. The Eynard-Orantin topological recursion	6
3. The generalized Catalan numbers and the topological recursion	10
4. The partition function for the generalized Catalan numbers and the Schrödinger equation	13
5. Single Hurwitz numbers	17
6. The Schur function expansion of the Hurwitz partition function	24
7. Conclusion	30
Appendix A. Proof of the Schrödinger equation for the Catalan case	31
References	34

1. INTRODUCTION AND THE MAIN RESULTS

In a series of remarkable papers of Mariño [51] and Bouchard, Klemm, Mariño, and Pasquetti [8], these authors have developed an inductive mechanism to calculate a variety of quantum invariants and solutions to enumerative geometry questions, based on the fundamental work of Eynard and Orantin [27, 28, 29]. The validity of their method, known as the *remodeled B-model* based on the *topological recursion* of Eynard-Orantin, has been established for many different enumerative geometry problems, such as single Hurwitz numbers ([7, 26, 60], based on the conjecture of Bouchard and Mariño [9]), open Gromov-Witten invariants of smooth toric Calabi-Yau threefolds ([29, 80], based on the *remodeling conjecture* of Mariño [51] and Bouchard, Klemm, Mariño, Pasquetti [8]), and the number of lattice points on $\mathcal{M}_{g,n}$ and its symplectic and Euclidean volumes ([11, 20, 58], based on [63, 64]). It is expected that double Hurwitz numbers, stationary Gromov-Witten invariants of \mathbb{P}^1

2000 *Mathematics Subject Classification*. Primary: 14H15, 14N35, 05C30, 11P21; Secondary: 81T30.

[65, 66], certain Donaldson-Thomas invariants, and many other quantum invariants would also fall into this category.

Unlike the familiar *Topological Recursion Relations* (TRR) of the Gromov-Witten theory, the Eynard-Orantin recursion is a B-model formula [8, 51]. The significant feature of the formula is its universality: independent of the A-model problem, the B-model recursion takes always the same form. The input data of this B-model consist of a holomorphic Lagrangian immersion

$$\begin{array}{ccc} \iota : \Sigma & \longrightarrow & T^*\mathbb{C} \\ & & \downarrow \pi \\ & & \mathbb{C} \end{array}$$

of an open Riemann surface Σ (called a *spectral curve* of the Eynard-Orantin recursion) into the cotangent bundle $T^*\mathbb{C}$ equipped with the tautological 1-form η , and the symmetric second derivative of the logarithm of *Riemann's prime form* [30, 62] defined on $\Sigma \times \Sigma$. The procedure of Eynard-Orantin [27] then defines, inductively on $2g - 2 + n$, a meromorphic symmetric differential n -form $W_{g,n}$ on Σ^n for every $g \geq 0$ and $n \geq 1$ subject to $2g - 2 + n > 0$. A particular choice of the Lagrangian immersion gives a different $W_{g,n}$, which then gives a generating function of the solution to a different enumerative geometry problem.

Thus the real question is how to find the right Lagrangian immersion from a given A-model.

Suppose we have a solution to an enumerative geometry problem (an A-model problem). Then we know a generating function of these quantities. In [20] we proposed an idea of identifying the spectral curve Σ , which states that the spectral curve is the *Laplace transform* of the disc amplitude of the A-model problem. Here the Laplace transform plays the role of mirror symmetry. Thus we obtain a Riemann surface Σ . Still we do not see the aspect of the Lagrangian immersion in this manner.

Every curve in $T^*\mathbb{C}$ is trivially a Lagrangian. But not every Lagrangian is realized as the mirror dual to an A-model problem. The obstruction seems to lie in the K-group $K_2(\mathbb{C}(\Sigma)) \otimes \mathbb{Q}$. When this obstruction vanishes, we call Σ a *K_2 -Lagrangian*, following Kontsevich's terminology. For a K_2 -Lagrangian Σ , we expect the existence of a *holonomic system* that characterizes the *partition function* of the B-model theory, and at the same time, the characteristic variety of this holonomic system recovers the spectral curve Σ as the Lagrangian immersion. A generator of this holonomic system is called a *quantum Riemann surface* [1, 16, 17], because it is a differential operator whose total symbol is the spectral curve realized as a Lagrangian immersion [18]. It is the work of Gukov and Sułkowski [39] that suggested the obstruction to the existence of the holonomic system with algebraic K-theory as an element of K_2 .

Another mysterious link of the Eynard-Orantin theory is its relation to integrable systems of the KP/KdV type [5, 27]. We note that the partition function of the B-model is always the *principal specialization* of a τ -function of the KP equations for all the examples we know by now.

The purpose of the present paper is to give the simplest non-trivial mathematical examples of the theory that exhibit these key features mentioned above. With these examples one can calculate all quantities involved, give proofs of the statements predicted in physics, and examine the mathematical structure of the theory. Our examples are based on enumeration problems of branched coverings of \mathbb{P}^1 .

The idea of *homological mirror symmetry* of Kontsevich [44] allows us to talk about the mirror symmetry *without* underlying spaces, because the formulation is based on the

derived equivalence of categories. Therefore, we can consider the mirror dual B-models corresponding to the enumeration problems of branched coverings on the A-model side. At the same time, being the *derived* equivalence, the homological mirror symmetry does not tell us any direct relations between the quantum invariants on the A-model side and the complex geometry on the B-model side. This is exactly where Mariño's idea of *remodeling* B-model comes in for rescue. The remodeled B-model of [8, 51] is not a derived category of coherent sheaves. Although its applicability is restricted to the case when there is a family of curves Σ that exhibits the geometry of the B-model, the new idea is to construct a network of inter-related differential forms on the symmetric powers of Σ via the Eynard-Orantin recursion, and to understand this infinite system as *the B-model*. The advantage of this idea is that we can relate the solution of the geometric enumeration problem on the A-model side and the symmetric differential forms on the B-model side through the *Laplace transform*. In this sense we consider the Laplace transform as a mirror symmetry.

The first example we consider in this paper is the generalized Catalan numbers of an arbitrary genus. This is equivalent to the “ $c = 1$ model” of [39, Section 5]. In terms of enumeration, we are counting the number of algebraic curves defined over $\overline{\mathbb{Q}}$ in a systematic way by using the dual graph of Grothendieck's *dessins d'enfants* [4, 74].

Let $D_{g,n}(\mu_1, \dots, \mu_n)$ denote the automorphism-weighted count of the number of connected cellular graphs on a closed oriented surface of genus g (i.e., the 1-skeleton of cell-decompositions of the surface), with n labeled vertices of degrees (μ_1, \dots, μ_n) . The letter D stands for ‘dessin.’ The *generalized Catalan numbers* of type (g, n) are defined by

$$C_{g,n}(\mu_1, \dots, \mu_n) = \mu_1 \cdots \mu_n D_{g,n}(\mu_1, \dots, \mu_n).$$

While $D_{g,n}(\vec{\mu})$ is a rational number due to the graph automorphisms, the generalized Catalan number $C_{g,n}(\vec{\mu})$ is always a non-negative integer. It gives the dimension of a linear skein space. In particular, the $(g, n) = (0, 1)$ case recovers the original Catalan numbers:

$$C_{0,1}(2m) = C_m = \frac{1}{m+1} \binom{2m}{m} = \dim \text{End}_{\mathcal{U}_q(\mathfrak{sl}_2)}(T^{\otimes m} \mathbb{C}^2).$$

As explained in [20], the mirror dual to the Catalan numbers C_m is the plane curve Σ defined by

$$(1.1) \quad \begin{cases} x = z + \frac{1}{z} \\ y = -z, \end{cases}$$

where

$$(1.2) \quad z(x) = \sum_{m=0}^{\infty} C_m \frac{1}{x^{2m+1}}.$$

Note that (1.1) also gives a Lagrangian immersion $\Sigma \rightarrow T^*\mathbb{C}$. Let us introduce the *free energies* by

$$(1.3) \quad F_{g,n}^C(z(x_1), \dots, z(x_n)) = \sum_{\vec{\mu} \in \mathbb{Z}_+^n} D_{g,n}(\vec{\mu}) e^{-(w_1 \mu_1 + \dots + w_n \mu_n)} = \sum_{\vec{\mu} \in \mathbb{Z}_+^n} D_{g,n}(\vec{\mu}) \prod_{i=1}^n \frac{1}{x_i^{\mu_i}}$$

as the Laplace transform of the number of dessins, where the coordinates are related by (1.2) and $x_i = e^{w_i}$. The free energy $F_{g,n}^C(z_1, \dots, z_n)$ is a symmetric function in n -variables, and its *principal specialization* is defined by $F_{g,n}^C(z, \dots, z)$. Now let

$$W_{g,n}^C(z_1, \dots, z_n) = d_1 \cdots d_n F_{g,n}^C(z_1, \dots, z_n).$$

It is proved in [20] that $W_{g,n}^C$'s satisfy the Eynard-Orantin topological recursion.

The *Catalan partition function* is given by the formula of [27]:

$$(1.4) \quad Z^C(z, \hbar) = \exp \left(\sum_{g=0}^{\infty} \sum_{n=1}^{\infty} \frac{1}{n!} \hbar^{2g-2+n} F_{g,n}^C(z, z, \dots, z) \right).$$

In this paper we prove

Theorem 1.1. *The Catalan partition function satisfies the Schrödinger equation*

$$(1.5) \quad \left(\hbar^2 \frac{d^2}{dx^2} + \hbar x \frac{d}{dx} + 1 \right) Z^C(z(x), \hbar) = 0.$$

The characteristic variety of this ordinary differential operator, $y^2 + xy + 1 = 0$ for every fixed choice of \hbar , is exactly the Lagrangian immersion (1.1), where we identify the xy -plane as the cotangent bundle $T^*\mathbb{C}$ with the fiber coordinate $y = \hbar \frac{d}{dx}$.

Remark 1.2. A purely geometric reason of our interest in the function appearing as the principal specialization $F_{g,n}^C(z, \dots, z)$ is that, in the stable range $2g - 2 + n > 0$, it is a *polynomial* in

$$(1.6) \quad s = \frac{z^2}{z^2 - 1}$$

of degree $6g - 6 + 3n$. It is indeed the virtual Poincaré polynomial of $\mathcal{M}_{g,n} \times \mathbb{R}_+^n$ [58], and its special value at $s = 1$ gives the Euler characteristic $(-1)^n \chi(\mathcal{M}_{g,n})$ of the moduli space $\mathcal{M}_{g,n}$ of smooth n -pointed curves of genus g . Thus $Z^C(z, \hbar)$ is the exponential generating function of the virtual Poincaré polynomials of $\mathcal{M}_{g,n} \times \mathbb{R}_+^n$.

As such, the generating function $Z^C(z, \hbar)$ is also expressible in terms of a Hermitian matrix integral

$$(1.7) \quad Z^C(z, \hbar) = \int_{\mathcal{H}_{N \times N}} \det(1 - \sqrt{s}X)^N e^{-\frac{N}{2} \text{trace}(X^2)} dX$$

with the identification (1.6) and $\hbar = 1/N$. Here dX is the normalized Lebesgue measure on the space of $N \times N$ Hermitian matrices $\mathcal{H}_{N \times N}$. It is a well-known fact that Eq.(1.7) is the principal specialization of a KP τ -function (see for example, [54]).

Another example we consider in this paper is based on single Hurwitz numbers. As a counting problem it is easier to state than the previous example, but the Lagrangian immersion requires a transcendental function, and hence the resulting Schrödinger equation exhibits a rather different nature.

Let $H_{g,n}(\mu_1, \dots, \mu_n)$ be the automorphism-weighted count of the number of topological types of Hurwitz covers $f : C \rightarrow \mathbb{P}^1$ of a connected non-singular algebraic curve C of genus g . A holomorphic map f is a *Hurwitz cover* of *profile* (μ_1, \dots, μ_n) if it has n labeled poles of orders (μ_1, \dots, μ_n) and is simply ramified otherwise. Introduce the Laplace transform of single Hurwitz numbers by

$$(1.8) \quad F_{g,n}^H(t(w_1), \dots, t(w_n)) = \sum_{\vec{\mu} \in \mathbb{Z}_+^n} H_{g,n}(\vec{\mu}) e^{-(w_1 \mu_1 + \dots + w_n \mu_n)},$$

where

$$t(w) = \sum_{m=0}^{\infty} \frac{m^m}{m!} e^{-mw}$$

is the tree-function. Here again $F_{g,n}^H(t_1, \dots, t_n)$ is a polynomial of degree $6g - 6 + 3n$ if $2g - 2 + n > 0$ [60]. Bouchard and Mariño have conjectured [9] that

$$W_{g,n}^H(t_1, \dots, t_n) = d_1 \cdots d_n F_{g,n}^H(t_1, \dots, t_n)$$

satisfy the Eynard-Orantin topological recursion, with respect to the Lagrangian immersion

$$(1.9) \quad \begin{cases} x = e^{-w} = ze^{-z} \in \mathbb{C}^* \\ y = z \in \mathbb{C}, \end{cases}$$

where

$$z = \frac{t-1}{t}$$

and we use $\eta = y \frac{dx}{x}$ as the tautological 1-form on $T^*\mathbb{C}^*$. The Bouchard-Mariño conjecture was proved in [7, 26, 60].

Now we define the *Hurwitz partition function*

$$(1.10) \quad Z^H(t, \hbar) = \exp \left(\sum_{g=0}^{\infty} \sum_{n=1}^{\infty} \frac{1}{n!} \hbar^{2g-2+n} F_{g,n}^H(t, \dots, t) \right).$$

Then we have the following

Theorem 1.3. *The Hurwitz partition function satisfy two equations:*

$$(1.11) \quad \left[\frac{1}{2} \hbar \frac{\partial^2}{\partial w^2} + \left(1 + \frac{1}{2} \hbar \right) \frac{\partial}{\partial w} - \hbar \frac{\partial}{\partial \hbar} \right] Z^H(t(w), \hbar) = 0,$$

$$(1.12) \quad \left(\hbar \frac{d}{dw} + e^{-w} e^{-\hbar \frac{d}{dw}} \right) Z^H(t(w), \hbar) = 0.$$

Moreover, each of the two equations recover the Lagrangian immersion (1.9) from the asymptotic analysis at $\hbar \sim 0$. And if we view \hbar as a fixed constant in (1.12), then its total symbol is the Lagrangian immersion (1.9) with the identification $z = -\hbar \frac{d}{dw}$.

Remark 1.4. The second order equation (1.11) is a consequence of the polynomial recursion of [60]. This situation is exactly the same as Theorem 1.1. The differential-difference equation (1.12), or a *delay* differential equation, follows from the principal specialization of the KP τ -function that gives another generating function of single Hurwitz numbers [41, 67]. We remark that (1.12) is also derived in [82]. The point of view of differential-difference equation is further developed in [59] for the case of double Hurwitz numbers and r -spin structures, where we generalize a result of [82].

Remark 1.5. Define two operators by

$$(1.13) \quad P = \hbar \frac{d}{dw} + e^{-w} e^{-\hbar \frac{d}{dw}} \quad \text{and}$$

$$(1.14) \quad Q = \frac{1}{2} \hbar \frac{\partial^2}{\partial w^2} + \left(1 + \frac{1}{2} \hbar \right) \frac{\partial}{\partial w} - \hbar \frac{\partial}{\partial \hbar}.$$

Then it is noted in [49] that

$$(1.15) \quad [P, Q] = P.$$

Thus the *heat equation* (1.11) preserves the space of solutions of the *Schrödinger equation* (1.12). In this sense, (1.12) is holonomic for every fixed \hbar . The analysis of these equations is further investigated in [49].

The existence of a holonomic system is particularly appealing when we consider the knot A-polynomial as the defining equation of a Lagrangian immersion, in connection to the AJ conjecture [33, 34, 38]. One can ask:

Question 1.6. *Let K be a knot in S^3 and A_K its A-polynomial [13]. Is there a concrete formula for the quantum knot invariants of K , such as the colored Jones polynomials, in terms of A_K ?*

The B-model developed in [15], [39], and more recently in [6], clearly shows that the answer is yes, and it should be given by the Eynard-Orantin formalism. Although our examples in the present paper are not related to any knots, they suggest the existence of a corresponding A-model. An interesting theory of generalized A-polynomials and quantum knot invariants from the point of view of mirror symmetry is recently presented in Aganagic and Vafa [2]. We also remark that there are further developments in this direction [10, 31, 32].

The paper is organized as follows. In Section 2, we give the definition of the Eynard-Orantin topological recursion. We emphasize the aspect of Lagrangian immersion in our presentation. In Section 3 we review the generalized Catalan numbers of [20]. Then in Section 4 we derive the Schrödinger equation for the Catalan partition function. The equation for the Hurwitz partition function is given in Section 5. Finally, in Section 6 we give the proof of (1.12) using the Schur function expansion of the Hurwitz generating function and its principal specialization.

2. THE EYNARD-ORANTIN TOPOLOGICAL RECURSION

We adopt the following definition for the topological recursion of Eynard-Orantin [27]. Our emphasis, which is different from the original, is the point of view of the Lagrangian immersion.

Definition 2.1. The **spectral curve** (Σ, ι) consists of an open Riemann surface Σ and a *Lagrangian immersion*

$$(2.1) \quad \begin{array}{ccc} \iota : \Sigma & \longrightarrow & T^*\mathbb{C} \\ & & \downarrow \pi \\ & & \mathbb{C} \end{array}$$

with respect to the canonical holomorphic symplectic structure $\omega = -d\eta$ on $T^*\mathbb{C}$, where η is the tautological 1-form on the cotangent bundle $\pi : T^*\mathbb{C} \rightarrow \mathbb{C}$. Recall that $p \in \Sigma$ is a *Lagrangian singularity* if $d(\pi \circ \iota) = 0$ at p , and that $\pi(\iota(p)) \in \mathbb{C}$ is a *caustic* of the Lagrangian immersion. We assume that the projection π restricted to the Lagrangian immersion is locally simply ramified around each Lagrangian singularity. We denote by $R = \{p_1, \dots, p_r\} \subset \Sigma$ the set of Lagrangian singularities, and by

$$U = \sqcup_{j=1}^r U_j$$

the disjoint union of small neighborhood U_j around p_j such that $\pi : U_j \rightarrow \pi(U_j) \subset \mathbb{C}$ is a double-sheeted covering ramified only at p_j . We denote by $s_j(z)$ the local Galois conjugate of $z \in U_j$.

Another key ingredient of the Eynard-Orantin theory is the **normalized fundamental differential of the second kind** $B_\Sigma(z_1, z_2)$ [30, Page 20], [62, Page 3.213], which is a symmetric differential 2-form on $\Sigma \times \Sigma$ with second-order poles along the diagonal. To define it, let us recall a few basic facts about algebraic curves. Let C be a nonsingular

complete algebraic curve over \mathbb{C} . We are considering a nonsingular compactification $C = \overline{\Sigma}$ of the Riemann surface Σ . We identify the Jacobian variety of C as $Jac(C) = Pic^0(C)$, which is isomorphic to $Pic^{g-1}(C)$. The *theta divisor* Θ of $Pic^{g-1}(C)$ is defined by

$$\Theta = \{L \in Pic^{g-1}(C) \mid \dim H^1(C, L) > 0\}.$$

We use the same notation for the translate divisor on $Jac(C)$, also called the theta divisor. Now consider the diagram

$$\begin{array}{ccc} & Jac(C) & \\ & \uparrow \delta & \\ & C \times C & \\ \swarrow pr_1 & & \searrow pr_2 \\ C & & C, \end{array}$$

where pr_j denotes the projection to the j -th components, and

$$\delta : C \times C \ni (p, q) \mapsto p - q \in Jac(C).$$

Then the *prime form* $E_C(z_1, z_2)$ [30, Page 16] is defined as a holomorphic section

$$E_C(p, q) \in H^0 \left(C \times C, pr_1^* K_C^{-\frac{1}{2}} \otimes pr_2^* K_C^{-\frac{1}{2}} \otimes \delta^*(\Theta) \right),$$

where K_C is the canonical line bundle of C and we choose Riemann's spin structure (or the Szegő kernel) $K_C^{\frac{1}{2}}$ (see [30, Theorem 1.1]). We do not need the precise definition of the prime form here, but its characteristic properties are important:

- (1) $E_C(p, q)$ vanishes only along the diagonal $\Delta \subset C \times C$, and has simple zeros along Δ .
- (2) Let z be a local coordinate on C such that $dz(p)$ gives the local trivialization of K_C around p . When q is near at p , $\delta^*(\Theta)$ is also trivialized around $(p, q) \in C \times C$, and we have a local behavior

$$(2.2) \quad E_C(z(p), z(q)) = \frac{z(p) - z(q)}{\sqrt{dz(p)} \cdot \sqrt{dz(q)}} (1 + O((z(p) - z(q))^2)).$$

$$(3) \quad E_C(z(p), z(q)) = -E_C(z(q), z(p)).$$

The fundamental 2-form $B_C(p, q)$ is then defined by

$$(2.3) \quad B_C(p, q) = d_1 \otimes d_2 \log E_C(p, q)$$

(see [30, Page 20], [62, Page 3.213]). We note that $dz(p)$ appears in (2.2) just as the indicator of our choice of the local trivialization. With this local trivialization, the square

$$E_C(p, q)^2 \in H^0 \left(C \times C, pr_1^* K_C^{-1} \otimes pr_2^* K_C^{-1} \otimes \delta^*(\Theta)^{\otimes 2} \right)$$

behaves better because of

$$(2.4) \quad E_C(z(p), z(q))^2 = \frac{(z(p) - z(q))^2}{dz(p) \cdot dz(q)} (1 + O((z(p) - z(q))^2)).$$

We then see that

$$(2.5) \quad B_C(z(p), z(q)) = \frac{1}{2} d_1 \otimes d_2 \log E^2(z(p), z(q))$$

$$= \frac{dz(p) \cdot dz(q)}{(z(p) - z(q))^2} + O(1) dz(p) \cdot dz(q) \in H^0(C \times C, pr_1^* K_C \otimes pr_2^* K_C \otimes \mathcal{O}(2\Delta)).$$

Definition 2.2. Let D be a divisor of Σ . A **meromorphic symmetric differential form** of degree n with poles along D is an element of the symmetric tensor product

$$\text{Sym}^n H^0(\Sigma, K_\Sigma(*D)).$$

Definition 2.3. Meromorphic differential forms

$$W_{g,n}(z_1, \dots, z_n) \in \text{Sym}^n H^0(\Sigma, K_\Sigma(*R))$$

for $g \geq 0$ and $n \geq 1$, subject to $2g - 2 + n > 0$, with poles along the Lagrangian singularities of the Lagrangian immersion $\Sigma \rightarrow T^*\mathbb{C}$, are said to satisfy the **Eynard-Orantin topological recursion** if they satisfy the recursion formula

$$(2.6) \quad W_{g,n}(z_1, z_2, \dots, z_n) = \frac{1}{2\pi i} \sum_{j=1}^r \oint_{\gamma_j} K_j(z, z_1) \left[W_{g-1, n+1}(z, s_j(z), z_2, \dots, z_n) + \sum_{\substack{\text{No } (0,1) \text{ terms} \\ g_1+g_2=g \\ I \sqcup J = \{2,3,\dots,n\}}} W_{g_1, |I|+1}(z, z_I) W_{g_2, |J|+1}(s_j(z), z_J) \right].$$

Here the integration is taken with respect to $z \in U_j$ along a positively oriented simple closed loop γ_j around p_j , and $z_I = (z_i)_{i \in I}$ for a subset $I \subset \{1, 2, \dots, n\}$. In the summation, “No (0, 1) terms” means the summand does not contain the terms with $g_1 = 0$ and $I = \emptyset$ or $g_2 = 0$ and $J = \emptyset$. The recursion kernels $K_j(z, z_1)$, $j = 1, \dots, r$, are defined as follows. First we define $W_{0,1}$ and $W_{0,2}$.

$$(2.7) \quad W_{0,1}(z) = \iota^* \eta = y dx \in H^0(\Sigma, K),$$

where x is a linear coordinate of \mathbb{C} and y is the dual coordinate of $T_0^*\mathbb{C}$ so that the Lagrangian immersion is given by

$$(x, y) : \Sigma \ni t \mapsto (x(z), y(z)) \in T^*\mathbb{C}.$$

$$(2.8) \quad W_{0,2}(z_1, z_2) = B_\Sigma(z_1, z_2) - \pi^* \frac{dx_1 \cdot dx_2}{(x_1 - x_2)^2}.$$

We note that $W_{0,2}(z_1, z_2)$ is holomorphic along the diagonal $z_1 = z_2$. The recursion kernel $K_j(z_1, z_2) \in H^0(U_j \times \Sigma, K_{U_j}^{-1} \otimes K_\Sigma)$ for $z_1 \in U_j$ and $z_2 \in \Sigma$ is defined by

$$(2.9) \quad K_j(z_1, z_2) = \frac{1}{2} \frac{1}{W_{0,1}(s_j(z_1)) - W_{0,1}(z_1)} \otimes \int_{z_1}^{s_j(z_1)} W_{0,2}(\cdot, z_2) = \frac{1}{2} \frac{d_2(\log E_\Sigma(s_j(z_1), z_2) - \log E_\Sigma(z_1, z_2))}{(y(s_j(z_1)) - y(z_1)) dx(z_1)}.$$

The kernel $K_j(z_1, z_2)$ is an algebraic operator that multiplies dz_2 while contracts $\partial/\partial z_1$.

The topological recursion is the reduction of $2g - 2 + n$ by 1, which is different from the usual boundary degeneration formula of $\overline{\mathcal{M}}_{g,n}$. As shown in Figure 2.1, the reduction corresponds to degeneration cycles of codimension 1 and 2, as in Arbarello-Cornalba-Griffiths [3, Chapter 17, Section 5, Page 589].

$$\mathcal{M}_{0,3} \times \overline{\mathcal{M}}_{g,n-1} \rightarrow \overline{\mathcal{M}}_{g,n}, \quad \mathcal{M}_{0,3} \times \overline{\mathcal{M}}_{g-1,n+1} \rightarrow \overline{\mathcal{M}}_{g,n},$$

$$\mathcal{M}_{0,3} \times \bigcup_{\substack{g_1+g_2=g \\ n_1+n_2=n-1}} \overline{\mathcal{M}}_{g_1, n_1+1} \times \overline{\mathcal{M}}_{g_2, n_2+1} \longrightarrow \overline{\mathcal{M}}_{g,n}$$




FIGURE 2.1. The topological recursion and degeneration.

The recursion starts from $W_{1,1}$ and $W_{0,3}$. If we modify (2.6) slightly, then these can also be calculated from $W_{0,2}$ [27].

$$(2.10) \quad W_{1,1}(z_1) = \frac{1}{2\pi i} \sum_{j=1}^r \oint_{\gamma_j} K_j(z, z_1) \left[W_{0,2}(u, v) + c\pi^* \frac{dx(u) \cdot dx(v)}{(x(u) - x(v))^2} \right] \Big|_{u=z, v=s_j(z)}.$$

Formula (2.6) does not give an apparently symmetric expression for $W_{0,3}$. In terms of the coordinate $(x, y) \in T^*\mathbb{C}$ we have an alternative formula for $W_{0,3}$ [27]:

$$(2.11) \quad W_{0,3}(z_1, z_2, z_3) = \frac{1}{2\pi i} \sum_{j=1}^r \oint_{\gamma_j} \frac{W_{0,2}(z, z_1)W_{0,2}(z, z_2)W_{0,2}(z, z_3)}{dx(z) \cdot dy(z)}.$$

Suppose we have a solution $W_{g,n}$ to the topological recursion. A **primitive functions** of the symmetric differential form $W_{g,n}$ is a symmetric meromorphic function $F_{g,n}$ on Σ^n such that its n -fold exterior derivative recovers the $W_{g,n}$, i.e.,

$$(2.12) \quad W_{g,n}(z_1, \dots, z_n) = d_{z_1} \cdots d_{z_n} F_{g,n}(z_1, \dots, z_n).$$

The **partition function** of the topological recursion for a genus 0 spectral curve is the formal expression in infinitely many variables

$$(2.13) \quad Z(z_1, z_2, \dots; \hbar) = \exp \left(\sum_{g \geq 0, n \geq 1} \frac{1}{n!} \hbar^{2g-2+n} F_{g,n}(z_1, z_2, \dots, z_n) \right).$$

The *principal specialization* of the partition function is also denoted by the same letter Z :

$$(2.14) \quad Z(z, \hbar) = \exp \left(\sum_{g \geq 0, n \geq 1} \frac{1}{n!} \hbar^{2g-2+n} F_{g,n}(z, z, \dots, z) \right).$$

Remark 2.4. The partition function depends on the choice of the primitive functions. When we consider the topological recursion as the B-model corresponding to an A-model counting problem, then there is always a canonical choice for the primitives, as the Laplace transform of the quantum invariants.

Remark 2.5. When the spectral curve Σ has a higher genus, the partition function requires a non-perturbative factor in terms of a theta function associated to the curve [5, 6]. In this case the algebraic K-theory condition of [39], probably similar to the *Boutroux condition* of [1], becomes essential for the existence of the quantum curve or the Schrödinger equation. Our consideration in paper is limited to the genus 0 case.

3. THE GENERALIZED CATALAN NUMBERS AND THE TOPOLOGICAL RECURSION


A **cellular graph** of type (g, n) is the one-skeleton of a cell-decomposition of a connected closed oriented surface of genus g with n 0-cells labeled by the index set $[n] = \{1, 2, \dots, n\}$. Two cellular graphs are identified if an orientation-preserving homeomorphism of a surface into another surface maps one cellular graph to another, honoring the labeling of each vertex. Let $D_{g,n}(\mu_1, \dots, \mu_n)$ denote the number of connected cellular graphs Γ of type (g, n) with n labeled vertices of degrees (μ_1, \dots, μ_n) , counted with the weight $1/|\text{Aut}(\Gamma)|$. It is generally a rational number. The orientation of the surface induces a cyclic order of incident half-edges at each vertex of a cellular graph Γ . Since $\text{Aut}(\Gamma)$ fixes each vertex, the automorphism group is a subgroup of the Abelian group $\prod_{i=1}^n \mathbb{Z}/\mu_i \mathbb{Z}$ that rotates each vertex. Therefore,

$$(3.1) \quad C_{g,n}(\mu_1, \dots, \mu_n) = \mu_1 \cdots \mu_n D_{g,n}(\mu_1, \dots, \mu_n)$$

is always an integer. The cellular graphs counted by (3.1) are connected graphs of genus g with n vertices of degrees (μ_1, \dots, μ_n) , and at the j -th vertex for every $j = 1, \dots, n$, an outgoing arrow is placed on one of the incident μ_j half-edges (see Figure 3.1). The placement of n arrows corresponds to the factors $\mu_1 \cdots \mu_n$ on the right-hand side. We call this integer the generalized **Catalan number** of type (g, n) . The reason for this naming comes from the following theorem.

Theorem 3.1. *The generalized Catalan numbers (3.1) satisfy the following equation.*

$$(3.2) \quad C_{g,n}(\mu_1, \dots, \mu_n) = \sum_{j=2}^n \mu_j C_{g,n-1}(\mu_1 + \mu_j - 2, \mu_2, \dots, \widehat{\mu_j}, \dots, \mu_n) \\ + \sum_{\alpha + \beta = \mu_1 - 2} \left[C_{g-1, n+1}(\alpha, \beta, \mu_2, \dots, \mu_n) + \sum_{\substack{g_1 + g_2 = g \\ I \sqcup J = \{2, \dots, n\}}} C_{g_1, |I|+1}(\alpha, \mu_I) C_{g_2, |J|+1}(\beta, \mu_J) \right],$$


FIGURE 3.1. A cellular graph of type $(1, 2)$.

where $\mu_I = (\mu_i)_{i \in I}$ for an index set $I \subset [n]$, $|I|$ denotes the cardinality of I , and the third sum in the formula is for all partitions of g and set partitions of $\{2, \dots, n\}$.

Proof. Let Γ be an arrowed cellular graph counted by the left-hand side of (3.2). Since all vertices of Γ are labeled, let $\{p_1, \dots, p_n\}$ denote the vertex set. We look at the half-edge incident to p_1 that carries an arrow.

Case 1. The arrowed half-edge extends to an edge E that connects p_1 and p_j for some $j > 1$.

In this case, we shrink the edge and join the two vertices p_1 and p_j together. By this process we create a new vertex of degree $\mu_1 + \mu_j - 2$. To make the counting bijective, we need to be able to go back from the shrunk graph to the original, provided that we know μ_1 and μ_j . Thus we place an arrow to the half-edge next to E around p_1 with respect to the counter-clockwise cyclic order that comes from the orientation of the surface. In this process we have μ_j different arrowed graphs that produce the same result, because we must remove the arrow placed around the vertex p_j in the original graph. This gives the right-hand side of the first line of (3.2). See Figure 3.2.

FIGURE 3.2. The process of shrinking the arrowed edge E that connects vertices p_1 and p_j , $j > 1$.

Case 2. The arrowed half-edge at p_1 is a loop E that goes out and comes back to p_1 .

The process we apply is again shrinking the loop E . The loop E separates all other half-edges into two groups, one consisting of α of them placed on one side of the loop, and the other consisting of β half-edges placed on the other side. It can happen that $\alpha = 0$ or $\beta = 0$. Shrinking a loop on a surface causes pinching. Instead of creating a pinched (i.e., singular) surface, we separate the double point into two new vertices of degrees α and β . Here again we need to remember the place of the loop E . Thus we place an arrow to the half-edge next to the loop in each group. See Figure 3.3.

After the pinching and separating the double point, the original surface of genus g with n vertices $\{p_1, \dots, p_n\}$ may change its topology. It may have genus $g - 1$, or it splits into two pieces of genus g_1 and g_2 . The second line of (3.2) records all such cases. This completes the proof. \square




FIGURE 3.3. The process of shrinking the arrowed loop E that is attached to p_1 .

Remark 3.2. For $(g, n) = (0, 1)$, the above formula reduces to

$$(3.3) \quad C_{0,1}(\mu_1) = \sum_{\alpha+\beta=\mu_1-2} C_{0,1}(\alpha)C_{0,1}(\beta).$$

When $n = 1$, the degree of the unique vertex μ_1 is always even. By defining $C_{0,1}(0) = 1$, we find that

$$C_{0,1}(2m) = C_m = \frac{1}{m+1} \binom{2m}{m}$$

is the m -th Catalan number. Only for $(g, n) = (0, 1)$ we have this irregular case of $\mu_1 = 0$ happens, because a degree 0 single vertex is *connected*, and gives a cell-decomposition of S^2 . We can imagine that a single vertex on S^2 has an infinite cyclic group as its automorphism, so that $C_{0,1}(0) = 1$ is consistent with

$$C_{0,1}(\mu_1) = \mu_1 D_{0,1}(\mu_1).$$

All other cases, if one of the vertices has degree 0, then the Catalan number $C_{g,n}$ is simply 0 because of the definition (3.1).

Let us introduce the generating function of the Catalan numbers by

$$(3.4) \quad z = z(x) = \sum_{m=0}^{\infty} \frac{1}{x^{2m+1}}.$$

Then by the quadratic recursion (3.3) we find that the inverse function of $z(x)$ that vanishes at $x = \infty$ is given by

$$(3.5) \quad x = z + \frac{1}{z}.$$

This defines a Lagrangian immersion

$$(3.6) \quad \Sigma = \mathbb{C} \ni z \mapsto (x(z), y(z)) \in T^*\mathbb{C}, \quad \begin{cases} x(z) = z + \frac{1}{z} \\ y(z) = -z \end{cases}.$$

The Lagrangian singularities are located at the points at which $dx = 0$, i.e., $z = \pm 1$. Often it is more convenient to use the coordinate

$$(3.7) \quad z = \frac{t+1}{t-1}.$$

The following theorem is established in [20].

Theorem 3.3 ([20]). *The Laplace transform of the Catalan numbers of type (g, n) defined as symmetric differential forms by*

$$(3.8) \quad W_{g,n}^C(t_1, \dots, t_n) = (-1)^n \sum_{(\mu_1, \dots, \mu_n) \in \mathbb{Z}_+^n} C_{g,n}(\mu_1, \dots, \mu_n) e^{-\langle w, \mu \rangle} dw_1 \cdots dw_n$$

satisfies the Eynard-Orantin recursion with respect to the Lagrangian immersion (3.6) and

$$(3.9) \quad W_{0,2}^C(t_1, t_2) = \frac{dt_1 \cdot dt_2}{(t_1 - t_2)^2} - \frac{dx_1 \cdot dx_2}{(x_1 - x_2)^2}.$$

Here the Laplace transform coordinate w is related to the coordinate t of the Lagrangian by

$$e^{w_i} = x_i = z_i + \frac{1}{z_i} = \frac{t_i + 1}{t_i - 1} + \frac{t_i - 1}{t_i + 1}, \quad i = 1, 2, \dots, n,$$

and $\langle w, \mu \rangle = w_1 \mu_1 + \dots + w_n \mu_n$.

In this case the Eynard-Orantin recursion formula is given by

$$(3.10) \quad W_{g,n}^C(t_1, \dots, t_n) = -\frac{1}{64} \frac{1}{2\pi i} \int_{\gamma} \left(\frac{1}{t+t_1} + \frac{1}{t-t_1} \right) \frac{(t^2-1)^3}{t^2} \cdot \frac{1}{dt} \cdot dt_1 \\ \times \left[\sum_{j=2}^n \left(W_{0,2}^C(t, t_j) W_{g,n-1}^C(-t, t_2, \dots, \hat{t}_j, \dots, t_n) + W_{0,2}^D(-t, t_j) W_{g,n-1}^C(t, t_2, \dots, \hat{t}_j, \dots, t_n) \right) \right. \\ \left. + W_{g-1,n+1}^C(t, -t, t_2, \dots, t_n) + \sum_{\substack{\text{stable} \\ g_1+g_2=g \\ I \sqcup J = \{2,3,\dots,n\}}} W_{g_1,|I|+1}^C(t, t_I) W_{g_2,|J|+1}^C(-t, t_J) \right].$$

The last sum is restricted to the stable geometries. In other words, the partition should satisfies $2g_1 - 1 + |I| > 0$ and $2g_2 - 1 + |J|$. The contour integral is taken with respect to t on the curve defined by Figure 3.4.




FIGURE 3.4. The integration contour γ . This contour encloses an annulus bounded by two concentric circles centered at the origin. The outer one has a large radius $r > \max_{j \in N} |t_j|$ and the negative orientation, and the inner one has an infinitesimally small radius with the positive orientation.

Remark 3.4. The recursion formula (3.10) is a genuine induction formula with respect to $2g - 2 + n$. Thus from the given $W_{0,1}^C$ and $W_{0,2}^C$, we can calculate all $W_{g,n}^C$ one by one. This is a big difference between (3.10) and (3.2). The latter relation contains the terms with $C_{g,n}$ in the right-hand side as well.

4. THE PARTITION FUNCTION FOR THE GENERALIZED CATALAN NUMBERS AND THE SCHRÖDINGER EQUATION

Let us now define

$$(4.1) \quad F_{g,n}^C(t_1, \dots, t_n) = \sum_{(\mu_1, \dots, \mu_n) \in \mathbb{Z}_+^n} D_{g,n}(\mu_1, \dots, \mu_n) e^{-\langle w, \mu \rangle}$$

for $2g - 2 + n > 0$. Then from (3.1) we have

$$W_{g,n}^C(t_1, \dots, t_n) = d_1 \cdots d_n F_{g,n}^C(t_1, \dots, t_n).$$

Therefore, we have a natural primitive function of $W_{g,n}(t_1, \dots, t_n)$ for every (g, n) . We note that

$$t = -1 \implies z = 0 \implies x = \infty.$$

Therefore,

$$(4.2) \quad F_{g,n}^C(t_1, \dots, t_n) \Big|_{t_i = -1} = 0$$

for every $i = 1, 2, \dots, n$. The following recursion formula of [61] is the key for our investigation.

Theorem 4.1. *The Laplace transform $F_{g,n}^C(t_{[n]})$ satisfies the following differential recursion equation for every (g, n) subject to $2g - 2 + n > 0$.*

$$(4.3) \quad \begin{aligned} & \frac{\partial}{\partial t_1} F_{g,n}^C(t_{[n]}) \\ &= -\frac{1}{16} \sum_{j=2}^n \left[\frac{t_j}{t_1^2 - t_j^2} \left(\frac{(t_1^2 - 1)^3}{t_1^2} \frac{\partial}{\partial t_1} F_{g,n-1}^C(t_{[\hat{j}]}) - \frac{(t_j^2 - 1)^3}{t_j^2} \frac{\partial}{\partial t_j} F_{g,n-1}^C(t_{[\hat{1}]}) \right) \right. \\ & \quad \left. - \frac{1}{16} \sum_{j=2}^n \frac{(t_1^2 - 1)^2}{t_1^2} \frac{\partial}{\partial t_1} F_{g,n-1}^C(t_{[\hat{j}]}) \right. \\ & \quad \left. - \frac{1}{32} \frac{(t_1^2 - 1)^3}{t_1^2} \left[\frac{\partial^2}{\partial u_1 \partial u_2} F_{g-1,n+1}^C(u_1, u_2, t_2, t_3, \dots, t_n) \right] \Big|_{u_1 = u_2 = t_1} \right. \\ & \quad \left. - \frac{1}{32} \frac{(t_1^2 - 1)^3}{t_1^2} \sum_{\substack{\text{stable} \\ g_1 + g_2 = g \\ I \sqcup J = \{2, 3, \dots, n\}}} \frac{\partial}{\partial t_1} F_{g_1, |I|+1}^C(t_1, t_I) \frac{\partial}{\partial t_1} F_{g_2, |J|+1}^C(t_1, t_J) \right]. \end{aligned}$$

Here we use the index convention $[n] = \{1, 2, \dots, n\}$ and $[\hat{j}] = \{1, 2, \dots, \hat{j}, \dots, n\}$.

Remark 4.2. We note that the above formula is identical to [58, Theorem 5.1], even though $F_{g,n}$ is a different function. There we considered the Laplace transform of the number of lattice points in $\mathcal{M}_{g,n}$, and hence $F_{0,1} = F_{0,2} = 0$.

Remark 4.3. Because of (4.2), the recursion (4.3) uniquely determines each $F_{g,n}^C$ by integrating from -1 to t_1 . With the same reason, $F_{g,n}^C$ is uniquely determined by $W_{g,n}^C$. Since we know exactly where $F_{g,n}^C$ vanishes, there is no discrepancy of the constants of integration in (2.12).

Let us now consider the principal specialization of the partition function for the Catalan numbers

$$(4.4) \quad Z^C(t, \hbar) = \exp \left(\sum_{g=0}^{\infty} \sum_{n=1}^{\infty} \frac{1}{n!} \hbar^{2g-2+n} F_{g,n}^C(t, t, \dots, t) \right).$$

Since unstable terms $F_{0,1}^C(t)$ and $F_{0,2}^C(t, t)$ are included in the above formula, we need to calculate them first.

Proposition 4.4. *In terms of the z -variable, we have*

$$(4.5) \quad F_{0,1}^C(t) = -\frac{1}{2}z^2 + \log z,$$

$$(4.6) \quad F_{0,2}^C(t_1, t_2) = -\log(1 - z_1 z_2).$$

Proof. Due to the irregularity of $\mu = 0$ for $D_{0,1}(\mu)$, we need to modify the definitions (3.8) and (4.1) for $(g, n) = (0, 1)$. It is natural to define

$$(4.7) \quad W_{0,1}(t) = -\sum_{m=0}^{\infty} C_{0,1}(2m) \frac{dx}{x^{2m+1}} = -z dx = \left(-z + \frac{1}{z}\right) dz$$

because of the consistency with (2.7). Since $W_{0,1}^C = dF_{0,1}^C$, we have

$$\begin{aligned} F_{0,1}(t) &= -\frac{1}{2}z^2 + \log z + \text{const} \\ &= \sum_{m=0}^{\infty} D_{0,1}(2m) \left(\frac{1}{x^{2m}} - \delta_{m,0}\right), \end{aligned}$$

where the $m = 0$ term is adjusted so that we do not have the infinity term $D_{0,1}(0)$ in $F_{0,1}^C$. Using the expression

$$D_{0,1}(2m) = \frac{1}{2m(m+1)} \binom{2m}{m},$$

we have

$$\lim_{m \rightarrow 0} D_{0,1}(2m) \left(\frac{1}{x^{2m}} - \delta_{m,0}\right) = -\log x.$$

Since $x = z + \frac{1}{z}$, by taking the limit $z \rightarrow 0$ we conclude that the constant term in $F_{0,1}^C(t)$ is 0, which establishes (4.5).

The computation of $F_{0,1}^C(t_1, t_2)$ is performed in [20, Proposition 4.1], where the idea is to use the Euler differential operator $x_1 \frac{d}{dx_1} + x_2 \frac{d}{dx_2}$. By definition $F_{0,1}^C(t_1, t_2)$ does not have any constant term as $x \rightarrow \infty$, therefore there is no constant correction in (4.6), either. \square

Theorem 4.5. *The principal specialization of the partition function satisfies the following Schrödinger equation.*

$$(4.8) \quad \left(\hbar^2 \frac{d^2}{dx^2} + \hbar x \frac{d}{dx} + 1\right) Z^C(t, \hbar) = 0,$$

where t is considered as a function in x by

$$t = t(x) = \frac{z(x) + 1}{z(x) - 1}$$

and (3.4).

The rest of this section is devoted to proving the above theorem. Let us define for $m \geq 0$

$$(4.9) \quad S_m = \sum_{2g-2+n=m-1} \frac{1}{n!} F_{g,n}^C(t, \dots, t),$$

and put

$$F = \sum_{m=0}^{\infty} \hbar^{m-1} S_m.$$

Then the Schrödinger equation (4.8) becomes

$$\hbar^2 \left(\frac{d^2 F}{dx^2} + \left(\frac{dF}{dx} \right)^2 \right) + \hbar x \frac{dF}{dx} + 1 = 0,$$

which is equivalent to

$$(4.10) \quad \sum_{m=0}^{\infty} S_m'' \hbar^{m+1} + \left(\sum_{m=0}^{\infty} S_m' \hbar^m \right)^2 + x \sum_{m=0}^{\infty} S_m' \hbar^m + 1 = 0,$$

where $' = \frac{d}{dx}$. Since $S_0 = F_{0,1}^C$ and

$$W_{0,1}^C = dF_{0,1}^C = ydx = -zdx,$$

we obtain

$$(4.11) \quad S_0' = \frac{d}{dx} F_{0,1}^C = -z = -\frac{t+1}{t-1}.$$

Using the Lagrangian immersion (3.5), we see that the constant terms of (4.10) then become

$$(4.12) \quad (-z)^2 + x(-z) + 1 = z^2 - xz + 1 = 0.$$

Collecting the \hbar^{m+1} -contributions in (4.10) for $m \geq 0$, we obtain

$$\begin{aligned} -x \frac{d}{dx} S_{m+1} &= \frac{d^2}{dx^2} S_m + \sum_{a+b=m+1} \frac{dS_a}{dx} \frac{dS_b}{dx} \\ &= \frac{d^2}{dx^2} S_m + 2S_0' S_{m+1}' + 2S_1' S_m' + \sum_{\substack{a+b=m+1 \\ a,b \geq 2}} \frac{dS_a}{dx} \frac{dS_b}{dx}. \end{aligned}$$

Therefore,

$$(4.13) \quad -(2S_0' + x) \frac{d}{dx} S_{m+1} = \frac{d^2}{dx^2} S_m + 2S_1' \frac{d}{dx} S_m + \sum_{\substack{a+b=m+1 \\ a,b \geq 2}} \frac{dS_a}{dx} \frac{dS_b}{dx}.$$

To use the closed formulas (4.5) and (4.6) for S_0 and S_1 here, we need to switch from the x -coordinate to the t -coordinate. Using the change of variable formulas (3.5) and (3.7), we have

$$dx = dz \left(1 - \frac{1}{z^2} \right) = -\frac{8t}{(t^2 - 1)^2} dt,$$

hence

$$\begin{aligned} \frac{d}{dx} &= \frac{z^2}{z^2 - 1} \frac{d}{dz} = -\frac{(t^2 - 1)^2}{8t} \frac{d}{dt}, \\ \frac{d^2}{dx^2} &= \frac{(t^2 - 1)^4}{64t^2} \frac{d^2}{dt^2} + \frac{(t^2 - 1)^2}{8t} \left(\frac{d}{dt} \frac{(t^2 - 1)^2}{8t} \right) \frac{d}{dt} \\ &= \frac{(t^2 - 1)^4}{64t^2} \frac{d^2}{dt^2} + \frac{(t^2 - 1)^3}{64t^3} (3t^2 + 1) \frac{d}{dt}. \end{aligned}$$

Therefore,

$$(4.14) \quad -(2S_0' + x) \frac{d}{dx} = -\frac{(t^2 - 1)^2}{8t} \left(z - \frac{1}{z} \right) \frac{d}{dt} = -\frac{(t^2 - 1)^2}{8t} \frac{4t}{t^2 - 1} \frac{d}{dt} = \frac{1}{2} (t^2 - 1) \frac{d}{dt},$$

$$(4.15) \quad S'_1 = \frac{1}{2} \frac{d}{dx} F_{0,2}^C(z, z) = \frac{1}{2} \frac{z^2}{z^2 - 1} \frac{d}{dz} (-\log(1 - z^2)) = -\frac{(t^2 - 1)(t + 1)^2}{16t^2}.$$

Substituting (4.11), (4.14) and (4.15) in (4.13), we obtain

Proposition 4.6. *For $m \geq 0$, the Schrödinger equation (4.8) is equivalent to a recursion formula*

$$(4.16) \quad \frac{d}{dt} S_{m+1} = -\frac{(t^2 - 1)^3}{32t^2} \left[\frac{d^2}{dt^2} S_m + \sum_{\substack{a+b=m+1 \\ a, b \geq 2}} \frac{dS_a}{dt} \frac{dS_b}{dt} \right] - \frac{(t^2 - 1)^2}{16t^3} (2t^2 + t + 1) \frac{d}{dt} S_m.$$

Remark 4.7. The recursion equation (4.16) is a direct consequence of the principal specialization applied to (4.3).

The proof of (4.16) is given in Appendix A.

5. SINGLE HURWITZ NUMBERS

The A-model problem that we are interested in this section is the automorphism-weighted count $H_{g,n}(\mu_1, \dots, \mu_n)$ of the number of the topological types of meromorphic functions $f : C \rightarrow \mathbb{P}^1$ of a nonsingular complete irreducible algebraic curve C of genus g that has n labeled poles of orders (μ_1, \dots, μ_n) such that all other critical points of f than these poles are unlabeled simple ramification points. Let r denote the number of such simple ramification points. Then the Riemann-Hurwitz formula gives

$$(5.1) \quad r = 2g - 2 + n + \sum_{i=1}^n \mu_i.$$

A remarkable formula due to Ekedahl, Lando, Shapiro and Vainshtein [22, 37, 68] relates Hurwitz numbers and Gromov-Witten invariants. For genus $g \geq 0$ and the number $n \geq 1$ of the poles subject to the stability condition $2g - 2 + n > 0$, the ELSV formula states that

$$(5.2) \quad H_{g,n}(\mu_1, \dots, \mu_n) = \prod_{i=1}^n \frac{\mu_i^{\mu_i}}{\mu_i!} \int_{\overline{\mathcal{M}}_{g,n}} \frac{\Lambda_g^\vee(1)}{\prod_{i=1}^n (1 - \mu_i \psi_i)} \\ = \sum_{j=0}^g (-1)^j \sum_{k_1, \dots, k_n \geq 0} \langle \tau_{k_1} \cdots \tau_{k_n} c_j(\mathbb{E}) \rangle_{g,n} \prod_{i=1}^n \frac{\mu_i^{\mu_i + k_i}}{\mu_i!},$$

where $\overline{\mathcal{M}}_{g,n}$ is the Deligne-Mumford moduli stack of stable algebraic curves of genus g with n distinct smooth marked points, $\Lambda_g^\vee(1) = 1 - c_1(\mathbb{E}) + \cdots + (-1)^g c_g(\mathbb{E})$ is the alternating sum of the Chern classes of the Hodge bundle \mathbb{E} on $\overline{\mathcal{M}}_{g,n}$, ψ_i is the i -th tautological cotangent class, and

$$(5.3) \quad \langle \tau_{k_1} \cdots \tau_{k_n} c_j(\mathbb{E}) \rangle_{g,n} = \int_{\overline{\mathcal{M}}_{g,n}} \psi_1^{k_1} \cdots \psi_n^{k_n} c_j(\mathbb{E})$$

is the linear Hodge integral, which is 0 unless $k_1 + \cdots + k_n + j = 3g - 3 + n$. Although the Deligne-Mumford stack $\overline{\mathcal{M}}_{g,n}$ is not defined for $2 - 2g - n < 0$, single Hurwitz numbers are well defined for *unstable* geometries $(g, n) = (0, 1)$ and $(0, 2)$, and their values are

$$(5.4) \quad H_0((d)) = \frac{d^{d-3}}{(d-1)!} = \frac{d^{d-2}}{d!} \quad \text{and} \quad H_0((\mu_1, \mu_2)) = \frac{1}{\mu_1 + \mu_2} \cdot \frac{\mu_1^{\mu_1}}{\mu_1!} \cdot \frac{\mu_2^{\mu_2}}{\mu_2!}.$$

The ELSV formula remains valid for unstable cases if we define

$$(5.5) \quad \int_{\overline{\mathcal{M}}_{0,1}} \frac{\Lambda_0^\vee(1)}{1-d\psi} = \frac{1}{d^2},$$

$$(5.6) \quad \int_{\overline{\mathcal{M}}_{0,2}} \frac{\Lambda_0^\vee(1)}{(1-\mu_1\psi_1)(1-\mu_2\psi_2)} = \frac{1}{\mu_1 + \mu_2}.$$

The ELSV formula predicts that the single Hurwitz numbers exhibit the polynomial behavior in terms of their Laplace transform. Following [26], and modifying our choice of the coordinates slightly, we define

$$(5.7) \quad \begin{aligned} F_{g,n}^H(t_1, \dots, t_n) &= \sum_{\mu \in \mathbb{Z}_+^n} H_{g,n}(\mu_1, \dots, \mu_n) e^{-(\mu_1 w_1 + \dots + \mu_n w_n)} \\ &= \sum_{\mu \in \mathbb{Z}_+^n} \sum_{k_1 + \dots + k_n \leq 3g-3+n} \langle \tau_{k_1} \cdots \tau_{k_n} \Lambda_g^\vee(1) \rangle_{g,n} \left(\prod_{i=1}^n \frac{\mu_i^{\mu_i + k_i}}{\mu_i!} \right) e^{-(\mu_1 w_1 + \dots + \mu_n w_n)} \\ &= \sum_{k_1 + \dots + k_n \leq 3g-3+n} \langle \tau_{k_1} \cdots \tau_{k_n} \Lambda_g^\vee(1) \rangle_{g,n} \prod_{i=1}^n \hat{\xi}_{k_i}(t_i), \end{aligned}$$

where the $\hat{\xi}$ -functions are given by

$$(5.8) \quad \hat{\xi}_k(t) = \sum_{\mu=1}^{\infty} \frac{\mu^{\mu+k}}{\mu!} e^{-\mu w} = \sum_{\mu=1}^{\infty} \frac{\mu^{\mu+k}}{\mu!} x^\mu$$

for $k \geq 0$, and $x = e^{-w}$. Although $\hat{\xi}_k$ are complicated functions in x , their behavior is simple in terms of $\hat{\xi}_0$. So we introduce

$$(5.9) \quad t = 1 + \sum_{\mu=1}^{\infty} \frac{\mu^\mu}{\mu!} x^\mu \quad \text{and} \quad z = \sum_{\mu=1}^{\infty} \frac{\mu^{\mu-1}}{\mu!} x^\mu.$$

Then by the Lagrange inversion formula, we have

$$(5.10) \quad x = z e^{-z} \quad \text{and} \quad z = \frac{t-1}{t},$$

and moreover, each $\hat{\xi}_k(t)$ is a polynomial of degree $2k+1$ in t for every $k \geq 0$, recursively defined by

$$(5.11) \quad \hat{\xi}_{k+1}(t) = t^2(t-1) \frac{d}{dt} \hat{\xi}_k(t).$$

This is because

$$(5.12) \quad -\frac{d}{dw} = x \frac{d}{dx} = t^2(t-1) \frac{d}{dt}.$$

Therefore, if $2g-2+n > 0$, then $F_{g,n}^H(t_1, \dots, t_n)$ is a symmetric polynomial of degree $6g-6+3n$, and satisfies

$$(5.13) \quad F_{g,n}^H(t_1, \dots, t_n) \Big|_{t_j=1} = 0$$

for every $j = 1, 2, \dots, n$.

The computation of [20] adjusted to our current convention of this paper,

$$(5.14) \quad F_{0,1}^H(t) = \frac{1}{2} \left(1 - \frac{1}{t^2} \right) = z - \frac{1}{2} z^2$$

and

$$(5.15) \quad F_{0,2}^H(t_1, t_2) = \log \left(\frac{z_1 - z_2}{x_1 - x_2} \right) - (z_1 + z_2),$$

determines the Lagrangian immersion

$$\iota : \Sigma = \mathbb{C}^* \longrightarrow T^*\mathbb{C}^*$$

by

$$(5.16) \quad \begin{cases} x = ze^{-z} \\ y = z, \end{cases}$$

and

$$W_{0,2}^H(t_1, t_2) = d_1 \otimes d_2 F_{0,2}^H(t_1, t_2) = \frac{dt_1 \cdot dt_2}{(t_1 - t_2)^2} - \frac{dx_1 \cdot dx_2}{(x_1 - x_2)^2}.$$

Here the tautological 1-form on $T^*\mathbb{C}^*$ is chosen to be

$$(5.17) \quad \eta = y \frac{dx}{x}.$$

It is consistent with

$$W_{0,1}^H(t) = dF_{0,1}^H(t) = z \frac{dx}{x} = \iota^* \eta.$$

We also note that

$$(5.18) \quad F_{0,1}^H(t)|_{t=1} = 0,$$

and

$$(5.19) \quad F_{0,2}^H(t_1, t_2)|_{t_j=1} = 0, \quad j = 1 \text{ or } 2.$$

The latter equality holds because $t_2 = 1 \implies z_2 = 0 \implies x_2 = 0$, and hence from (5.15) we have

$$F_{0,2}^H(t_1, 1) = \log \frac{z_1}{x_1} - z_1 = 0.$$

Since

$$\left. \frac{x_1 - x_2}{z_1 - z_2} \right|_{z_1=z_2=z} = (1 - z)e^{-z},$$

the diagonal value of $F_{0,2}^H$ is calculated as

$$(5.20) \quad F_{0,2}^H(t, t) = \log \left(\frac{e^z}{-z} \right) - 2z = -z - \log(1 - z) = \frac{1-t}{t} + \log t.$$

The single Hurwitz numbers $H_{g,n}(\vec{\mu})$ satisfy the *cut-and-join* equation [36, 77]

$$(5.21) \quad \left(2g - 2 + n + \sum_{i=1}^n \mu_i \right) H_{g,n}(\mu_1, \dots, \mu_n) = \frac{1}{2} \sum_{i \neq j} (\mu_i + \mu_j) H_{g,n-1}(\mu_i + \mu_j, \mu_{[\hat{i}, \hat{j}]}) \\ + \frac{1}{2} \sum_{i=1}^n \sum_{\alpha+\beta=\mu_i} \alpha\beta \left[H_{g-1,n+1}(\alpha, \beta, \mu_{[\hat{i}]}) + \sum_{\substack{g_1+g_2=g \\ I \sqcup J = [\hat{i}]}} H_{g_1,|I|+1}(\alpha, \mu_I) H_{g_2,|J|+1}(\beta, \mu_J) \right],$$

where we use the convention for indices as in Section 3. The Laplace transform of (5.21) is the polynomial recursion of [60] and takes the form

$$\begin{aligned}
(5.22) \quad & \left(2g - 2 + n + \sum_{i=1}^n t_i(t_i - 1) \frac{\partial}{\partial t_i} \right) F_{g,n}^H(t_1, \dots, t_n) \\
&= \frac{1}{2} \sum_{i \neq j} \frac{t_i t_j}{t_i - t_j} \left(t_i^2 (t_i - 1)^2 \frac{\partial}{\partial t_i} F_{g,n-1}^H(t_{[\hat{j}]}), - t_j^2 (t_j - 1)^2 \frac{\partial}{\partial t_j} F_{g,n-1}^H(t_{[\hat{i}]}) \right) \\
&\quad - \sum_{i \neq j} t_i^3 (t_i - 1) \frac{\partial}{\partial t_i} F_{g,n-1}^H(t_{[\hat{j}]}) \\
&\quad + \frac{1}{2} \sum_{i=1}^n (t_i^2 (t_i - 1))^2 \frac{\partial^2}{\partial u_1 \partial u_2} F_{g-1,n+1}^H(u_1, u_2, t_{[\hat{i}]}) \Big|_{u_1=u_2=t_i} \\
&\quad + \frac{1}{2} \sum_{i=1}^n (t_i^2 (t_i - 1))^2 \sum_{\substack{\text{stable} \\ g_1+g_2=g \\ I \sqcup J = [\hat{i}]}} \frac{\partial}{\partial t_i} F_{g_1,|I|+1}^H(t_i, t_I) \cdot \frac{\partial}{\partial t_i} F_{g_2,|J|+1}^H(t_i, t_J).
\end{aligned}$$

It is proved in [26] that the n -fold exterior differentiation of the above formula is exactly the Eynard-Orantin recursion, as predicted by Bouchard and Mariño [9]. Thus we obtain a natural integration of the Eynard-Orantin recursion by taking the Laplace transform of the A-model quantity again, which is the single Hurwitz umber $H_{g,n}(\vec{\mu})$.

Theorem 5.1. *Let us define*

$$(5.23) \quad S_m^H(t) = \sum_{2g-2+n=m-1} \frac{1}{n!} F_{g,n}^H(t, \dots, t).$$

Then S_m^H 's satisfy the following second order differential equations:

$$(5.24) \quad \left(e^{mw} \frac{d}{dw} e^{-mw} \right) S_{m+1}^H = -\frac{1}{2} \left[\frac{d^2}{dw^2} S_m^H + \sum_{a+b=m+1} \frac{dS_a^H}{dw} \cdot \frac{dS_b^H}{dw} + \frac{d}{dw} S_m^H \right].$$

Here the w -dependence of t is given by $x = e^{-w}$ and (5.9). We also note that $S_m^H(t)$ is a polynomial of degree $3m-3$ for every $m \geq 2$, and for all values of m we have

$$(5.25) \quad S_m^H(t) \Big|_{t=1} = 0.$$

The proof is similar to the case of the Catalan numbers (Section 4 and Appendix A). First we compute the principal specialization of the differential equation (5.22). We then assemble them according to (5.23). By adjusting the unstable geometry terms $(g, n) = (0, 1)$ and $(0, 2)$, we obtain (5.24). However, due to the difference between the cut-and-join equation and the edge-shrinking operation of Section 3, the resulting equation becomes quite different.

Choose $m \geq 2$ and (g, n) so that $2g - 2 + n = m$. Then the principal specialization of the left-hand side of (5.22) is

$$\left(m + t(t-1) \frac{d}{dt} \right) F_{g,n}^H(t, \dots, t).$$

The first line of the right-hand side of (5.22) gives

$$\begin{aligned}
& \frac{1}{2} t^2 \sum_{i \neq j} \frac{\partial}{\partial t_i} \left(t_i^2 (t_i - 1)^2 \frac{\partial}{\partial t_i} F_{g,n-1}^H(t_i, t, \dots, t) \right) \Big|_{t_i=t} \\
&= \frac{n(n-1)}{2} t^2 \frac{d}{dt} (t^2 (t-1)^2) \cdot \frac{1}{n-1} \frac{d}{dt} F_{g,n-1}^H(t, \dots, t)
\end{aligned}$$

$$\begin{aligned}
& + \frac{n(n-1)}{2} t^4 (t-1)^2 \frac{\partial^2}{\partial u^2} F_{g,n-1}^H(u, t, \dots, t) \Big|_{u=t} \\
& = \frac{1}{2} n! t^2 \frac{d}{dt} (t^2 (t-1)^2) \cdot \frac{1}{(n-1)!} \frac{d}{dt} F_{g,n-1}^H(t, \dots, t) \\
& \quad + \frac{1}{2} n! (n-1) t^4 (t-1)^2 \frac{1}{(n-1)!} \frac{\partial^2}{\partial u^2} F_{g,n-1}^H(u, t, \dots, t) \Big|_{u=t}.
\end{aligned}$$

The second line of the right-hand side of (5.22) becomes

$$-n! t^3 (t-1) \frac{1}{(n-1)!} \frac{d}{dt} F_{g,n-1}^H(t, \dots, t).$$

The third line simply produces

$$\frac{n!}{2} (t^2 (t-1))^2 \frac{1}{(n+1)!} (n+1)n \frac{\partial^2}{\partial u_1 \partial u_2} F_{g-1, n+1}^H(u_1, u_2, t, \dots, t) \Big|_{u_1=u_2=t}.$$

Finally, since the set partition becomes the partition of numbers because all variables are set to be equal, the fourth line of the right-hand side of (5.22) gives

$$\frac{n!}{2} (t^2 (t-1))^2 \sum_{\substack{\text{stable} \\ g_1+g_2=g \\ n_1+n_2=n-1}} \frac{1}{(n_1+1)!} \frac{d}{dt} F_{g_1, n_1+1}^H(t, \dots, t) \cdot \frac{1}{(n_2+1)!} \frac{d}{dt} F_{g_2, n_2+1}^H(t, \dots, t).$$

We now apply the operation

$$\sum_{2g-2+n=m} \frac{1}{n!}$$

to the above terms. The left-hand side becomes

$$\left(m + t(t-1) \frac{d}{dt} \right) S_{m+1}^H.$$

The right-hand side terms are re-assembled into the sum of (g', n') subject to $2g' - 2 + n' = m - 1$, following the topological structure of the recursion (5.22). Noticing that unstable geometers are contained only in S_0^H and S_1^H , we obtain

$$\begin{aligned}
(5.26) \quad & \left(m + t(t-1) \frac{d}{dt} \right) S_{m+1}^H(t) \\
& = \frac{1}{2} \left(t^2 \frac{d}{dt} (t^2 (t-1)^2) \cdot \frac{d}{dt} S_m^H(t) + (t^2 (t-1))^2 \frac{d^2}{dt^2} S_m^H(t) - 2t^3 (t-1) \frac{d}{dt} S_m^H(t) \right) \\
& \quad + \frac{1}{2} (t^2 (t-1)^2)^2 \sum_{\substack{a+b=m+1 \\ a, b \geq 2}} \frac{d}{dt} S_a^H(t) \cdot \frac{d}{dt} S_b^H(t) \\
& = \frac{1}{2} (t^2 (t-1))^2 \left(\frac{d^2}{dt^2} S_m^H(t) + \sum_{\substack{a+b=m+1 \\ a, b \geq 2}} \frac{d}{dt} S_a^H(t) \cdot \frac{d}{dt} S_b^H(t) \right) + 2t^3 (t-1)^2 \frac{d}{dt} S_m^H(t).
\end{aligned}$$

Proposition 5.2. *The functions $S_m(t)$ are recursively determined by*

$$(5.27) \quad S_0^H(t) = \frac{1}{2} \left(1 - \frac{1}{t^2} \right),$$

$$(5.28) \quad S_1^H(t) = \frac{1}{2} \left(\frac{1-t}{t} + \log t \right),$$

and

$$(5.29) \quad S_{m+1}^H(t) = \left(\frac{t-1}{t} \right)^{-m} \int_1^t \left[\frac{1}{2} t^{3-m} (t-1)^{m+1} \left(\frac{d^2}{dt^2} S_m^H(t) + \sum_{\substack{a+b=m+1 \\ a,b \geq 2}} \frac{d}{dt} S_a^H(t) \cdot \frac{d}{dt} S_b^H(t) \right) + 2t^{2-m} (t-1)^{m+1} \frac{d}{dt} S_m^H(t) \right] dt.$$

Proof. As a differential operator,

$$m + t(t-1) \frac{d}{dt} = t(t-1) \left(\frac{t-1}{t} \right)^{-m} \frac{d}{dt} \left(\frac{t-1}{t} \right)^m.$$

Therefore, (5.26) is equivalent to

$$\begin{aligned} & \left[\left(\frac{t-1}{t} \right)^{-m} \frac{d}{dt} \left(\frac{t-1}{t} \right)^m \right] S_{m+1}^H(t) \\ &= \frac{1}{2} t^3 (t-1) \left(\frac{d^2}{dt^2} S_m^H(t) + \sum_{\substack{a+b=m+1 \\ a,b \geq 2}} \frac{d}{dt} S_a^H(t) \cdot \frac{d}{dt} S_b^H(t) \right) + 2t^2 (t-1) \frac{d}{dt} S_m^H(t). \end{aligned}$$

On the right-hand side only $S_k^H(t)$ with $k \leq m$ appear. Using the fact of the zero (5.25), we obtain (5.29). \square

On the third line of (5.26) the terms with S_0^H and S_1^H are not included. More precisely, these omitted terms are

$$t(t-1)^2 \frac{d}{dt} S_{m+1}^H(t) + \frac{1}{2} t^2 (t-1)^3 \frac{d}{dt} S_m^H(t).$$

When we add these terms to (5.26), and adjust the second order differentiation as

$$\left(t^2 (t-1) \frac{d}{dt} \right)^2 = (t^2 (t-1))^2 \frac{d^2}{dt^2} + t^3 (t-1) (3t-2) \frac{d}{dt},$$

we finally obtain

$$(5.30) \quad \begin{aligned} & \left(m + t^2 (t-1) \frac{d}{dt} \right) S_{m+1}^H(t) \\ &= \frac{1}{2} \left(\left(t^2 (t-1) \frac{d}{dt} \right)^2 S_m^H(t) + (t^2 (t-1))^2 \sum_{a+b=m+1} \frac{dS_a^H(t)}{dt} \cdot \frac{dS_b^H(t)}{dt} - t^2 (t-1) \frac{d}{dt} S_m^H(t) \right). \end{aligned}$$

Then (5.24) follows from (5.30) and (5.12). This completes the proof Theorem 5.1.

Theorem 5.3. *Let us define the partition function for the single Hurwitz numbers in a similar way:*

$$(5.31) \quad Z^H(t, \hbar) = \exp \left(\sum_{m=0}^{\infty} S_m^H \hbar^{m-1} \right) = \exp \left(\sum_{g=0}^{\infty} \sum_{n=1}^{\infty} \frac{1}{n!} \hbar^{2g-2+n} F_{g,n}^H(t, \dots, t) \right).$$

Then we have

$$(5.32) \quad \left[\frac{1}{2} \frac{\partial^2}{\partial w^2} + \left(\frac{1}{2} + \frac{1}{\hbar} \right) \frac{\partial}{\partial w} - \frac{\partial}{\partial \hbar} \right] Z^H(t, \hbar) = 0.$$

Remark 5.4. Eq.(5.32) is a heat equation, where \hbar is considered as the *time* variable of the heat conduction. It determines the solution uniquely with the “initial condition”

$$Z^H(t(w), \hbar)|_{\hbar \sim 0} = \exp \left(\frac{1}{\hbar} S_0^H + S_1^H \right)$$

given by (5.27) and (5.28).

Proof. Let

$$F^H(t, \hbar) = \sum_{m=0}^{\infty} S_m^H \hbar^{m-1} = \sum_{g=0}^{\infty} \sum_{n=1}^{\infty} \frac{1}{n!} \hbar^{2g-2+n} F_{g,n}^H(t, \dots, t).$$

In terms of F^H , (5.32) is equivalent to

$$(5.33) \quad \frac{\hbar}{2} \frac{\partial^2 F^H}{\partial w^2} + \frac{\hbar}{2} \left(\frac{\partial F^H}{\partial w} \right)^2 + \left[\left(1 + \frac{\hbar}{2} \right) \frac{\partial}{\partial w} - \hbar \frac{\partial}{\partial \hbar} \right] F^H = 0.$$

We apply the operation

$$\sum_{m=0}^{\infty} \hbar^m$$

to (5.24). The left-hand side is

$$(5.34) \quad \sum_{m=0}^{\infty} \hbar^m \left(-m + \frac{d}{dw} \right) S_{m+1}^H = \left(\frac{\partial}{\partial w} - \hbar \frac{\partial}{\partial \hbar} \right) \left(F^H - \frac{1}{\hbar} S_0^H \right).$$

The right-hand side gives

$$(5.35) \quad -\frac{\hbar}{2} \left(\frac{\partial^2}{\partial w^2} \sum_{m=0}^{\infty} S_m^H \hbar^{m-1} + \sum_{m=0}^{\infty} \sum_{a+b=m+1} \frac{\partial S_a^H}{\partial w} \hbar^{a-1} \cdot \frac{\partial S_b^H}{\partial w} \hbar^{b-1} + \frac{\partial}{\partial w} \sum_{m=0}^{\infty} S_m^H \hbar^{m-1} \right) \\ = -\frac{\hbar}{2} \left(\frac{\partial^2}{\partial w^2} F^H + \left(\frac{\partial F^H}{\partial w} \right)^2 - \frac{1}{\hbar^2} \left(\frac{\partial S_0^H}{\partial w} \right)^2 + \frac{\partial}{\partial w} F^H \right).$$

If we collect all terms that contain $S_0^H(t)$ in (5.34) and (5.35), we have an equation

$$\left(t^2(t-1) \frac{\partial}{\partial t} + \hbar \frac{\partial}{\partial \hbar} \right) \frac{1}{\hbar} S_0^H = \frac{1}{2\hbar} \left(t^2(t-1) \frac{\partial}{\partial t} S_0^H \right)^2,$$

or equivalently,

$$(5.36) \quad S_0^H(t) = t^2(t-1) \frac{d}{dt} S_0^H(t) - \frac{1}{2} (t^2(t-1))^2 \left(\frac{dS_0^H(t)}{dt} \right)^2.$$

We see that (5.27) is a solution to (5.36). After eliminating (5.36) from (5.34) = (5.35), we obtain (5.33). This completes the proof. \square

6. THE SCHUR FUNCTION EXPANSION OF THE HURWITZ PARTITION FUNCTION

Let us introduce the *free energy* of single Hurwitz numbers by a formal sum as

$$(6.1) \quad \begin{aligned} F^H(t_1, t_2, t_3, \dots; \hbar) &= \sum_{g \geq 0, n \geq 1} \frac{1}{n!} \hbar^{2g-2+n} F_{g,n}^H(t_1, \dots, t_n) \\ &= \sum_{g \geq 0, n \geq 1} \frac{1}{n!} \hbar^{2g-2+n} \sum_{(\mu_1, \dots, \mu_n) \in \mathbb{Z}_+^n} H_{g,n}(\mu_1, \dots, \mu_n) e^{-(\mu_1 + \dots + \mu_n)} \prod_{i=1}^n x_i^{\mu_i}. \end{aligned}$$

The partition function we considered in Section 5 is the principal specialization

$$Z^H(t, \hbar) = \exp(F^H(t, t, \dots; \hbar)).$$

Recall another generating function of the Hurwitz numbers [41, 42, 67] defined by

$$(6.2) \quad \mathbf{H}(s, \mathbf{p}) = \sum_{g \geq 0, n \geq 1} \mathbf{H}_{g,n}(s, \mathbf{p}),$$

$$(6.3) \quad \mathbf{H}_{g,n}(s, \mathbf{p}) = \frac{1}{n!} \sum_{\vec{\mu} \in \mathbb{Z}_+^n} H_{g,n}(\vec{\mu}) \mathbf{p}_\mu s^{r(g, \mu)},$$

where $\mathbf{p}_\mu = p_{\mu_1} \cdots p_{\mu_n}$, and

$$r = r(g, \mu) = 2g - 2 + n + \sum_{i=1}^n \mu_i$$

is again the number of simple ramification point of a Hurwitz cover of genus g and profile μ .

At this point we wish to go back and forth between the following two distinct points of view: One is to regard $\vec{\mu} = (\mu_1, \dots, \mu_n)$ as a vector consisting of positive integers, and the other is to view μ as a *partition* of length n . For any function $f(\vec{\mu})$ in $\vec{\mu}$ as a vector, we have a change of summation formula

$$(6.4) \quad \sum_{\vec{\mu} \in \mathbb{Z}_+^n} f(\vec{\mu}) = \sum_{\mu: \ell(\mu) = n} \frac{1}{|\text{Aut}(\mu)|} \sum_{\sigma \in S_n} f(\vec{\mu}_\sigma).$$

Here the first sum in the right-hand side runs over partitions μ of a fixed length n , the second sum is over the symmetric group S_n of n letters,

$$\vec{\mu}_\sigma = (\mu_{\sigma(1)}, \dots, \mu_{\sigma(n)}) \in \mathbb{Z}_+^n$$

is the integer vector obtained by permuting the parts of μ by $\sigma \in S_n$, and $\text{Aut}(\mu)$ is the permutation group interchanging the equal parts of μ . As a partition, the length of μ is denoted by $\ell(\mu)$, and its *size* by

$$|\mu| = \sum_{i=1}^{\ell(\mu)} \mu_i.$$

Often single Hurwitz numbers are labeled by the genus g and a *partition* μ . In this case the expression

$$h_{g, \mu} = \frac{r(g, \mu)!}{|\text{Aut}(\mu)|} H_{g, \ell(\mu)}(\mu_1, \dots, \mu_{\ell(\mu)})$$

is used in the literature, when we do not label the poles, but label the simple ramification points. The generating function then has an expression in terms of sumes over partitions:

$$\mathbf{H}(s, \mathbf{p}) = \sum_{g=0}^{\infty} \sum_{\mu} h_{g,\mu} \mathbf{p}_{\mu} \frac{s^{r(g,\mu)}}{r(g,\mu)!}.$$

In terms of the ELSV formula (5.2) we have

$$(6.5) \quad \mathbf{H}(s, \mathbf{p}) = \sum_{g \geq 0, n \geq 1} \frac{1}{n!} s^{2g-2+n} \sum_{k_1 + \dots + k_n \leq 3g-3+n} \langle \tau_{k_1} \cdots \tau_{k_n} \Lambda_g^{\vee}(1) \rangle_{g,n} \prod_{i=1}^n \sum_{\mu_i=1}^{\infty} \frac{\mu_i^{\mu_i+k_i}}{\mu_i!} s^{\mu_i} p_{\mu_i}$$

with an appropriate incorporation of (5.5) and (5.6). Recall the Laplace transform (5.7) here for comparison that is assembled into the *free energy*

$$(6.6) \quad F^H(t_1, t_2, \dots; \hbar) := \sum_{g \geq 0, n \geq 1} \frac{1}{n!} \hbar^{2g-2+n} F_{g,n}^H(t_1, \dots, t_n) = \sum_{g \geq 0, n \geq 1} \frac{1}{n!} \hbar^{2g-2+n} \sum_{k_1 + \dots + k_n \leq 3g-3+n} \langle \tau_{k_1} \cdots \tau_{k_n} \Lambda_g^{\vee}(1) \rangle_{g,n} \prod_{i=1}^n \sum_{\mu_i=1}^{\infty} \frac{\mu_i^{\mu_i+k_i}}{\mu_i!} x_i^{\mu_i}.$$

It is easy to see that the relation between the two sets of variables is exactly the power-sum symmetric functions. Let us re-scale the usual power-sum symmetric function p_j of degree j in x_i 's with a scale parameter s as follows:

$$(6.7) \quad p_j(s) := s^{-j} \left(x_1^j + x_2^j + x_3^j + \dots \right).$$

Here we consider $p_j(s)$ as a degree j polynomial defined on \mathbb{C}^n , but the dimension n is unspecified. Then for every $\mu \in \mathbb{Z}_+^n$, we have

$$(6.8) \quad d_1 \cdots d_n \mathbf{p}_{\mu}(s) = s^{-(\mu_1 + \dots + \mu_n)} \left(\sum_{\sigma \in S_n} \prod_{i=1}^n \mu_i x_{\sigma(i)}^{\mu_i-1} \right) dx_1 \cdots dx_n$$

as a differential form on \mathbb{C}^n .

Now from (6.7), (6.8) and (6.4), we obtain

$$(6.9) \quad \begin{aligned} d_1 \cdots d_n \mathbf{H}_{g,n}(s, \mathbf{p}(s)) &= \sum_{\mu: \ell(\mu)=n} h_{g,\mu} d_1 \cdots d_n \mathbf{p}_{\mu}(s) \frac{s^r}{r!} \\ &= \sum_{\mu: \ell(\mu)=n} h_{g,\mu} \frac{s^r}{r!} s^{-|\mu|} \sum_{\sigma \in S_n} \prod_{i=1}^n \mu_i x_{\sigma(i)}^{\mu_i-1} dx_1 \cdots dx_n \\ &= \sum_{\mu: \ell(\mu)=n} \frac{1}{|\text{Aut}(\mu)|} H_{g,n}(\mu_1, \dots, \mu_n) s^{2g-2+n} \sum_{\sigma \in S_n} \prod_{i=1}^n \mu_i x_{\sigma(i)}^{\mu_i-1} dx_1 \cdots dx_n \\ &= \sum_{\mu \in \mathbb{Z}_+^n} H_{g,n}(\mu_1, \dots, \mu_n) s^{2g-2+n} \prod_{i=1}^n \mu_i x_i^{\mu_i-1} dx_1 \cdots dx_n \\ &= s^{2g-2+n} d_1 \cdots d_n \sum_{\mu \in \mathbb{Z}_+^n} H_{g,n}(\mu_1, \dots, \mu_n) \prod_{i=1}^n x_i^{\mu_i} \end{aligned}$$

$$= s^{2g-2+n} d_1 \cdots d_n F_{g,n}^H(t_1, \dots, t_n) = s^{2g-2+n} W_{g,n}^H(t_1, \dots, t_n).$$

This formula tells us that the Eynard-Orantin differential form $W_{g,n}^H$ is the exterior derivative of $\mathbf{H}_{g,n}(s, \mathbf{p}(s))$ with the identification (6.7). Moreover, we have

$$(6.10) \quad s^{2g-2+n} F_{g,n}^H(t_1, \dots, t_n) \equiv \mathbf{H}_{g,n}(s, \mathbf{p}(s)) \pmod{\text{Ker}(d_1 \cdots d_n)}$$

as functions on \mathbb{C}^n .

Remark 6.1. Let us examine (6.10). For $n = 1$, the power sum (6.7) contains only one term and we have

$$(6.11) \quad \mathbf{H}_{g,1}(s, \mathbf{p}(s)) = s^{2g-1} \sum_{k=1}^{\infty} H_{g,1}(k) p_k(s) s^k = s^{2g-1} \sum_{k=1}^{\infty} H_{g,1}(k) x^k = s^{2g-1} F_{g,1}^H(t_1).$$

Thus (6.10) is an equality for $n = 1$. In general what happens is

$$(6.12) \quad \begin{aligned} \mathbf{H}_{g,n}(s, \mathbf{p}) &= \frac{1}{n!} s^{2g-2+n} \sum_{\vec{\mu} \in \mathbb{Z}_+^n} H_{g,n}(\vec{\mu}) \prod_{i=1}^n (x_1^{\mu_1} + x_2^{\mu_2} + \cdots + x_n^{\mu_n}) \\ &= s^{2g-2+n} \sum_{\vec{\mu} \in \mathbb{Z}_+^n} H_{g,n}(\vec{\mu}) x_1^{\mu_1} x_2^{\mu_2} \cdots x_n^{\mu_n} + (\text{terms with less than } n \text{ variables}). \end{aligned}$$

Therefore, (6.10) is never an equality for $n > 1$.

However, the principal specialization $t = t_1 = t_2 = t_3 = \cdots$ corresponds to evaluating

$$(6.13) \quad p_j = \left(\frac{x}{s}\right)^j.$$

With this identification we have again an equality

$$(6.14) \quad \begin{aligned} \mathbf{H}(s, \mathbf{p}) &= \sum_{g \geq 0, n \geq 1} \frac{1}{n!} s^{2g-2+n} \sum_{\vec{\mu} \in \mathbb{Z}_+^n} H_{g,n}(\vec{\mu}) x^{(\mu_1 + \cdots + \mu_n)} = \sum_{g \geq 0, n \geq 1} \frac{1}{n!} s^{2g-2+n} F_{g,n}^H(t, t, \dots, t). \end{aligned}$$

In Section 5 we noted that the Eynard-Orantin recursion for Hurwitz numbers is the Laplace transform of the cut-and-join equation (5.21) [26, 60]. Another consequence of the same combinatorial equation is a *heat equation* [35, 41, 79]

$$(6.15) \quad \frac{\partial}{\partial s} e^{\mathbf{H}(s, \mathbf{p})} = \frac{1}{2} \left[\sum_{i, j \geq 1} \left((i+j) p_i p_j \frac{\partial}{\partial p_{i+j}} + i j p_{i+j} \frac{\partial^2}{\partial p_i \partial p_j} \right) \right] e^{\mathbf{H}(s, \mathbf{p})},$$

with the initial condition $\mathbf{H}(0, \mathbf{p}) = p_1$. An important and fundamental fact here is that the heat equation (6.15) implies that $e^{\mathbf{H}(s, \mathbf{p})}$ is a KP τ -function for each value of s [41, 42, 67, 79]. Let us recall this fact here. We note that a solution of the heat equation is expanded by the eigenfunction of the second order operator. In our case of (6.15), the eigenfunctions of the cut-and-join operator on the right-hand side are given by the *Schur functions*.

For a partition $\mu = (\mu_1 \geq \mu_2 \geq \cdots)$ of a finite length $\ell(\mu)$, we define the **shifted power-sum function** by

$$(6.16) \quad \mathbf{p}_r[\mu] := \sum_{i=1}^{\infty} \left[\left(\mu_i - i + \frac{1}{2} \right)^r - \left(-i + \frac{1}{2} \right)^r \right].$$

This is a finite sum of $\ell(\mu)$ terms. In this paper we consider $\mathbf{p}_r[\mu]$ as a number associated with a partition μ . Then we have [35, 79]

$$(6.17) \quad \sum_{i,j \geq 1} \left((i+j)p_i p_j \frac{\partial}{\partial p_{i+j}} + ij p_{i+j} \frac{\partial^2}{\partial p_i \partial p_j} \right) s_\mu(\mathbf{p}) = \mathbf{p}_2[\mu] \cdot s_\mu(\mathbf{p}),$$

where $s_\mu(\mathbf{p})$ is the Schur function defined by

$$s_\mu(\mathbf{p}) = \sum_{|\lambda|=|\mu|} \frac{\chi_\mu(\lambda)}{z_\lambda} \mathbf{p}_\lambda,$$

$$z_\mu = \prod_{i=1}^{\ell(\mu)} m_i! i^{m_i},$$

$$m_i = \text{the number of parts in } \mu \text{ of length } i,$$

and $\chi_\mu(\lambda)$ is the value of the irreducible character of the representation μ of the symmetric group evaluated at the conjugacy class λ . If we write

$$\Delta = \frac{1}{2} \sum_{i,j \geq 1} \left((i+j)p_i p_j \frac{\partial}{\partial p_{i+j}} + ij p_{i+j} \frac{\partial^2}{\partial p_i \partial p_j} \right),$$

Then

$$\Delta s_\mu(\mathbf{p}) = \frac{1}{2} \mathbf{p}_2[\mu] \cdot s_\mu(\mathbf{p})$$

and

$$\frac{\partial}{\partial s} e^{\mathbf{H}(s,\mathbf{p})} = \Delta e^{\mathbf{H}(s,\mathbf{p})}.$$

Therefore, we have an expansion formula

$$e^{\mathbf{H}(s,\mathbf{p})} = \sum_{\mu} c_\mu s_\mu(\mathbf{p}) e^{\frac{1}{2} \mathbf{p}_2[\mu] s}$$

for a constant c_μ associated with every partition μ . The constants are determined by the initial value. Since the initial condition is

$$e^{\mathbf{H}(s,\mathbf{p})} \Big|_{s=0} = e^{p_1} = \sum_{\mu} c_\mu s_\mu(\mathbf{p}),$$

we conclude that

$$c_\mu = s_\mu(1, 0, 0, \dots, 0).$$

This follows from the Cauchy identity

$$(6.18) \quad \frac{1}{\prod_{i,j} (1 - x_i y_j)} = \sum_{\mu} s_\mu(\mathbf{x}) s_\mu(\mathbf{y}),$$

where

$$p_j = \sum_i x_i^j \quad \text{and} \quad p_j^y = \sum_i y_i^j.$$

Since we have

$$(6.19) \quad \sum_{\mu} s_\mu(\mathbf{p}) s_\mu(\mathbf{p}^y) = \frac{1}{\prod_{i,j} (1 - x_i y_j)} = \exp \left(- \sum_{i,j} \log(1 - x_i y_j) \right)$$

$$= \exp \left(\sum_{i,j} \sum_{m \geq 1} \frac{1}{m} x_j^m y_i^m \right) = \exp \left(\sum_{m \geq 1} \frac{1}{m} p_m p_m^y \right),$$

the restriction of (6.19) to $p_1^y = 1$ and $p_m^y = 0$ for all $m \geq 2$ reduces to

$$e^{p_1} = \sum_{\mu} s_{\mu}(1, 0, \dots, 0) s_{\mu}(\mathbf{p}).$$

Because of the determinantal formula for the Schur functions, $s_{\mu}(1, 0, 0, \dots, 0)$'s are the Plücker coordinate of a point of the Sato Grassmannian. It follows that

$$s_{\mu}(1, 0, 0, \dots, 0) e^{\frac{1}{2} \mathbf{p}_2[\mu]s}$$

for all μ also form the Plücker coordinates because of (6.16). Then by a theorem of Sato [73],

$$(6.20) \quad e^{\mathbf{H}(s, \mathbf{p})} = \sum_{\mu} s_{\mu}(1, 0, 0, \dots, 0) s_{\mu}(\mathbf{p}) e^{\frac{1}{2} \mathbf{p}_2[\mu]s} = \sum_{\mu} \frac{\dim \mu}{|\mu|!} e^{\frac{1}{2} \mathbf{p}_2[\mu]s} s_{\mu}(\mathbf{p})$$

is a τ -function of the KP equations. Here $\dim \mu$ is the dimension of the irreducible representation of the symmetric group $S_{|\mu|}$ belonging to the partition μ .

Thus we have established

Theorem 6.2. *The Hurwitz partition function $Z^H(t, \hbar)$ of (5.31) is obtained by evaluation of the KP τ -function $e^{\mathbf{H}(s, \mathbf{p})}$ at $(s, \mathbf{p}) = (\hbar, \mathbf{p}(\hbar))$:*

$$(6.21) \quad Z^H(t, \hbar) = e^{\mathbf{H}(\hbar, \mathbf{p}(\hbar))}.$$

Here $\mathbf{p}(\hbar)$ means the principal specialization

$$(6.22) \quad p_j = \left(\frac{x}{\hbar} \right)^j$$

for every $j = 1, 2, 3, \dots$. The t -variable and the x -variable are related by

$$x = \frac{t-1}{t} e^{\frac{1}{t}-1}.$$

Since we have a concrete expansion formula (6.20) for $e^{\mathbf{H}(s, \mathbf{p})}$, it is straightforward to find a formula for its principal specialization. Let us look at (6.19) again. This time we apply the principal specialization to both \mathbf{p} and \mathbf{p}^y , meaning that we substitute

$$p_m = x^m \quad \text{and} \quad p_m^y = y^m.$$

Then we have

$$\sum_{\mu} s_{\mu}(\mathbf{p}) s_{\mu}(\mathbf{p}^y) = \sum_{m=0}^{\infty} x^m y^m$$

after the double principal specialization. Therefore, the sum with respect to all partitions μ is reduced to the sum with respect to only one-part partitions, i.e., $\mu = (m)$. All other partitions contribute 0. Thus $s_{\mu} = s_m = h_m$, which is the m -th complete symmetric function. But because of the principal specialization, we simply have $h_m = x^m$. We note that if $\mu = (m)$, then (6.16) reduces to

$$\mathbf{p}_2[(m)] = \left(m - 1 + \frac{1}{2} \right)^2 - \left(-1 + \frac{1}{2} \right)^2 = m(m-1).$$

We therefore conclude that

$$(6.23) \quad Z^H(t, \hbar) = e^{\mathbf{H}(\hbar, \mathbf{p}(\hbar))} = \sum_{m=0}^{\infty} \frac{1}{m!} e^{\frac{1}{2}m(m-1)\hbar} \left(\frac{x}{\hbar}\right)^m.$$

We have now established a theorem of Zhou [82].

Theorem 6.3 ([82]). *The same Hurwitz partition function satisfies a differential-difference equation*

$$(6.24) \quad \left(\hbar \frac{\partial}{\partial w} + e^{-w} e^{-\hbar \frac{\partial}{\partial w}} \right) Z^H(t, \hbar) = 0.$$

Here again t is a function in w , which is given by

$$t = 1 + \sum_{k=1}^{\infty} \frac{k^k}{k!} x^k = 1 + \sum_{k=1}^{\infty} \frac{k^k}{k!} e^{-wk}.$$

The characteristic variety of this equation, with the identification of

$$z = -\hbar \frac{\partial}{\partial w},$$

is the Lagrangian immersion $e^{-w} = ze^{-z}$.

Proof. Let us denote

$$a_m = e^{\frac{1}{2}m(m-1)\hbar} \left(\frac{x}{\hbar}\right)^m = e^{(1+2+\dots+(m-1))\hbar} \left(\frac{x}{\hbar}\right)^m.$$

Then

$$Z^H(t, \hbar) = \sum_{m=0}^{\infty} \frac{1}{m!} a_m, \quad a_{m+1} = e^{m\hbar} a_m \frac{x}{\hbar}, \quad \text{and} \quad x \frac{d}{dx} a_m = m a_m.$$

We note that $x \frac{d}{dx}$ operates as the multiplication of m to a_m . Therefore,

$$\begin{aligned} -\hbar \frac{\partial}{\partial w} Z^H(t, \hbar) &= \hbar x \frac{d}{dx} \sum_{m=0}^{\infty} \frac{1}{m!} a_m = \hbar \sum_{m=0}^{\infty} \frac{1}{m!} a_{m+1} = x \sum_{m=0}^{\infty} \frac{1}{m!} e^{m\hbar} a_m \\ &= x e^{\hbar x \frac{d}{dx}} Z^H(t, \hbar) = e^{-w} e^{-\hbar \frac{\partial}{\partial w}} Z^H(t, \hbar). \end{aligned}$$

This completes the proof. \square

Remark 6.4. The asymptotic behavior of $Z^H(t, \hbar)$ near $\hbar = 0$ is determined by

$$\exp\left(\frac{1}{\hbar} S_0^H + S_1^H\right).$$

From (6.24) we have

$$\begin{aligned} 0 &= \exp\left(-\frac{1}{\hbar} S_0^H - S_1^H\right) \left(\hbar \frac{d}{dw} + e^{-w} e^{-\hbar \frac{d}{dw}}\right) \exp\left(\frac{1}{\hbar} S_0^H + S_1^H\right) \Big|_{\hbar=0} \\ &= \frac{d}{dw} S_0^H(t(w)) + e^{-w} \exp\left(\frac{S_0^H(t(w-\hbar)) - S_0^H(t(w))}{\hbar}\right) \Big|_{\hbar=0} \\ &= \frac{d}{dw} S_0^H + e^{-w} e^{-\frac{d}{dw} S_0^H} = -z + x e^z. \end{aligned}$$

We thus recover the Lagrangian immersion in this way as well.

Remark 6.5. We can directly verify that the expression (6.23) of $Z^H(t, \hbar)$ satisfies the Schrödinger equation (5.32). Indeed, since $x = e^{-w}$,

$$\begin{aligned} & \left[\frac{1}{2} \frac{\partial^2}{\partial w^2} + \left(\frac{1}{2} + \frac{1}{\hbar} \right) \frac{\partial}{\partial w} - \frac{\partial}{\partial \hbar} \right] \sum_{m=0}^{\infty} \frac{1}{m!} e^{\frac{1}{2}m(m-1)\hbar} \frac{1}{\hbar^m} e^{-mw} \\ &= \sum_{m=0}^{\infty} \frac{1}{m!} e^{\frac{1}{2}m(m-1)\hbar} \frac{1}{\hbar^m} e^{-mw} \left[\frac{1}{2}m^2 - m \left(\frac{1}{2} + \frac{1}{\hbar} \right) - \frac{1}{2}m(m-1) + \frac{m}{\hbar} \right] = 0. \end{aligned}$$

7. CONCLUSION

The main purpose of this paper is to derive the Schrödinger equation of the partition function from the integrated Eynard-Orantin topological recursion, when there is an A-model counting problem whose mirror dual is the Eynard-Orantin theory. We examined two different types of counting problem of ramified covering of \mathbb{P}^1 : one is Grothendieck's dessins d'enfants, and the other single Hurwitz numbers. The first example leads to a Lagrangian immersion in $T^*\mathbb{C}$ defined by a Laurent polynomial equation (3.6), while the latter corresponds to the Lambert curve in $T^*\mathbb{C}^*$ given by an exponential equation (5.16).

If we start with the Eynard-Orantin recursion (2.6) and define the primitive functions $F_{g,n}$ by (2.12), then we have the ambiguity in the constants of integration. However, if we start with an A-model, then the primitive functions $F_{g,n}$'s are given by the Laplace transform of the solution to the A-model problem. The examples we have studied in this paper show that always there is a natural zero $t_i = a$ of $F_{g,n}(t_1, \dots, t_n)$ in each variable, such as (4.2) and (5.13), when $2g - 2 + n > 0$. Thus the integration formula

$$F_{g,n}(t_1, \dots, t_n) = \int_a^{t_1} \cdots \int_a^{t_n} W_{g,n}(t_1, \dots, t_n)$$

uniquely determines $F_{g,n}$ from $W_{g,n}$. The integral transform equation (2.6) for $W_{g,n}$ is then equivalent to a differential transform equation for $F_{g,n}$, such as (4.3) and (5.22) for our examples. Because of our assumption for the Lagrangian immersion that the Lagrangian singularities are simply ramified, the differential recursion equation for $F_{g,n}$ is expected to be a second order PDE. Then by taking the principal specialization, we obtain a second order differential equation in t and \hbar . For our examples we have thus established (4.8) and (5.32).

Although (4.8) is holonomic if we consider \hbar a constant, (5.32) is a PDE containing the \hbar -differentiation as well. Therefore, it is not holonomic. This difference comes from the constant term $2g - 2 + n$ in the differential operator of the right-hand side of (5.22). After taking the n -fold symmetric exterior differentiation, this term drops, and thus the Eynard-Orantin recursion for Hurwitz numbers [26] takes the same shape as that of the Catalan case (3.10).

With the analysis of our examples, we notice that the issue of the constants of integration in (2.12) is *not* a simple matter. Only the corresponding A-model can dictate which constants of integration we should choose. Otherwise, the Schrödinger equation we wish to establish would take a totally different shape, depending on the choice of the constants.

Our second equation (6.24) for Hurwitz numbers is much similar to (4.8) in many ways, such as it is holonomic for each fixed \hbar . But this differential-difference equation is not a direct consequence of the differential equation (5.22), while it recovers the Lagrangian immersion more directly than (5.32). We also note here the commutator relation $[P, Q] = P$

of [49] that we mentioned in Introduction. We refer to [49] for a further integrable system theoretic analysis of these equations.

Interestingly, we derive (6.24) from the fact that there is another generating function for single Hurwitz numbers, which admits a Schur function expansion. This last point is a more general feature. In [59] we discover that there is a generalization of (6.24) for the case of double Hurwitz numbers and r -spin structures, which reduces to (6.24) as a special case. A detailed analysis of double Hurwitz numbers, or orbifold Hurwitz numbers, will be given elsewhere.

So far all these examples have a genus 0 spectral curve. Hence the proposed algebraic K-theory obstruction of Gukov and Sułkowski [39] automatically vanishes. It is also pointed out by Borot and Eynard [6] that, for a higher genus spectral curve, the definition of the partition function of the B-model needs to be modified, by including a theta function factor known as a *non-perturbative* sector. This modification also assures the modular invariance of the partition function. A further investigation is awaited here.

The examples we have carried out in this paper suggest that the expected Schrödinger equations for the knot A-polynomials should form a rather special class of the general Eynard-Orantin mechanism. This seems to be due to the integer-coefficient Laurent polynomial expression of the A-polynomials, and their K_2 Lagrangian property of Kontsevich.

APPENDIX A. PROOF OF THE SCHRÖDINGER EQUATION FOR THE CATALAN CASE

In this Appendix we give the proof of Theorem 4.5. In Section 4 we reduced the proof to verifying the recursion formula (4.16). To prove this, we need the following trivial lemma.

Lemma A.1. *Let $f(t_1, \dots, t_n)$ be a symmetric function in n variables. Then*

$$(A.1) \quad \begin{aligned} \frac{d}{dt} f(t, t, \dots, t) &= n \left[\frac{\partial}{\partial u} f(u, t, \dots, t) \right] \Big|_{u=t}; \\ \frac{d^2}{dt^2} f(t, t, \dots, t) &= n \left[\frac{\partial^2}{\partial u^2} f(u, t, \dots, t) \right] \Big|_{u=t} \\ &\quad + n(n-1) \left[\frac{\partial^2}{\partial u_1 \partial u_2} f(u_1, u_2, t, \dots, t) \right] \Big|_{u_1=u_2=t}. \end{aligned}$$

For two functions in one variable $g(x)$ and $f(x)$, we have

$$(A.2) \quad \left[\frac{1}{x-y} \left(g(x) \frac{df(x)}{dx} - g(y) \frac{df(y)}{dy} \right) \right] \Big|_{x=y} = g'(x) f'(x) + g(x) f''(x).$$

Proof. For any function f we have

$$\frac{d}{dt} f(t, t, \dots, t) = \left[\sum_{j=1}^n \frac{\partial}{\partial t_j} f(t_1, t_2, \dots, t_n) \right] \Big|_{t_1=t_2=\dots=t_n=t}.$$

Therefore,

$$\begin{aligned} \frac{d^2}{dt^2} f(t, t, \dots, t) &= \left[\left(\sum_{j=1}^n \frac{\partial}{\partial t_j} \right)^2 f(t_1, t_2, \dots, t_n) \right] \Big|_{t_1=t_2=\dots=t_n=t} \\ &= \left[\left(\sum_{j=1}^n \frac{\partial^2}{\partial t_j^2} + 2 \sum_{i \neq j} \frac{\partial^2}{\partial t_i \partial t_j} \right) f(t_1, t_2, \dots, t_n) \right] \Big|_{t_1=t_2=\dots=t_n=t}. \end{aligned}$$

Eq.(A.1) holds when f is symmetric. The second equation (A.2) follows from l'Hôpital's rule. \square

Now we are ready to give the proof of the recursion (4.16).

Proof of (4.16). The left-hand side of (4.16) is

$$\frac{d}{dt}S_{m+1} = \sum_{2g-2+n=m} \frac{1}{n!} \frac{d}{dt}F_{g,n}(t, \dots, t) = \sum_{2g-2+n=m} \frac{1}{(n-1)!} \left(\frac{\partial}{\partial t_1} F_{g,n}(t_1, t, \dots, t) \right) \Big|_{t_1=t}.$$

Thus we apply the operation

$$\sum_{2g-2+n=m} \frac{1}{(n-1)!}$$

to each line of the right-hand side of (4.3) and set all variables equal to t .

From Line 1 of the right-hand side of (4.3), we have

$$\begin{aligned} & -\frac{1}{16} \sum_{2g-2+n=m} \frac{1}{(n-1)!} \sum_{j=2}^n \frac{t_j}{(t_1+t_j)(t_1-t_j)} \\ & \quad \times \left(\frac{(t_1^2-1)^3}{t_1^2} \frac{\partial}{\partial t_1} F_{g,n-1}^C(t_{[j]}) - \frac{(t_j^2-1)^3}{t_j^2} \frac{\partial}{\partial t_j} F_{g,n-1}^C(t_{[\hat{1}]}) \right) \Big|_{t_1=\dots=t_n=t} \\ & = -\frac{1}{32} \sum_{2g-2+n=m} \frac{1}{(n-2)!} \left[\left(\frac{d}{dt} \frac{(t^2-1)^3}{t^2} \right) \frac{\partial}{\partial t_1} F_{g,n-1}^C(t_1, t, \dots, t) \right. \\ & \quad \left. + \frac{(t^2-1)^3}{t^2} \frac{\partial^2}{\partial t_1^2} F_{g,n-1}^C(t_1, t, \dots, t) \right] \Big|_{t_1=t} \\ & = -\frac{1}{32} \left(\frac{d}{dt} \frac{(t^2-1)^3}{t^2} \right) \sum_{2g-2+n=m} \frac{1}{(n-1)!} \frac{d}{dt} F_{g,n-1}^C(t, \dots, t) \\ & \quad - \frac{1}{32} \frac{(t^2-1)^3}{t^2} \sum_{2g-2+n=m} \frac{1}{(n-2)!} \frac{\partial^2}{\partial t_1^2} F_{g,n-1}^C(t_1, t, \dots, t) \Big|_{t_1=t} \\ & = -\frac{1}{32} \left(\frac{d}{dt} \frac{(t^2-1)^3}{t^2} \right) \sum_{2g'-2+n'=m-1} \frac{1}{n'!} \frac{d}{dt} F_{g',n'}^C(t, \dots, t) \\ & \quad - \frac{1}{32} \frac{(t^2-1)^3}{t^2} \sum_{2g'-2+n'=m-1} \frac{1}{(n'-1)!} \frac{\partial^2}{\partial t_1^2} F_{g',n'}^C(t_1, t, \dots, t) \Big|_{t_1=t} \\ & = -\frac{1}{32} \left(\frac{d}{dt} \frac{(t^2-1)^3}{t^2} \right) \frac{d}{dt} S_m \\ & \quad - \frac{1}{32} \frac{(t^2-1)^3}{t^2} \sum_{2g'-2+n'=m-1} \frac{1}{(n'-1)!} \frac{\partial^2}{\partial t_1^2} F_{g',n'}^C(t_1, t, \dots, t) \Big|_{t_1=t}. \end{aligned}$$

From Line 2 we obtain

$$-\frac{1}{16} \sum_{2g-2+n=m} \frac{1}{(n-1)!} \sum_{j=2}^n \frac{(t_1^2-1)^2}{t_1^2} \frac{\partial}{\partial t_1} F_{g,n-1}^C(t_{[j]}) \Big|_{t_1=\dots=t_n=t}$$

$$\begin{aligned}
&= -\frac{1}{16} \frac{(t^2 - 1)^2}{t^2} \sum_{2g-2+n=m} \frac{1}{(n-2)!} \frac{\partial}{\partial t_1} F_{g,n-1}^C(t_1, t, \dots, t) \Big|_{t_1=t} \\
&= -\frac{1}{16} \frac{(t^2 - 1)^2}{t^2} \frac{d}{dt} S_m.
\end{aligned}$$

Line 3 produces

$$\begin{aligned}
&-\frac{1}{32} \sum_{2g-2+n=m} \frac{1}{(n-1)!} \left[\frac{(t_1^2 - 1)^3}{t_1^2} \frac{\partial^2}{\partial u_1 \partial u_2} F_{g-1,n+1}^C(u_1, u_2, t_2, t_3, \dots, t_n) \right] \Big|_{u_1=u_2=t_1=\dots=t_n=t} \\
&= -\frac{1}{32} \frac{(t^2 - 1)^3}{t^2} \sum_{2g-2+n=m} \frac{1}{(n-1)!} \left[\frac{\partial^2}{\partial u_1 \partial u_2} F_{g-1,n+1}^C(u_1, u_2, t, \dots, t) \right] \Big|_{u_1=u_2=t} \\
&= -\frac{1}{32} \frac{(t^2 - 1)^3}{t^2} \sum_{2g'-2+n'=m-1, n' \geq 2} \frac{1}{(n'-2)!} \left[\frac{\partial^2}{\partial u_1 \partial u_2} F_{g',n'}^C(u_1, u_2, t, \dots, t) \right] \Big|_{u_1=u_2=t}.
\end{aligned}$$

Finally, Line 4 gives

$$\begin{aligned}
&-\frac{1}{32} \sum_{2g-2+n=m} \frac{1}{(n-1)!} \left[\frac{(t_1^2 - 1)^3}{t_1^2} \right. \\
&\quad \times \left. \sum_{\substack{\text{stable} \\ g_1+g_2=g \\ I \sqcup J = \{2,3,\dots,n\}}} \frac{\partial}{\partial t_1} F_{g_1,|I|+1}^C(t_1, t_I) \frac{\partial}{\partial t_1} F_{g_2,|J|+1}^C(t_1, t_J) \right] \Big|_{t_1=\dots=t_n=t} \\
&= -\frac{1}{32} \frac{(t^2 - 1)^3}{t^2} \sum_{2g-2+n=m} \frac{1}{(n-1)!} \\
&\quad \times \sum_{\substack{\text{stable} \\ g_1+g_2=g \\ n_1+n_2=n-1}} \left[\binom{n-1}{n_1} \frac{\partial}{\partial t_1} F_{g_1,n_1+1}^C(t_1, t, \dots, t) \frac{\partial}{\partial t_1} F_{g_2,n_2+1}^C(t_1, t, \dots, t) \right] \Big|_{t_1=t} \\
&= -\frac{1}{32} \frac{(t^2 - 1)^3}{t^2} \\
&\quad \times \sum_{2g-2+n=m} \sum_{\substack{\text{stable} \\ g_1+g_2=g \\ n_1+n_2=n-1}} \left[\frac{1}{n_1!} \frac{\partial}{\partial t_1} F_{g_1,n_1+1}^C(t_1, t, \dots, t) \frac{1}{n_2!} \frac{\partial}{\partial t_1} F_{g_2,n_2+1}^C(t_1, t, \dots, t) \right] \Big|_{t_1=t} \\
&= -\frac{1}{32} \frac{(t^2 - 1)^3}{t^2} \\
&\quad \times \sum_{2g-2+n=m} \sum_{\substack{\text{stable} \\ g_1+g_2=g \\ n_1+n_2=n-1}} \frac{1}{(n_1+1)!} \frac{d}{dt} F_{g_1,n_1+1}^C(t, \dots, t) \frac{1}{(n_2+1)!} \frac{d}{dt} F_{g_2,n_2+1}^C(t, \dots, t) \\
&= -\frac{1}{32} \frac{(t^2 - 1)^3}{t^2} \sum_{\substack{a+b=m+1 \\ a,b \geq 2}} \frac{dS_a}{dt} \frac{dS_b}{dt}.
\end{aligned}$$

In the last line we note that unstable geometries $(g, n) = (0, 1)$ and $(0, 2)$ are included only in S_0 and S_1 . This line gives the correct contribution of the product term in (4.16).

Since

$$-\frac{1}{32} \left(\frac{d}{dt} \frac{(t^2 - 1)^3}{t^2} \right) - \frac{1}{16} \frac{(t^2 - 1)^2}{t^3} = -\frac{1}{16} \frac{(t^2 - 1)^2}{t^3} (2t^2 + t + 1),$$

we have the correct term for dS_m/dt in (4.16). The remaining terms are second derivatives, and we calculate

$$\begin{aligned} & -\frac{1}{32} \frac{(t^2 - 1)^3}{t^2} \sum_{2g'-2+n'=m-1} \frac{1}{(n'-1)!} \frac{\partial^2}{\partial t_1^2} F_{g',n'}^C(t_1, t, \dots, t) \Big|_{t_1=t} \\ & -\frac{1}{32} \frac{(t^2 - 1)^3}{t^2} \sum_{2g'-2+n'=m-1, n' \geq 2} \frac{1}{(n'-2)!} \left[\frac{\partial^2}{\partial u_1 \partial u_2} F_{g',n'}^C(u_1, u_2, t, \dots, t) \right] \Big|_{u_1=u_2=t} \\ & = -\frac{1}{32} \frac{(t^2 - 1)^3}{t^2} \sum_{2g'-2+n'=m-1} \frac{1}{n'!} \\ & \times \left(n' \frac{\partial^2}{\partial t_1^2} F_{g',n'}^C(t_1, t, \dots, t) \Big|_{t_1=t} + n'(n'-1) \left[\frac{\partial^2}{\partial u_1 \partial u_2} F_{g',n'}^C(u_1, u_2, t, \dots, t) \right] \Big|_{u_1=u_2=t} \right) \\ & = -\frac{1}{32} \frac{(t^2 - 1)^3}{t^2} \frac{d^2}{dt^2} S_m. \end{aligned}$$

This completes the derivation of (4.16) from (4.3). \square

Acknowledgement. The authors thank the Banff International Research Center in Alberta and the Hausdorff Research Institute for Mathematics in Bonn for their support and hospitality, where this collaboration was started. They also thank Gaëtan Borot, Vincent Bouchard, Bertrand Eynard, Sergei Gukov, Jerry Kaminker, Maxim Kontsevich, Xiaojun Liu, Marcos Mariño, Michael Penkava, Anne Schilling, Sergey Shadrin, and Don Zagier for useful discussions. The research of M.M. has been supported by NSF DMS-1104734, DMS-1104751, Max-Planck Institut für Mathematik in Bonn, the Beijing International Center for Mathematical Research, American Institute of Mathematics in Palo Alto, the University of Salamanca, and Universiteit van Amsterdam. The research of P.S. has been supported by the DOE grant DE-FG03-92-ER40701FG-02, the European Commission under the Marie-Curie International Outgoing Fellowship Programme, and the Foundation for Polish Science.

REFERENCES

- [1] M. Aganagic, R. Dijkgraaf, A. Klemm, M. Mariño, and C. Vafa, *Topological Strings and Integrable Hierarchies*, [arXiv:hep-th/0312085], Commun. Math. Phys. **261**, 451–516 (2006).
- [2] M. Aganagic and C. Vafa, *Large N duality, mirror symmetry, and a Q-deformed A-polynomial for knots*, arXiv:1204.4709v4 [physics.hep-th] (2012).
- [3] E. Arbarello, M. Cornalba, and P.A. Griffiths, *Geometry of Algebraic Curves*, Volume II, Grundlehren der mathematischen Wissenschaften **268**, Springer-Verlag 2011.
- [4] G. V. Belyi, *On galois extensions of a maximal cyclotomic fields*, Math. U.S.S.R. Izvestija **14**, 247–256 (1980).
- [5] G. Borot and B. Eynard, *Geometry of spectral curves and all order dispersive integrable system*, arXiv:1110.4936 (2011).
- [6] G. Borot and B. Eynard, *All-order asymptotics of hyperbolic knot invariants from non-perturbative topological recursion of A-polynomials*, arXiv:1205.2261v1 [math-ph] (2012).
- [7] G. Borot, B. Eynard, M. Mulase and B. Safnuk, *Hurwitz numbers, matrix models and topological recursion*, Journal of Geometry and Physics **61**, 522–540 (2011).
- [8] V. Bouchard, A. Klemm, M. Mariño, and S. Pasquetti, *Remodeling the B-model*, Commun. Math. Phys. **287**, 117–178 (2008).
- [9] V. Bouchard and M. Mariño, *Hurwitz numbers, matrix models and enumerative geometry*, Proc. Symposia Pure Math. **78**, 263–283 (2008).

- [10] A. Brini, B. Eynard, and M. Mariño, *Torus knots and mirror symmetry* arXiv:1105.2012 (2011).
- [11] K. Chapman, M. Mulase, and B. Safnuk, *Topological recursion and the Kontsevich constants for the volume of the moduli of curves*, Communications in Number Theory and Physics **5**, 643–698 (2011).
- [12] L. Chen, *Bouchard-Klemm-Marino-Pasquetti Conjecture for \mathbb{C}^3* , arXiv:0910.3739 (2009).
- [13] D. Cooper, D.M. Culler, H. Gillet, D. Long, and P. Shalen, *Plane curves associated to character varieties of 3-manifolds*, Invent. Math. **118**, 47–84 (1994).
- [14] R.M. Corless, G.H. Gonnet, D.E.G. Hare, D.J. Jeffrey and D.E. Knuth, *On the Lambert W-function*, Adv. Computational Math. **5**, 329–359 (1996).
- [15] R. Dijkgraaf, H. Fuji, and M. Manabe, *The volume conjecture, perturbative knot invariants, and recursion relations for topological strings*, arXiv:1010.4542 [hep-th] (2010).
- [16] R. Dijkgraaf, L. Hollands, and P. Sułkowski, *Quantum curves and D-modules*, Journal of High Energy Physics **0810.4157**, 1–58 (2009).
- [17] R. Dijkgraaf, L. Hollands P. Sułkowski, and C. Vafa, *Supersymmetric gauge theories, intersecting branes and free Fermions*, Journal of High Energy Physics **0802.106**, (2008).
- [18] R. Dijkgraaf and C. Vafa, *Two Dimensional Kodaira-Spencer Theory and Three Dimensional Chern-Simons Gravity*, arXiv:0711.1932 [hep-th] (2007).
- [19] R. Dijkgraaf, E. Verlinde, and H. Verlinde, *Loop equations and Virasoro constraints in non-perturbative two-dimensional quantum gravity*, Nucl. Phys. **B348**, 435–456 (1991).
- [20] O. Dumitsrescu, M. Mulase, A. Sorkin and B. Safnuk, *The spectral curve of the Eynard-Orantin recursion via the Laplace transform*, arXiv:1202.1159 [math.AG] (2012).
- [21] P. Dunin-Barkowski, A. Mironov, A. Morozov, A. Sleptsov, and A. Smirnov, *Superpolynomials for toric knots from evolution induced by cut-and-join operators*, arXiv:1106.4305 [hep-th] (2011).
- [22] T. Ekedahl, S. Lando, M. Shapiro, A. Vainshtein, *Hurwitz numbers and intersections on moduli spaces of curves*, Invent. Math. **146**, 297–327 (2001).
- [23] B. Eynard, *Topological expansion for the 1-hermitian matrix model correlation functions*, arXiv:0407261 [hep-th] (2004).
- [24] B. Eynard, *Recursion between volumes of moduli spaces*, arXiv:0706.4403 [math-ph] (2007).
- [25] B. Eynard, *Intersection numbers of spectral curves*, arXiv:1104.0176 (2011).
- [26] B. Eynard, M. Mulase and B. Safnuk, *The Laplace transform of the cut-and-join equation and the Bouchard-Mariño conjecture on Hurwitz numbers*, Publications of the Research Institute for Mathematical Sciences **47**, 629–670 (2011).
- [27] B. Eynard and N. Orantin, *Invariants of algebraic curves and topological expansion*, Communications in Number Theory and Physics **1**, 347–452 (2007).
- [28] B. Eynard and N. Orantin, *Weil-Petersson volume of moduli spaces, Mirzakhani’s recursion and matrix models*, arXiv:0705.3600 [math-ph] (2007).
- [29] B. Eynard and N. Orantin, *Computation of open Gromov-Witten invariants for toric Calabi-Yau 3-folds by topological recursion, a proof of the BKMP conjecture*, arXiv:1205.1103v1 [math-ph] (2012).
- [30] J. Fay, *Theta functions on Riemann surfaces*, Lecture Notes in Mathematics **352**, Springer, 1973.
- [31] H. Fuji, S. Gukov, and P. Sułkowski, *Volume conjecture: refined and categorified*, arXiv:1203.2182v1 [hep-th] (2012).
- [32] H. Fuji, S. Gukov, and P. Sułkowski, *Super-A-polynomial for knots and BPS states*, arXiv:1205.1515v2 [hep-th] (2012).
- [33] S. Garoufalidis, *On the characteristic and deformation varieties of a knot*, Geometry & Topology Monographs **7**, 291–309 (2004).
- [34] S. Garoufalidis and T.T.Q. Lê, *The colored Jones function is qholonomic*, Geometry and Topology **9**, 1253–1293 (2005).
- [35] I.P. Goulden, *A differential operator for symmetric functions and the combinatorics of multiplying transpositions*, Trans. A.M.S., **344**, 421–440 (1994).
- [36] I.P. Goulden and D.M. Jackson, *Transitive factorisations into transpositions and holomorphic mappings on the sphere*, Proc. A.M.S., **125**, 51–60 (1997).
- [37] T. Graber and R. Vakil, *Hodge integrals and Hurwitz numbers via virtual localization*, Compositio Math. **135**, 25–36 (2003).
- [38] S. Gukov, *Three-dimensional quantum gravity, Chern-Simons theory, and the A-polynomial*, Commun. Math. Phys. **255**, 577–627 (2005).
- [39] S. Gukov and P. Sułkowski, *A-polynomial, B-model, and quantization*, arXiv:1108.0002v1 [hep-th] (2011).
- [40] J. L. Harer and D. Zagier, *The Euler characteristic of the moduli space of curves*, Inventiones Mathematicae **85**, 457–485 (1986).
- [41] M. Kazarian, *KP hierarchy for Hodge integrals*, arXiv:0809.3263 (2008).
- [42] M. Kazarian, S. Lando, *An algebro-geometric proof of Witten’s conjecture*, J. Amer. Math. Soc. **20**, 1079–1089 (2007).
- [43] M. Kontsevich, *Intersection theory on the moduli space of curves and the matrix Airy function*, Communications in Mathematical Physics **147**, 1–23 (1992).

- [44] M. Kontsevich, *Homological algebra of mirror symmetry*, arXiv:alg-geom/9411018 (1994).
- [45] J. Li, C.-C. M. Liu, K. Liu, and J. Zhou, *A mathematical theory of the topological vertex*, *Geom. Topol.* **13**, 527–621 (2009).
- [46] C.-C. M. Liu, *Formulae of one-partition and two-partition Hodge Integrals*, *Geom. & Top. Monographs* **8**, 105–128 (2006).
- [47] C.-C. M. Liu, *Lectures on the ELSV formula*, arXiv:1004.0853. In “Transformation Groups and Moduli Spaces of Curves,” *Adv. Lect. Math. (ALM)* **16**, 195–216, Higher Education Press and International Press, Beijing-Boston, 2010.
- [48] C.-C. M. Liu, K. Liu, J. Zhou, *A proof of a conjecture of Mariño-Vafa on Hodge Integrals*, *J. Differential Geom.* **65**, no. 2, 289–340 (2003).
- [49] X. Liu, M. Mulase, and P. Sulkowski, *Differential equations and the Eynard-Orantin recursion*, in preparation.
- [50] M. Mariño, *Chern-Simons theory, matrix models, and topological strings*, Oxford University Press, 2005.
- [51] M. Mariño, *Open string amplitudes and large order behavior in topological string theory*, *J. High Energy Physics* **0803-060**, 1–33 (2008).
- [52] M. Mirzakhani, *Simple geodesics and Weil-Petersson volumes of moduli spaces of bordered Riemann surfaces*, *Invent. Math.* **167**, 179–222 (2007).
- [53] M. Mirzakhani, *Weil-Petersson volumes and intersection theory on the moduli space of curves*, *J. Amer. Math. Soc.* **20**, 1–23 (2007).
- [54] M. Mulase, *Algebraic theory of the KP equations*, in “Perspectives in Mathematical Physics,” R. Penner and S.-T. Yau, Editors, International Press Company, 157–223 (1994).
- [55] M. Mulase, *Asymptotic analysis of a Hermitian matrix integral*, *International Journal of Mathematics* **6**, 881–892 (1995).
- [56] M. Mulase, *Lectures on the Asymptotic Expansion of a Hermitian Matrix Integral*, in Springer Lecture Notes in Physics **502**, H. Aratyn et al., Editors, 91–134 (1998).
- [57] M. Mulase and M. Penkava, *Ribbon graphs, quadratic differentials on Riemann surfaces, and algebraic curves defined over \mathbb{Q}* , *The Asian Journal of Mathematics* **2** (4), 875–920 (1998).
- [58] M. Mulase and M. Penkava, *Topological recursion for the Poincaré polynomial of the combinatorial moduli space of curves*, *Advances in Mathematics* **230**, 1322–1339 (2012).
- [59] M. Mulase, S. Shadrin, and L. Spitz, *The spectral curve and the Schrödinger equation of double Hurwitz numbers and higher spin structures*, in preparation.
- [60] M. Mulase and N. Zhang, *Polynomial recursion formula for linear Hodge integrals*, *Communications in Number Theory and Physics* **4**, 267–294 (2010).
- [61] M. Mulase and M. Zhou, in preparation.
- [62] D. Mumford, *Tata lectures on theta II, Jacobian theta functions and differential equations*, *Progress in Mathematics* **43**, Birkhäuser 1984.
- [63] P. Norbury, *Counting lattice points in the moduli space of curves*, arXiv:0801.4590 (2008).
- [64] P. Norbury, *String and dilaton equations for counting lattice points in the moduli space of curves*, arXiv:0905.4141 (2009).
- [65] P. Norbury and N. Scott, *Polynomials representing Eynard-Orantin invariants*, arXiv:1001.0449 (2010).
- [66] P. Norbury and N. Scott, *Gromov-Witten invariants of \mathbb{P}^1 and Eynard-Orantin invariants*, arXiv:1106.1337 (2011).
- [67] A. Okounkov, *Toda equations for Hurwitz numbers*, *Math. Res. Lett.* **7**, 447 (2000).
- [68] A. Okounkov and R. Pandharipande, *Gromov-Witten theory, Hurwitz numbers, and matrix models, I*, *Proc. Symposia Pure Math.* **80**, 325–414 (2009).
- [69] A. Okounkov and R. Pandharipande, *Gromov-Witten theory, Hurwitz theory, and completed cycles*, *Ann. Math.* **163**, 517–560 (2006).
- [70] A. Okounkov and R. Pandharipande, *The equivariant Gromov-Witten theory of \mathbb{P}^1* , arXiv:math/0207233 [math.AG] (2002).
- [71] H. Ooguri, P. Sulkowski, M. Yamazaki, *Wall Crossing As Seen By Matrix Models*, arXiv:1005.1293 (2010).
- [72] R. Penner, *Perturbation series and the moduli space of Riemann surfaces*, *J. Differ. Geom.* **27**, 35–53 (1988).
- [73] M. Sato, *Soliton equations as dynamical systems on an infinite dimensional Grassmannian manifold*, *Kokyuroku, Res. Inst. Math. Sci., Kyoto Univ.* **439**, 30–46 (1981).
- [74] L. Schneps and P. Lochak, *Geometric Galois actions*, *London Mathematical Society Lecture Notes Series* **242**, 1997.
- [75] G. Segal and G. Wilson, *Loop groups and equations of KdV type*, *Publ. Math. I.H.E.S.*, **61**, 5–65 (1985).
- [76] G. 't Hooft, *A planer diagram theory for strong interactions*, *Nuclear Physics* **B 72**, 461–473 (1974).
- [77] R. Vakil, *Harvard Thesis* 1997.
- [78] E. Witten, *Two dimensional gravity and intersection theory on moduli space*, *Surveys in Differential Geometry* **1**, 243–310 (1991).
- [79] J. Zhou, *Hodge integrals, Hurwitz numbers, and symmetric groups*, preprint, arXiv:math/0308024 [math.AG] (2003).
- [80] J. Zhou, *On computations of Hurwitz-Hodge integrals*, arXiv:0710.1679 (2007).

- [81] J. Zhou, *Local Mirror Symmetry for One-Legged Topological Vertex*, arXiv:0910.4320 (2009).
- [82] J. Zhou, *Quantum Mirror Curves for \mathbb{C}^3 and the Resolved Conifold*, arXiv:1207.0598v1 [math.AG] (2012).
- [83] S. Zhu, *The Laplace transform of the cut-and-join equation of Mariño-Vafa formula and its applications* arXiv:1001.0618 (2010).
- [84] D. Zvonkine, *An algebra of power series arising in the intersection theory of moduli spaces of curves and in the enumeration of ramified coverings of the sphere*, arXiv:math.AG/0403092 (2004).

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, DAVIS, CA 95616–8633, U.S.A.
E-mail address: `mulase@math.ucdavis.edu`

FACULTY OF PHYSICS, UNIVERSITY OF WARSAW, UL. HOŻA 69, 00-681 WARSAW, POLAND,

INSTITUTE FOR THEORETICAL PHYSICS, UNIVERSITY OF AMSTERDAM, SCIENCE PARK 904, 1090 GL, AMSTERDAM,
THE NETHERLANDS, AND

DEPARTMENT OF PHYSICS, CALIFORNIA INSTITUTE OF TECHNOLOGY, PASADENA, CA 91125, U.S.A.
E-mail address: `psulkows@theory.caltech.edu`