

Local extrema in random permutations and the structure of longest alternating subsequences

Dan Romik^{1†}

¹Department of Mathematics, University of California, One Shields Ave., Davis, CA 95616, USA

Abstract. Let \mathbf{as}_n denote the length of a longest alternating subsequence in a uniformly random permutation of order n . Stanley studied the distribution of \mathbf{as}_n using algebraic methods, and showed in particular that $\mathbb{E}(\mathbf{as}_n) = (4n+1)/6$ and $\text{Var}(\mathbf{as}_n) = (32n - 13)/180$. From Stanley’s result it can be shown that after rescaling, \mathbf{as}_n converges in the limit to the Gaussian distribution. In this extended abstract we present a new approach to the study of \mathbf{as}_n by relating it to the sequence of local extrema of a random permutation, which is shown to form a “canonical” longest alternating subsequence. Using this connection we reprove the abovementioned results in a more probabilistic and transparent way. We also study the distribution of the values of the local minima and maxima, and prove that in the limit the joint distribution of successive minimum-maximum pairs converges to the two-dimensional distribution whose density function is given by $f(s, t) = 3(1 - s)te^{t-s}$.

Résumé. Pour une permutation aléatoire d’ordre n , on désigne par \mathbf{as}_n la longueur maximale d’une de ses sous-suites alternantes. Stanley a étudié la distribution de \mathbf{as}_n en utilisant des méthodes algébriques, et il a démontré en particulier que $\mathbb{E}(\mathbf{as}_n) = (4n + 1)/6$ et $\text{Var}(\mathbf{as}_n) = (32n - 13)/180$. A partir du résultat de Stanley on peut montrer que \mathbf{as}_n converge à la distribution normale. Nous présentons ici une approche nouvelle à \mathbf{as}_n , par une connection à la suite de points d’extremum locaux d’une permutation aléatoire. En utilisant cette connection, nous re-prouvons les résultats mentionnés ci-dessus d’une façon plus probabiliste et transparente. En plus, nous prouvons un résultat asymptotique sur la distribution limite de paires successives de valeurs d’un minimum local suivi par un maximum local dans la permutation aléatoire.

Keywords: Longest alternating subsequences, permutation statistics, random permutation

1 Introduction

Let x_1, \dots, x_n be a sequence of distinct real numbers. A subsequence x_{i_1}, \dots, x_{i_k} , where $1 \leq i_1 < \dots < i_k \leq n$, is called an *alternating subsequence* if it satisfies

$$x_{i_1} > x_{i_2} < x_{i_3} > \dots x_{i_k}.$$

[†]Supported by the National Science Foundation under grant DMS-0955584.

Let $\text{as}_{\max}(x_1, \dots, x_n)$ be the maximal length of an alternating subsequence of x_1, \dots, x_n . Note that there may be more than one alternating subsequence of this maximal length. For example, the sequence 4, 2, 1, 3 has three longest alternating subsequences, namely (4, 1, 3), (4, 2, 3) and (2, 1, 3).

In this extended abstract we are concerned with the random variable

$$\mathbf{as}_n = \text{as}_{\max}(\sigma(1), \sigma(2), \dots, \sigma(n)),$$

where σ is a uniformly random permutation in the symmetric group S_n . Equivalently, for our purposes it will be convenient to realize \mathbf{as}_n as

$$\mathbf{as}_n = \text{as}_{\max}(X_1, X_2, \dots, X_n), \quad (1)$$

where X_1, X_2, \dots, X_n is a sequence of independent and identically distributed random variables with the uniform distribution $U[0, 1]$. The fact that this realization gives a random variable with the same distribution is a consequence of the well-known fact that the order structure of (X_1, \dots, X_n) is a uniformly random element of S_n .

Stanley (2008) studied \mathbf{as}_n , and proved, among other results, the following exact formulas for the expectation and variance of \mathbf{as}_n :

$$\begin{aligned} \mathbb{E}(\mathbf{as}_n) &= \frac{2}{3}n + \frac{1}{6}, & (n \geq 2), \\ \text{Var}(\mathbf{as}_n) &= \frac{8}{45}n - \frac{13}{180}, & (n \geq 4). \end{aligned} \quad (2) \quad (3)$$

In particular, for large values of n , \mathbf{as}_n takes values which are with high probability concentrated around its mean value of approximately $2n/3$, with a standard deviation of order \sqrt{n} . Stanley also noted that the distribution of \mathbf{as}_n converges in the limit as $n \rightarrow \infty$ to the Gaussian distribution. This was proved by Widom (2006) using a generating function identity derived by Stanley. Stanley also pointed out (see also Stanley (2010)) that the limiting Gaussian distribution follows using the same generating function identity from general results of Pemantle and Wilson (2002), and sketches a different proof based on unpublished results of Wilf (1998), which also rely on properties of specific generating functions related to \mathbf{as}_n .

In this extended abstract, we present an alternative approach to the study of the distribution of \mathbf{as}_n . The main idea is that one can construct a specific longest alternating subsequence of a given sequence x_1, \dots, x_n in a simple way using the ‘‘local extrema’’ of the sequence. In the probabilistic setting, the length \mathbf{as}_n is then equal (modulo some boundary corrections) to the number of local extrema of the sequence of random variables X_1, \dots, X_n . This number can be represented explicitly as a sum of Bernoulli random variables with simple correlations, which leads immediately to a new and more transparent derivation of the relations (2), (3) and the limiting Gaussian distribution. Furthermore, with this approach it is natural to try to understand the structure of this distinguished longest alternating subsequence formed from the local extrema; we will derive explicit formulas for the limiting densities of the local minima and maxima, and for the limiting two-dimensional density of ‘‘minimum-maximum pairs’’, which consist of a local minimum and the local maximum that follows it.

Note added in proof. Recently we were informed of a paper by Houdré and Restrepo (2010) where a similar approach to studying the longest alternating subsequences is presented and used to re-prove Stanley’s results mentioned above in a manner identical to ours. Thus, the material in Sections 2–5 below is not new, though it was discovered independently of Houdré and Restrepo’s work. The results in Sections 6 and 7 seem to be new.

2 Local extrema and the canonical alternating subsequence

Let us start with the combinatorial description of the sequence of local extrema and its relation to longest alternating subsequences. Let x_1, \dots, x_n be a sequence of distinct numbers. For $2 \leq k \leq n-1$, we say that x_k is a **local minimum** if $x_{k-1} > x_k < x_{k+1}$. We say that x_k is a **local maximum** if $x_{k-1} < x_k > x_{k+1}$, and we say that it is a **local extremum** if it is a local minimum or maximum. Define the **canonical alternating subsequence** to be the sequence of local extrema of x_1, \dots, x_n , together with the last element x_n , and together with the first element x_1 if it satisfies $x_1 > x_2$.

Lemma 1 *The canonical alternating subsequence is in fact an alternating subsequence, and its length is $\text{as}_{\max}(x_1, \dots, x_n)$.*

Proof: Because local minima and maxima appear in alternation, the sequence of local extrema meets the definition of an alternating subsequence, except possibly the requirement that the subsequence starts with a descent rather than an ascent. To make sure this requirement is also met, we add x_1 to the subsequence if the first local extremum is a minimum, which happens exactly if $x_1 > x_2$. It is also easy to check that adding x_n does not damage the alternating property in any case, so the canonical subsequence is alternating.

Denote the canonical alternating subsequence by x_{j_1}, \dots, x_{j_m} , and denote $A = \text{as}_{\max}(x_1, \dots, x_n)$. By definition we have that $m \leq A$. Conversely, to show that $m \geq A$, observe that the sequence of elements between successive local extrema is necessarily monotone. Therefore any alternating subsequence can contain at most one index from each of $[1, j_1), [j_1, j_2), \dots, [j_{m-1}, j_m)$ and $[j_m, n] = \{n\}$ and therefore has length $\leq m$. \square

3 A new probabilistic representation of as_n

As a corollary to the last result, we get the following convenient representation for the random variable as_n as defined in (1).

Corollary 2 *Define events A_1, \dots, A_{n-1} depending on the random variables X_1, \dots, X_n by*

$$\begin{aligned} A_1 &= \{X_1 > X_2\}, \\ A_k &= \{X_{k-1} < X_k > X_{k+1}\} \cup \{X_{k-1} > X_k < X_{k+1}\}, \quad (2 \leq k \leq n-1). \end{aligned}$$

Let I_A denote the indicator random variable of an event A . Then we have

$$\text{as}_n = 1 + \sum_{k=1}^{n-1} I_{A_k}. \quad (4)$$

Proof: For $2 \leq k \leq n-1$, A_k represents the event that X_k is a local extremum in the sequence X_1, \dots, X_n . So, the right-hand side of (4) exactly counts the number of terms in the canonical alternating subsequence. \square

4 Computation of the mean and variance

Next, we compute the expectations of the indicator random variables I_{A_k} , $1 \leq k \leq n-1$, and their correlations.

Lemma 3 *The means of the indicator random variables I_{A_k} , $1 \leq k \leq n-1$, are given by*

$$\mathbb{E}(I_{A_k}) = \mathbb{P}(A_k) = \begin{cases} 1/2 & k = 1, \\ 2/3 & 2 \leq k \leq n-1. \end{cases}$$

The covariances $\text{Cov}(I_{A_j}, I_{A_k}) = \mathbb{E}(I_{A_j}I_{A_k}) - \mathbb{E}(I_{A_j})\mathbb{E}(I_{A_k})$, $1 \leq j, k \leq n-1$, are given in the following covariance matrix:

$$\left(\text{Cov}(I_{A_j}, I_{A_k}) \right)_{j,k=1}^{n-1} = \begin{pmatrix} \frac{1}{4} & 0 & 0 & 0 & 0 & 0 & 0 & 0 & \dots & 0 \\ 0 & \frac{2}{9} & \frac{-1}{36} & \frac{1}{180} & 0 & 0 & 0 & 0 & \dots & 0 \\ 0 & \frac{-1}{36} & \frac{2}{9} & \frac{-1}{36} & \frac{1}{180} & 0 & 0 & 0 & \dots & 0 \\ 0 & \frac{1}{180} & \frac{-1}{36} & \frac{2}{9} & \frac{-1}{36} & \frac{1}{180} & 0 & 0 & \dots & 0 \\ 0 & 0 & \frac{1}{180} & \frac{-1}{36} & \frac{2}{9} & \frac{-1}{36} & \frac{1}{180} & 0 & \dots & 0 \\ 0 & 0 & 0 & \frac{1}{180} & \frac{-1}{36} & \frac{2}{9} & \frac{-1}{36} & \frac{1}{180} & \dots & 0 \\ \vdots & \vdots & & \ddots & \ddots & \ddots & \ddots & \ddots & & \vdots \\ 0 & 0 & \dots & & 0 & \frac{1}{180} & \frac{-1}{36} & \frac{2}{9} & \frac{-1}{36} & \frac{1}{180} \\ 0 & 0 & \dots & & & 0 & \frac{1}{180} & \frac{-1}{36} & \frac{2}{9} & \frac{-1}{36} \\ 0 & 0 & \dots & & & & 0 & \frac{1}{180} & \frac{-1}{36} & \frac{2}{9} \end{pmatrix}.$$

Proof: The computation of the means is trivial and is omitted. For the covariances, note that the events A_j and A_k are independent (and therefore the corresponding covariance is 0) if $|j-k| > 2$, since each A_j is a function of X_{j-1} , X_j and X_{j+1} only; this leaves the diagonal strip $|j-k| \leq 2$ containing potentially non-zero terms, which are computed as follows. By symmetry assume that $j \leq k$. On the main diagonal $j = k$, it is easy to compute the variances

$$\text{Cov}(I_{A_k}, I_{A_k}) = \text{Var}(I_{A_k}) = \mathbb{P}(A_k)(1 - \mathbb{P}(A_k)) = \begin{cases} 1/4 & k = 1, \\ 2/9 & 2 \leq k \leq n-1. \end{cases}$$

Next, if $j = 1$, we have for $k = 2$ that

$$\text{Cov}(I_{A_1}, I_{A_2}) = \mathbb{P}(A_1 \cap A_2) - \mathbb{P}(A_1)\mathbb{P}(A_2) = \mathbb{P}(X_1 > X_2 < X_3) - \frac{1}{2} \cdot \frac{2}{3} = \frac{1}{3} - \frac{1}{3} = 0,$$

and similarly for $k = 3$,

$$\begin{aligned} \text{Cov}(I_{A_1}, I_{A_3}) &= \mathbb{P}(A_1 \cap A_3) - \frac{1}{2} \cdot \frac{2}{3} = \mathbb{P}(X_1 > X_2 > X_3 < X_4) + \mathbb{P}(X_1 > X_2 < X_3 > X_4) - \frac{1}{3} \\ &= \frac{1}{8} + \frac{5}{24} - \frac{1}{3} = 0. \end{aligned}$$

(The number $5/24$ comes from the fact that there are 5 alternating permutations of order 4.) Next, if $k = j + 1 \geq 3$, we have

$$\begin{aligned} \text{Cov}(I_{A_j}, I_{A_{j+1}}) &= \mathbb{P}(A_2 \cap A_3) - \frac{2}{3} \cdot \frac{2}{3} \\ &= \mathbb{P}(X_1 > X_2 < X_3 > X_4) + \mathbb{P}(X_1 < X_2 > X_3 < X_4) - \frac{4}{9} = \frac{10}{24} - \frac{4}{9} = \frac{-1}{36}. \end{aligned}$$

Finally, for $k = j + 2 \geq 4$ we have

$$\begin{aligned} \text{Cov}(I_{A_j}, I_{A_{j+2}}) &= \mathbb{P}(A_2 \cap A_4) - \frac{2}{3} \cdot \frac{2}{3} \\ &= \mathbb{P}(X_1 < X_2 > X_3 < X_4 > X_5) + \mathbb{P}(X_1 > X_2 < X_3 > X_4 < X_5) \\ &\quad + \mathbb{P}(X_1 < X_2 > X_3 > X_4 < X_5) + \mathbb{P}(X_1 > X_2 < X_3 < X_4 > X_5) - \frac{4}{9} \\ &= 2 \cdot \frac{16}{120} + 2 \cdot \frac{11}{120} - \frac{4}{9} = \frac{1}{180}, \end{aligned}$$

where the fraction $16/120$ comes from the fact that there are 16 alternating permutations of order 5, and the fraction $11/120$ comes from the fact that there are 11 permutations σ of order 5 satisfying the order relations

$$\sigma(1) < \sigma(2) > \sigma(3) > \sigma(4) < \sigma(5)$$

(see also Lemma 6 below for an alternative way to compute this number). \square

Using Lemma 3 and the representation (4), we can now easily compute the mean and variance of \mathbf{as}_n to recover the relations (2), (3):

$$\begin{aligned} \mathbb{E}(\mathbf{as}_n) &= \mathbb{E} \left(1 + \sum_{k=1}^{n-1} I_{A_k} \right) = 1 + \frac{1}{2} + (n-2) \frac{2}{3} = \frac{2}{3}n + \frac{1}{6}, \\ \text{Var}(\mathbf{as}_n) &= \text{Var} \left(1 + \sum_{k=1}^{n-1} I_{A_k} \right) = \sum_{j,k=1}^n \text{Cov}(I_{A_j}, I_{A_k}) \\ &= \frac{1}{4} + (n-2) \frac{2}{9} + 2(n-3) \frac{-1}{36} + 2(n-4) \frac{1}{180} = \frac{8}{45}n - \frac{13}{180}, \end{aligned}$$

where in the computation of the mean we assume that $n \geq 2$, and for the variance computation we assume that $n \geq 4$. Note that in principle, higher moments of \mathbf{as}_n can also be computed in the same way, although the computations require higher-order correlations (e.g., of the form $\mathbb{E}(I_{A_j} I_{A_k} I_{A_\ell})$) and therefore become more tedious.

5 The limiting Gaussian distribution

We now show how the representation (4) can be used to deduce the limiting Gaussian distribution of \mathbf{as}_n .

Proposition 4 *For all $t \in \mathbb{R}$ we have*

$$\mathbb{P} \left(\frac{\mathbf{as}_n - \mathbb{E}(\mathbf{as}_n)}{(\text{Var}(\mathbf{as}_n))^{1/2}} \leq t \right) \rightarrow \frac{1}{\sqrt{2\pi}} \int_{-\infty}^t e^{-x^2/2} dx \quad \text{as } n \rightarrow \infty.$$

Proof: First, note that in (4) we can ignore the constant 1 and the first summand I_{A_1} in the sum of indicators, since their contribution is negligible compared to the scale of \sqrt{n} . Thus, if we consider the modified random variable $\mathbf{as}'_n = \sum_{k=2}^{n-1} I_{A_k}$, it will be enough to prove that \mathbf{as}'_n converges after scaling to the standard Gaussian distribution $N(0, 1)$. This sum is a sum of a sequence of identically distributed random variables. The sequence of random variables is *not* independent, but it is “3-dependent”, meaning that any two contiguous blocks of random variables that are separated by a gap of length 3 or more are independent of each other. It is a well-known fact that for bounded and identically distributed random variables (and even under much weaker conditions), one can replace the condition of independence by “ m -dependence” (where m is the size of the gap needed to ensure independence of the two blocks), and this is enough to ensure that the standard central limit theorem from probability theory holds. Formally, our sequence satisfies the assumptions of Theorem 1 of Hoeffding and Robbins (1948) (whose proof is based on a simple reduction to the standard central limit theorem for independent sequences), which gives the desired conclusion on convergence to the Gaussian distribution. \square

We emphasize that, while the proofs of the convergence to Gaussian distribution given by Stanley (2008) and Widom (2006) also rely on fairly straightforward techniques, namely asymptotic analysis of generating functions, they provide little intuition regarding why one should expect to see a Gaussian limit. The proof above makes the situation much clearer, since it is based on the observation that \mathbf{as}_n is a sum of many weakly independent components.

6 The distribution of local extrema

Having demonstrated the relevance of the sequence of local extrema to understanding the longest alternating subsequence statistic \mathbf{as}_n , it now makes sense to try to understand aspects of the behavior of the extrema, other than just how many of them there are. In particular, it is natural to look at the distribution of the extrema *values* — i.e., how small or large can we expect them to be? Of course, the extrema consist of local minima and local maxima appearing in alternation, so it makes sense to answer this question for the minima and maxima separately. The precise result is as follows.

Theorem 5 *Let N_{\min} be the (random) number of local minima in the sequence of random variables X_1, \dots, X_n , and let $m_1, m_2, \dots, m_{N_{\min}}$ denote the values of the local minima in the sequence, i.e., $m_j = X_{k_j}$ where k_j is the position where the j -th local minimum appears. For all $0 \leq a < b \leq 1$ we have the convergence in probability*

$$\frac{1}{N_{\min}} \#\{1 \leq j \leq N_{\min} : m_j \in [a, b]\} \xrightarrow[n \rightarrow \infty]{P} \int_a^b 3(1-t)^2 dt = (1-a)^3 - (1-b)^3. \quad (5)$$

In other words, the local minima in the limit are distributed in $[0, 1]$ according to the limiting probability density function $3(1-t)^2$. Similarly (and symmetrically), if we denote by N_{\max} the number of local maxima, and denote the values of the local maxima by $M_1, \dots, M_{N_{\max}}$, then for all $0 \leq a < b \leq 1$ we have the convergence in probability

$$\frac{1}{N_{\max}} \#\{1 \leq j \leq N_{\max} : M_j \in [a, b]\} \xrightarrow[n \rightarrow \infty]{P} \int_a^b 3t^2 dt = b^3 - a^3. \quad (6)$$

That is, the local maxima are in the limit distributed according to the density $3t^2$.

Proof: First, note that N_{\min} and N_{\max} differ by at most 1, and their sum is equal to $\mathbf{as}_n - 1$ or $\mathbf{as}_n - 2$ (depending on whether the term I_{A_1} in (4) is 0 or 1), so, by the results on the mean and variance of \mathbf{as}_n , we see that both N_{\min} and N_{\max} are equal to $n/3 + o(n)$ with asymptotically high probability as $n \rightarrow \infty$. Second, note that it is enough to prove (5) (which by symmetry implies (6)) in the case $b = 1$. Next, observe that we can write the number of local minima falling in the interval $[a, 1]$ as a sum of indicator random variables, namely

$$\#\{1 \leq j \leq N_{\min} : m_j \in [a, 1]\} = \sum_{k=2}^{n-1} 1_{\{X_{k-1} > X_k < X_{k+1}, X_k \geq a\}} \quad (7)$$

It is trivial to compute the mean of each of these (identically distributed) indicators. It is equal to

$$\mathbb{E} \left(1_{\{X_{k-1} > X_k < X_{k+1}, X_k \geq a\}} \right) = \mathbb{P}(X_1, X_2, X_3 \geq a, X_2 = \min(X_1, X_2, X_3)) = \frac{1}{3}(1-a)^3.$$

It is also easy to see that the variance of the sum of the indicators is $O(n)$, since, as before, each of the indicators is correlated with only the two adjacent ones on each side. Therefore we get (for example using Chebyshev's inequality) that as $n \rightarrow \infty$ the left-hand side of (7) is with high probability equal to $\frac{1}{3}(n-2)(1-a)^3 + o(n)$ (in fact, the error is of order $O(n^{1/2})$, and one can use the standard results from probability theory mentioned above to get a limiting Gaussian convergence for this random variable as well). Combining this with the previous observation about the asymptotic behavior of N_{\min} gives (5). \square

7 The joint distribution of a minimum-maximum pair

Our final result concerns a formula for the limiting *joint distribution* of a local minimum and the local maximum that follows it. We start with a lemma.

Lemma 6 *Let $d \geq 4$, and let A be an open subset of $\{(s, t) : 0 \leq s \leq t \leq 1\}$. We have*

$$\begin{aligned} & \mathbb{P}\left(X_1 > X_2 < X_3 < \dots < X_{d-1} > X_d, (X_2, X_{d-1}) \in A\right) \\ &= \frac{1}{(d-4)!} \iint_A (1-s)t(t-s)^{d-4} ds dt. \end{aligned} \quad (8)$$

Proof: The probability on the left-hand side can be expressed in an obvious way as a d -dimensional multiple integral of the constant function 1 on the subset of $[0, 1]^d$ consisting of vectors (x_1, \dots, x_d) satisfying $x_1 > x_2 < \dots < x_{d-1} > x_d$ and $(x_2, x_{d-1}) \in A$. Choosing x_2 and x_{d-1} as the outer variables of integration, we can compute this integral as the iterated integral

$$\begin{aligned} & \iint_A \left(\int_{x_2}^1 dx_1 \int_0^{x_{d-1}} dx_d \iint \dots \int_{\{x_2 \leq x_3 \leq \dots \leq x_{d-2} \leq x_{d-1}\}} dx_3 \dots dx_{d-2} \right) dx_2 dx_{d-1} \\ &= \iint_A (1-x_2)x_{d-1} \frac{(x_{d-1}-x_2)^{d-4}}{(d-4)!} dx_2 dx_{d-1}. \end{aligned}$$

This is equal to the right-hand side of (8). \square

The result on the joint distribution of a minimum-maximum pair is as follows.

Theorem 7 Let A be an open subset of $\{(s, t) : 0 \leq s \leq t \leq 1\}$. Denote by $N_{\min\text{-max}}$ the number of minimum-maximum pairs, which are defined as pairs (i, j) of positions where $i < j$, X_i is a local minimum of the sequence X_1, \dots, X_n , X_j is a local maximum, and X_i and X_j are not separated by another local extremum. Denote the values of these minimum-maximum pairs (X_i, X_j) (arranged in order of their appearance) by $(m_1, \mu_1), (m_2, \mu_2), \dots, (m_{N_{\min\text{-max}}}, \mu_{N_{\min\text{-max}}})$ (in the notation of Theorem 5, $\mu_j = M_j$ or M_{j+1} depending on whether the first local maximum appears after the first local minimum or before it). Then we have the convergence in probability

$$\frac{1}{N_{\min\text{-max}}} \#\{1 \leq k \leq N_{\min} : (m_j, \mu_j) \in A\} \xrightarrow[n \rightarrow \infty]{P} \iint_A 3(1-s)t e^{t-s} ds dt. \quad (9)$$

That is, the joint distribution of a local minimum-maximum pair is represented in the limit by the density function $f(s, t) = 3(1-s)te^{t-s}$, $(0 < s < t < 1)$.

Proof: It is easy to see that $N_{\min\text{-max}}$ differs from N_{\min} by at most 1, so as before, we know that it is with high probability approximately equal to $n/3 + o(n)$. Denote

$$T_n = \#\{1 \leq k \leq N_{\min\text{-max}} : (m_j, \mu_j) \in A\}.$$

The key observation is that we can decompose T_n based on the size of the gap between the position of the local minimum and the subsequent maximum. This leads to a representation

$$T_n = Y_{n,1} + Y_{n,2} + \dots + Y_{n,n-3},$$

where we define random variables $Y_{n,1}, \dots, Y_{n,n-3}$ by

$$Y_{n,1} = \#\{2 \leq k \leq n-2 : X_{k-1} > X_k < X_{k+1} > X_{k+2}, (X_k, X_{k+1}) \in A\},$$

$$Y_{n,2} = \#\{2 \leq k \leq n-3 : X_{k-1} > X_k < X_{k+1} < X_{k+2} > X_{k+3}, (X_k, X_{k+2}) \in A\},$$

⋮

$$Y_{n,j} = \#\{2 \leq k \leq n-j-1 : X_{k-1} > X_k < X_{k+1} < \dots < X_{k+j} > X_{k+j+1}, (X_k, X_{k+j}) \in A\},$$

⋮

$$Y_{n,n-3} = \#\{2 \leq k \leq 2 : X_1 > X_2 < X_3 < \dots < X_{n-1} > X_n, (X_2, X_{n-1}) \in A\}.$$

Now observe that each $Y_{n,j}$ can in turn be represented as a sum of $n-j-2$ indicator random variables of events of the form

$$B_{n,j,k} = \{X_{k-1} > X_k < X_{k+1} < \dots < X_{k+j} > X_{k+j+1}, (X_k, X_{k+j}) \in A\}. \quad (10)$$

For a fixed j , all of these events have the same expectation, given by the right-hand side of (8) with $d = j+3$. It follows that the expectation of T_n is given by

$$\mathbb{E}(T_n) = \sum_{j=1}^{n-3} (n-j-2) \frac{1}{(j-1)!} \iint_A (1-s)t(t-s)^{j-1} ds dt. \quad (11)$$

Because of the fast decay of the coefficients $1/(j-1)!$, it is an easy exercise to sum this series asymptotically (for example by truncating it around $j \approx \log n$; see below for a related estimate), to obtain that

$$\mathbb{E}(T_n) = n \iint_A (1-s)te^{t-s} ds dt + O(n^{-10}) \quad \text{as } n \rightarrow \infty.$$

So, we get that at least the mean of the variable $\frac{1}{N_{\min\text{-max}}}T_n$ approaches the expression on the right-hand side of (9) in the limit. It remains to show that this random variable is concentrated around its mean. To see this, set $p_n = \lfloor \log n \rfloor$ (where $\lfloor x \rfloor$ denotes the integer part of x), and define

$$T'_n = \sum_{j=1}^{p_n} Y_{n,j}.$$

Observe that $T_n = T'_n$ with high probability, since, by Markov's inequality,

$$\begin{aligned} \mathbb{P}(T_n \neq T'_n) &= \mathbb{P}\left(\sum_{j=p_n+1}^{n-3} Y_{n,j} > 0\right) \leq \mathbb{E}\left(\sum_{j=p_n+1}^{n-3} Y_{n,j}\right) \\ &\leq n \sum_{j=p_n+1}^{\infty} \frac{1}{(j-1)!} = O(n^{-10}). \end{aligned} \quad (12)$$

But we know that the variance of T'_n is given by

$$\text{Var}(T'_n) = \sum_{i,j=1}^{p_n} \text{Cov}(Y_{n,i}, Y_{n,j}).$$

We claim that these covariances satisfy $\text{Cov}(Y_{n,i}, Y_{n,j}) \leq 10n \log n$; if true, this implies that $\text{Var}(T'_n) \leq 10n(\log n)^3$, so that T'_n has standard deviation of order at most $n^{1/2}(\log n)^{3/2}$ and is therefore concentrated around its mean. To see where the covariance bound comes from, consider that each $Y_{n,j}$ is represented by a sum of indicator random variables $I_{B_{n,j,k}}$, where $B_{n,j,k}$ is defined in (10) above. The number of indicators is at most n , and furthermore the only nonzero correlations between $I_{B_{n,j,k}}$ and $I_{B_{n,i,k'}}$ can appear when $|k - k'| \leq \log n$, since otherwise the indicators are functions of independent blocks of random variables from the sequence X_1, \dots, X_n . This easily implies the stated covariance bound. To summarize, we have shown concentration of T'_n (and therefore also of T_n , by (12)) around its mean, which is also very close to the mean of T_n . Combining these facts with (11) and dividing by $n/3$, the approximate value of $N_{\min\text{-max}}$ with high probability, gives the result. \square

8 Concluding remarks

It is interesting to contrast, as Stanley (2008) did, the results for longest alternating subsequences of random permutations with the well-developed theory of longest *increasing* subsequences (see Aldous and Diaconis (1999), Stanley (2007)). In all honesty, it must be admitted that the latter subject leads to a richer and more interesting theory... Still, the study of longest alternating subsequences is not without its own rewards, and provides a nice example of the interaction of algebraic-combinatorial and probabilistic

ideas. In particular, the connection made in this extended abstract between this permutation statistic and the study of the sequence of local extrema of permutations raises interesting new questions. If one starts with a doubly-infinite sequence $\dots, X_{-1}, X_0, X_1, X_2, \dots$ of i.i.d. $U[0, 1]$ random variables and considers the local extrema (so that the restriction to any finite block says something about longest alternating subsequences in that block), the sequence of local extrema can be thought of as an interesting “stationary point process” on \mathbb{Z} . This is analogous to the sequence of descents in random sequences, which was studied by many authors. Most recently, Borodin et al. (2010) proved that the sequence of descents is a determinantal point process. One natural question that seems worthy of further study is whether this property is shared by the sequence of local extrema.

References

- D. Aldous and P. Diaconis. Longest increasing subsequences: from patience sorting to the baik-deift-johansson. *Bull. Amer. Math. Soc.*, 36:413–432, 1999.
- A. Borodin, P. Diaconis, and J. Fulman. On adding a list of numbers (and other one-dependent determinantal processes). *Bull. Amer. Math. Soc.*, 47:639–670, 2010.
- W. Hoeffding and H. Robbins. The central limit theorem for dependent random variables. *Duke Math. J.*, 15:773–780, 1948.
- C. Houdré and R. Restrepo. A probabilistic approach to the asymptotics of the length of the longest alternating subsequence. *Elec. J. Combin.*, 17:#R168, 2010.
- R. Pemantle and M. C. Wilson. Asymptotics of multivariate sequences: I. smooth points of the singular variety. *J. Combin. Theory Ser. A*, 97:129–161, 2002.
- R. P. Stanley. Increasing and decreasing subsequences and their variants. *Proc. Internat. Cong. Math. Madrid 2006*, pages 545–579, 2007.
- R. P. Stanley. Longest alternating subsequences of permutations. *Michigan Math. J.*, 57:675–687, 2008.
- R. P. Stanley. A survey of alternating permutations. *Contemp. Math.*, to appear, 2010.
- H. Widom. On the limiting distribution for the longest alternating subsequence in a random permutation. *Electron. J. Combin.*, 13:Article R25, 2006.
- H. S. Wilf. Real zeroes of polynomials that count runs and descending runs. *Unpublished preprint*, 1998.