

# MAT 207B: Methods of Applied Mathematics: Supplementary Notes I by Naoki Saito

## A Brief History of the Convergence of the Fourier Series

**Theorem 1** (Dirichlet, 1829) Suppose  $f$  is 1-periodic, piecewise smooth on  $\mathbb{R}$ . Then,  $n$ th partial sum,

$$S_n[f](x) := \sum_{-n}^n c_k e^{2\pi i k x}, \text{ satisfies}$$

$$\lim_{n \rightarrow \infty} S_n[f](x) = \frac{1}{2} [f(x+) + f(x-)].$$

In particular, if  $x$  is a point of continuity, then  $\lim_{n \rightarrow \infty} S_n[f](x) = f(x)$ .

**Theorem 2** (du Bois Reymond, 1876) There exists  $f \in C(I)$  such that  $\{S_n[f](0)\}$  diverges, where  $I$  is an interval of unit length.

**Theorem 3** (A weak version of Fejér's Theorem) If  $f$  is 1-periodic, *continuous*, and piecewise smooth on  $\mathbb{R}$ , then the Fourier series of  $f$  converges to  $f$  *absolutely* and *uniformly*.

**Definition:** Suppose a series of functions  $\sum_1^\infty g_n(x)$  converges to  $g(x)$  on a set  $x \in I$ . Then, the convergence is called *absolute* if  $\sum_1^\infty |g_n(x)|$  also converges for  $x \in I$ .

If we have  $\sup_{x \in I} \left| g(x) - \sum_1^N g_n(x) \right| \rightarrow 0$  as  $N \rightarrow \infty$ , then we call this a *uniform* convergence.

**Theorem 4** (Fejér 1904) If  $f \in C(I)$ , then the Cesàro means of  $S_n[f]$  converge *uniformly* to  $f$ .

**Definition:** The  $m$ th *Cesàro mean* of partial sums is the mean of the first  $m + 1$  partial sums, i.e.,

$$\sigma_m[f](x) := \frac{1}{m+1} \sum_{n=0}^m S_n[f](x).$$

**Theorem 5** (Size of the Fourier coefficients and the smoothness of the functions) Suppose  $f$  is 1-periodic. If  $f \in C^{k-1}(\mathbb{R})$  and  $f^{(k-1)}$  is piecewise smooth (i.e.,  $f^{(k)}$  exists and piecewise continuous), then the Fourier coefficients of  $f$ ,  $c_n$ , satisfy  $\sum_n |n^k c_n|^2 < \infty$ . In particular,  $n^k c_n \rightarrow 0$ . On the other hand, suppose  $c_n, n \neq 0$ , satisfy  $|c_n| \leq C|n|^{-(k+\gamma)}$  for some  $C > 0$  and  $\gamma > 1$ . Then  $f \in C^k(\mathbb{R})$ .

**Theorem 6** (Kolmogorov, 1926) There exists  $f \in L^1(I)$  such that  $\{S_n[f](x)\}$  diverges for every  $x$ .

**Theorem 7** (Carleson, 1966) If  $f \in L^2(I)$ , then  $S_n[f](x)$  converges to  $f(x)$  almost everywhere.

**Theorem 8** (Hunt, 1967) If  $f \in L^p(I), p > 1$ , then  $S_n[f](x)$  converges to  $f(x)$  almost everywhere.

Mathematicians are still trying to simplify the proof of the Carlson-Hunt theorem as of today.

For the details of the above facts, see [1, Chap. 1,2], [2, Chap. 1], [3, Part 1], and [4, Chap. 1]. [5, Chap. 1].

## References

- [1] J. M. ASH, ed., *Studies in Harmonic Analysis*, vol. 13 of MAA Studies in Mathematics, Math. Assoc. Amer., 1976.

- [2] H. DYM AND H. P. MCKEAN, *Fourier Series and Integrals*, Academic Press, 1972.
- [3] T. W. KÖRNER, *Fourier Analysis*, Cambridge Univ. Press, 1988.
- [4] S. G. KRANTZ, *A Panorama of Harmonic Analysis*, no. 27 in The Carus Mathematical Monographs, Math. Assoc. Amer., Washington, D.C., 1999.
- [5] M. A. PINSKY, *Introduction to Fourier Analysis and Wavelets*, Amer. Math. Soc., Providence, RI, 2002. Republished by AMS, 2009.