

MAT 271: Applied & Computational Harmonic Analysis Homework 1: due Thursday, 01/23/14

Problem 1: Prove that the dilation operator:

$$\delta_s f(x) := \frac{1}{\sqrt{s}} f\left(\frac{x}{s}\right), \quad s > 0,$$

is an isometry (i.e., norm-preserving) in $L^2(\mathbb{R})$.

Problem 2: Suppose $f, g \in L^1$. Prove the following Fourier transform formulas:

- (a) $\mathcal{F}\{\tau_a f\}(\xi) = e^{-2\pi i a \xi} \widehat{f}(\xi)$, where $a \in \mathbb{R}$.
- (b) $\mathcal{F}\{\delta_s f\}(\xi) = \delta_{1/s} \widehat{f}(\xi) = \sqrt{s} \widehat{f}(s\xi)$, where $s > 0$.
- (c) If $f \in C^1(\mathbb{R})$ and $f'(x) \rightarrow 0$ as $|x| \rightarrow \infty$, then $\mathcal{F}\{f'\}(\xi) = (2\pi i \xi) \widehat{f}(\xi)$.
- (d) $\mathcal{F}\{f * g\}(\xi) = \widehat{f}(\xi) \cdot \widehat{g}(\xi)$.

Problem 3: Compute the Fourier transform of the Gaussian function:

$$g(x; \sigma) := \frac{1}{\sqrt{2\pi\sigma}} e^{-x^2/2\sigma^2}, \quad x \in \mathbb{R}, \sigma > 0.$$

Problem 4: Prove that the following *Plancherel identities* for any functions $f, g \in \mathcal{X} := \{f \in L^1 \mid \widehat{f} \in L^1\}$:

$$\langle f, g \rangle = \langle \widehat{f}, \widehat{g} \rangle, \quad \|f\| = \|\widehat{f}\|.$$

Problem 5: Prove that the *sinc* function, $\text{sinc}(x) := \frac{\sin \pi x}{\pi x}$, $x \in \mathbb{R}$, belongs to $L^2(\mathbb{R})$, but not to $L^1(\mathbb{R})$.