

- 6 (a) Describe every vector $w = (w_1, w_2)$ that is perpendicular to $v = (2, -1)$.
 (b) All vectors perpendicular to $V = (1, 1, 1)$ lie on a _____ in 3 dimensions.
 (c) The vectors perpendicular to both $(1, 1, 1)$ and $(1, 2, 3)$ lie on a _____.
- 7 Find the angle θ (from its cosine) between these pairs of vectors:
- (a) $v = \begin{bmatrix} 1 \\ \sqrt{3} \end{bmatrix}$ and $w = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$ (b) $v = \begin{bmatrix} 2 \\ 2 \\ -1 \end{bmatrix}$ and $w = \begin{bmatrix} 2 \\ -1 \\ 2 \end{bmatrix}$
 (c) $v = \begin{bmatrix} 1 \\ \sqrt{3} \end{bmatrix}$ and $w = \begin{bmatrix} -1 \\ \sqrt{3} \end{bmatrix}$ (d) $v = \begin{bmatrix} 3 \\ 1 \end{bmatrix}$ and $w = \begin{bmatrix} -1 \\ -2 \end{bmatrix}$.
- 8 True or false (give a reason if true or find a counterexample if false):
- (a) If $u = (1, 1, 1)$ is perpendicular to v and w , then v is parallel to w .
 (b) If u is perpendicular to v and w , then u is perpendicular to $v + 2w$.
 (c) If u and v are perpendicular unit vectors then $\|u - v\| = \sqrt{2}$. *Yes!*
- 9 The slopes of the arrows from $(0, 0)$ to (v_1, v_2) and (w_1, w_2) are v_2/v_1 and w_2/w_1 . **Suppose the product v_2w_2/v_1w_1 of those slopes is -1 .** Show that $v \cdot w = 0$ and the vectors are perpendicular. (The line $y = 4x$ is perpendicular to $y = -\frac{1}{4}x$.)
- 10 Draw arrows from $(0, 0)$ to the points $v = (1, 2)$ and $w = (-2, 1)$. Multiply their slopes. That answer is a signal that $v \cdot w = 0$ and the arrows are _____.
- 11 If $v \cdot w$ is negative, what does this say about the angle between v and w ? Draw a 3-dimensional vector v (an arrow), and show where to find all w 's with $v \cdot w < 0$.
- 12 With $v = (1, 1)$ and $w = (1, 5)$ choose a number c so that $w - cv$ is perpendicular to v . Then find the formula for c starting from *any* nonzero v and w .
- 13 Find nonzero vectors v and w that are perpendicular to $(1, 0, 1)$ and to each other.
- 14 Find nonzero vectors u, v, w that are perpendicular to $(1, 1, 1, 1)$ and to each other.
- 15 The geometric mean of $x = 2$ and $y = 8$ is $\sqrt{xy} = 4$. The arithmetic mean is larger: $\frac{1}{2}(x + y) = \underline{\hspace{1cm}}$. This would come in Example 6 from the Schwarz inequality for $v = (\sqrt{2}, \sqrt{8})$ and $w = (\sqrt{8}, \sqrt{2})$. Find $\cos \theta$ for this v and w .
- 16 **How long is the vector $v = (1, 1, \dots, 1)$ in 9 dimensions?** Find a unit vector u in the same direction as v and a unit vector w that is perpendicular to v .
- 17 What are the cosines of the angles α, β, θ between the vector $(1, 0, -1)$ and the unit vectors i, j, k along the axes? Check the formula $\cos^2 \alpha + \cos^2 \beta + \cos^2 \theta = 1$.