

NO L_1 -CONTRACTIVE METRICS FOR SYSTEMS OF CONSERVATION LAWS¹

BY

BLAKE TEMPLE

ABSTRACT. Let

$$(*) \quad u_t + F(u)_x = 0$$

be any 2×2 system of conservation laws satisfying certain generic assumptions on F in a neighborhood \mathcal{N} of u -space. We prove that for every nondegenerate metric D on u -space there exists states u_1 and u_2 in \mathcal{N} such that $\int_{-\infty}^{\infty} D(u(x, t), u_1) dx$ is a strictly increasing function of t in a neighborhood of $t = 0$, where u is the admissible solution of $(*)$ with initial data

$$u(x, 0) = \begin{cases} u_1, & x \leq 0, \\ u_2, & 0 < x < 1, \\ u_1, & x \geq 1. \end{cases}$$

This contrasts with the case of a scalar equation in which $\int_{-\infty}^{\infty} D(u(x, t), v(x, t)) dx$ is a decreasing function of t for all admissible solution pairs u and v when D is taken to be the absolute value norm.

1. Introduction. Let

$$(1) \quad u_t + F(u)_x = 0, \quad u \equiv u(x, t),$$

be any 2×2 system of conservation laws which is strictly hyperbolic and genuinely nonlinear (cf. [8]) in both characteristic fields in some open neighborhood \mathcal{N}_0 of a state $u_0 \in \mathbb{R}^2$. Assume also that the shock and rarefaction curves do not coincide in at least one characteristic family, say the first (cf. [14]). (By [14] this is equivalent to assuming that each integral curve to the first eigenvector of dF is not a straight line in any open subset of \mathcal{N}_0 ; i.e., $(r \cdot \nabla)r \neq 0$ in \mathcal{N}_0 is sufficient, where r is an eigenvector field, $|r| = 1$.) We prove that for such systems there does not exist a metric D compatible with state space such that

$$(2) \quad \int_{-\infty}^{+\infty} D(u(x, t), v(x, t)) dx$$

is a decreasing function of time for all physical weak solutions u and v that agree off a compact set.

More precisely we prove the stronger statement that there does not exist a metric that is L^1 -contractive relative to a constant state for a simple class of noninteracting solutions. In particular, let dF denote the matrix derivative of F with respect to u , let

Received by the editors August 25, 1983.

1980 *Mathematics Subject Classification*. Primary 35L65, 76N10.

¹Work done while author was a visiting member at the Courant Institute.

$\lambda_1 \equiv \lambda_1(u) < \lambda_2(u) \equiv \lambda_2$ denote the eigenvalues of dF , and let

$$\Lambda = \sup\{|\lambda_2(u) - \lambda_1(v)|, u, v \in \mathcal{N}_0\}.$$

Let

$$(1') \quad u_0(\varepsilon, u_1, u_2; x) = \begin{cases} u_1 & \text{for } -\infty \leq x \leq 0, \\ u_2 & \text{for } 0 < x \leq \varepsilon, \\ u_1 & \text{for } \varepsilon < x \leq +\infty. \end{cases}$$

The Riemann problems posed in (1) and (1') can be solved and are noninteracting for times $t \leq \Lambda^{-1}\varepsilon$ (cf. [8]). Let $u(\varepsilon, u_1, u_2; x, t)$ denote this solution, $0 \leq t \leq \Lambda\varepsilon$. We prove the following theorem.

THEOREM 1. *Let u be any weak solution of the form*

$$u(x, t) = u(\varepsilon, u_1, u_2; x, t).$$

Then there does not exist a metric D compatible with u -space such that

$$(3) \quad \int_{-\infty}^{\infty} D(u(x, t), u_1) dx$$

is a decreasing function of time for all $u_1, u_2 \in \mathcal{N}_0$, $0 \leq \varepsilon \leq \Lambda$, $0 \leq t \leq \Lambda^{-1}\varepsilon$.

By a metric compatible with u -space we mean any symmetric function

$$(4) \quad D: \mathcal{N}_0 \times \mathcal{N}_0 \rightarrow \mathbf{R}$$

such that, for all u, v and w in \mathcal{N}_0 ,

$$(5) \quad (\text{triangle inequality}) \quad D(u, v) + D(v, w) \geq D(u, w)$$

and

$$(6) \quad (\text{compatibility}) \quad C_0^{-1}|u - v| \leq D(u, v) \leq C_0|u - v|$$

for some uniform constant $C_0 > 0$.

Because of the scale invariance of (1), the following corollary is a direct consequence of Theorem 1 (see the Appendix).

COROLLARY 1. *There does not exist a constant ω and a metric D compatible with u -space such that either of the following Gronwall-type inequalities holds for all weak solutions*

$$(7) \quad u(x, t) \equiv u(\varepsilon, u_1, u_2; x, t),$$

where $u_1, u_2 \in \mathcal{N}_0$, $0 \leq \varepsilon \leq \Lambda$ and $0 \leq t \leq \Lambda^{-1}\varepsilon$:

$$(8) \quad \begin{aligned} \frac{d}{dt} \int_{-\infty}^{\infty} D(u(x, t), u_1) dx &\leq \omega \int_{-\infty}^{\infty} D(u(x, t), u_1) dx, \\ \int_{-\infty}^{\infty} D(u(x, t), u_1) dx &\leq e^{\omega t} \int_{-\infty}^{\infty} D(u(x, 0), u_1) dx. \end{aligned}$$

Moreover, because the class of functions given in (7) contains only noninteracting solutions, we can also conclude from Theorem 1 that there does not exist a metric such that (2) will decrease during each iteration of the random choice method of Glimm (cf. [3]).

This result for systems contrasts with the case of a scalar conservation law for which the integral (2) is decreasing when D is taken to be the usual absolute value norm (cf. [6, 15]). In particular, Theorem 1 implies that such a result for systems will fail when D is taken to be the absolute value norm in any variables which are obtained by a regular transformation of the conserved variables.

We remark that the class of systems for which the shock and rarefaction curves do not coincide in one family includes the equations of elasticity and gas dynamics, but excludes certain equations that arise in chemical engineering (cf. [1, 4, 5, 7, 10, 13, 14]). Restricting to this class ensures that the system does not uncouple, in which case an L_1 -contractive metric does exist by results on scalar equations.

We prove Theorem 1 by contradiction, i.e., we obtain a set of inequalities by assuming that (3) decreases for certain solutions of form (7). These inequalities are then shown to be inconsistent with our other assumptions.

2. Proof of the theorem. To prove Theorem 1, assume that D is an L_1 -contractive metric for a 2×2 system (1) that satisfies our other assumptions. The four lemmas to follow are used to obtain a contradiction, and thus a proof of Theorem 1.

We adopt the following notation. Let $r_i \equiv r_i(u)$, $i = 1, 2$, denote the eigenvectors of the matrix $dF(u)$ for corresponding eigenvalues $\lambda_i(u)$, $i = 1, 2$. Let $R_i(u)$ denote the integral curve of $r_i(u)$ in \mathcal{N}_0 , and for $v \in R_i(u)$, let $R_i(u, v)$ denote the set of points on $R_i(u)$ that lie strictly between u and v . Let $S_1(u)$ denote the 1-shock curve in \mathcal{N}_0 associated with the left-hand state u . Such curves exist locally in \mathcal{N}_0 , and for $v \in S_1(u)$, $\lambda_1(v) \leq \lambda_1(u)$ (cf. [8]). Our assumption that the shock and rarefaction curves do not coincide in the first family states that $S_1(u) \not\subset R_1(u)$ for any $u \in \mathcal{N}_0$. By taking \mathcal{N}_0 sufficiently small we can, without loss of generality, assume that (cf. [14]) $\mathcal{N}_0 \cap S_1(u) \cap R_1(u) = \{u\}$.

We now let u, \bar{u}, v, \bar{v} denote any states in \mathcal{N}_0 related in the following way (see Figure 1):

$$(9) \quad \bar{v} \in S_1(u), \quad \{v\} = R_1(u) \cap R_2(\bar{v}), \quad \{\bar{u}\} = R_2(u) \cap R_1(\bar{v}),$$

and such that all states on the various shock and integral curves of r_i that connect these points pairwise also lie in \mathcal{N}_0 . In this case we say that u, \bar{u}, v, \bar{v} have the configuration of Figure 1. For example, by our assumptions, in any neighborhood of $u \in \mathcal{N}_0$ there exist states \bar{u}, v and \bar{v} such that u, \bar{u}, v, \bar{v} have the configuration of Figure 1.




FIGURE 1




FIGURE 2

If u, \bar{u}, v, \bar{v} have the configuration of Figure 1, then by strict hyperbolicity and genuine nonlinearity, λ_1 increases monotonically from v to u and from \bar{v} to \bar{u} along $R_1(v, u)$ and $R_1(\bar{v}, \bar{u})$, respectively. Without loss of generality assume further that λ_2 increases monotonically from \bar{v} to v and from \bar{u} to u along $R_2(\bar{v}, v)$ and $R_2(\bar{u}, u)$, respectively. (If λ_2 decreases on $R_2(\bar{v}, v)$ and $R_2(\bar{u}, u)$, then the following argument goes through with straightforward modifications. By our assumptions of strict hyperbolicity and genuine nonlinearity, one of these two monotonicity assumptions must occur for all u, \bar{u}, v , and \bar{v} in \mathcal{N}_0 if \mathcal{N}_0 is sufficiently small.) Let $\mu(\lambda)$ (resp. $\bar{\mu}(\lambda)$) denote the parametrization of $R_1(v, u)$ (resp. $R_1(\bar{v}, \bar{u})$) with respect to λ_1 , and let $\nu(\lambda)$ (resp. $\bar{\nu}(\lambda)$) denote the parametrization of $R_2(\bar{v}, v)$ (resp. $R_2(\bar{u}, u)$) with respect to λ_2 . Finally, let s denote the speed of the 1-shock that connects u to \bar{v} .

LEMMA 1. *The following inequalities hold for any states u, \bar{u}, v, \bar{v} which have the configuration of Figure 1:*

$$(10a) \quad \int_{\lambda_1(v)}^{\lambda_1(u)} [D(\mu(\lambda), u) + D(\mu(\lambda), v)] d\lambda - (\lambda_1(u) - \lambda_1(v)) D(u, v) \geq 0,$$

$$(10b) \quad \int_{\lambda_1(\bar{v})}^{\lambda_1(\bar{u})} [D(\bar{\mu}(\lambda), u) + D(\bar{\mu}(\lambda), \bar{v})] d\lambda - (\lambda_1(\bar{u}) - \lambda_1(\bar{v})) D(u, \bar{v}) \geq 0.$$

PROOF. By the triangle inequality,

$$(11) \quad D(\mu(\lambda), u) + D(\mu(\lambda), v) \geq D(u, v)$$

for all $\mu(\lambda) \in R_1(v, u)$. Integrating (11) from $\lambda_1(v)$ to $\lambda_1(u)$ verifies (10a). Statement (10b) is derived similarly.

LEMMA 2. *There exists a neighborhood \mathcal{N}_1 of u_0 and a constant $C_1 > 0$, $\mathcal{N}_1 \subset \mathcal{N}_0$, such that, if u, \bar{u}, v, \bar{v} are states in \mathcal{N}_1 and have the configuration of Figure 1, then*

$$(12) \quad D(\bar{v}, u) - D(v, u) \leq -C_1 |u - \bar{u}|.$$

PROOF. Consider first the following solution U defined for $0 \leq t \leq 1$:

$$(13) \quad U(x, 0) = \begin{cases} u & \text{for } x \in [0, \lambda_1(u) - s], \\ v & \text{otherwise.} \end{cases}$$

Let

$$(14) \quad I_1(t) = \int_{-\infty}^{\infty} D(u(x, t), v) dx.$$

Then

$$(15) \quad I_1(1) = \int_{\lambda_1(v)}^{\lambda_1(u)} D(\mu(\lambda), v) d\lambda + (\lambda_2(\bar{v}) - \lambda_2(\bar{u})) D(\bar{v}, v) \\ + \int_{\lambda_2(\bar{v})}^{\lambda_2(v)} D(\nu(\lambda), v) d\lambda$$




FIGURE 3

and

$$(16) \quad I_1(0) = (\lambda_1(u) - s)D(u, v).$$

But by our assumption, $I_1(1) - I_1(0) \leq 0$, so we have

$$(17) \quad \int_{\lambda_1(v)}^{\lambda_1(u)} D(\mu(\lambda), v) d\lambda \leq (\lambda_1(u) - s)D(u, v) - (\lambda_2(\bar{v}) - \cancel{\lambda_1(\bar{v})})D(\bar{v}, v).$$

We now obtain a similar estimate for a complementary solution V defined for $0 \leq t \leq 1$:

$$(18) \quad V(x, 0) = \begin{cases} v & \text{for } x \in [0, \lambda_2(v) - \lambda_1(v)], \\ u & \text{otherwise.} \end{cases}$$

Let

$$(19) \quad I_2(t) = \int_{-\infty}^{\infty} D(V(x, t), u) dx,$$

(20)

$$I_2(1) = (\lambda_2(\bar{v}) - s)D(\bar{v}, u) + \int_{\lambda_2(\bar{v})}^{\lambda_2(v)} D(v(\lambda), u) d\lambda + \int_{\lambda_1(v)}^{\lambda_1(u)} D(\mu(\lambda), u) d\lambda$$

and

$$(21) \quad I_2(0) = (\lambda_2(v) - \lambda_1(v))D(v, u) \\ = [(s - \lambda_1(v)) + (\lambda_2(\bar{v}) - s) + (\lambda_2(v) - \lambda_2(\bar{v}))]D(v, u).$$

Thus $I_2(1) - I_2(0) \leq 0$ implies

$$(22) \quad \int_{\lambda_1(v)}^{\lambda_1(u)} D(\mu(\lambda), u) d\lambda \\ \leq (s - \lambda_1(v))D(v, u) + (\lambda_2(\bar{v}) - s)[D(v, u) - D(\bar{v}, u)] \\ + \left\{ (\lambda_2(v) - \lambda_2(\bar{v}))D(v, u) - \int_{\lambda_2(\bar{v})}^{\lambda_2(v)} D(v(\lambda), u) d\lambda \right\}.$$

Adding (17) to (22) gives

$$(23) \quad \left\{ \int_{\lambda_1(v)}^{\lambda_1(u)} [D(\mu(\lambda), u) + D(\mu(\lambda), v)] d\lambda - (\lambda_1(u) - \lambda_1(v))D(u, v) \right\} \\ \leq (s - \lambda_2(\bar{v}))[D(\bar{v}, u) - D(v, u)] - (\lambda_2(\bar{v}) - \cancel{\lambda_1(\bar{v})})D(\bar{v}, v) \\ + \left\{ (\lambda_2(v) - \lambda_2(\bar{v}))D(v, u) - \int_{\lambda_2(\bar{v})}^{\lambda_2(v)} D(v(\lambda), u) d\lambda \right\}.$$

But by (10a) of Lemma 1, the left-hand side of (23) is positive, so we obtain

$$(24) \quad (\lambda_2(\bar{v}) - s)[D(\bar{v}, u) - D(v, u)] \leq -(\lambda_2(\bar{v}) - \cancel{\lambda_1(\bar{v})})D(\bar{v}, v) \\ + \left\{ (\lambda_2(v) - \lambda_2(\bar{v}))D(v, u) - \int_{\lambda_2(\bar{v})}^{\lambda_2(v)} D(v(\lambda), u) d\lambda \right\}.$$




FIGURE 5

LEMMA 3. *There exists a neighborhood \mathcal{N}_2 of u_0 and a constant $C_2 > 0$, $\mathcal{N}_2 \subset \mathcal{N}_0$, such that, if u, \bar{u}, v and \bar{v} are states in \mathcal{N}_2 and have the configuration of Figure 1, then*

$$(34) \quad D(\bar{u}, \bar{v}) - D(u, \bar{v}) \leq -C_2|\bar{u} - u|.$$

PROOF. As in the proof of Lemma 2, consider first the following solution W defined for $0 \leq t \leq 1$:

$$(35) \quad W(x, 0) = \begin{cases} \bar{v} & \text{for } x \in [0, s - \lambda_1(\bar{v})], \\ u & \text{otherwise.} \end{cases}$$

Let

$$(36) \quad I_3(t) = \int_{-\infty}^{\infty} D(W(x, t), u) dx.$$

Then

$$(37) \quad I_3(1) = \int_{\lambda_1(\bar{v})}^{\lambda_1(\bar{u})} D(\bar{\mu}(\lambda), u) d\lambda \\ + (\lambda_2(\bar{u}) - \lambda_1(\bar{u}))D(\bar{u}, u) + \int_{\lambda_2(\bar{u})}^{\lambda_2(u)} D(\bar{v}(\lambda), u) d\lambda,$$

$$(38) \quad I_3(0) = (s - \lambda_1(\bar{v}))D(\bar{v}, u).$$

By our assumption $I_3(1) - I_3(0) \leq 0$, so

$$(39) \quad \int_{\lambda_1(\bar{v})}^{\lambda_1(\bar{u})} D(\bar{\mu}(\lambda), u) d\lambda - (s - \lambda_1(\bar{v}))D(\bar{v}, u) \leq -(\lambda_2(\bar{u}) - \lambda_1(\bar{u}))D(\bar{u}, u).$$

We now obtain a similar estimate for a complementary solution X defined for $0 \leq t \leq 1$:

$$(40) \quad X(x, 0) = \begin{cases} u & \text{for } x \in [0, \lambda_2(u) - s], \\ \bar{v} & \text{otherwise.} \end{cases}$$

Let

$$(41) \quad I_4(t) = \int_{-\infty}^{\infty} D(X(x, t), \bar{v}) dx.$$

Then

$$(42) \quad I_4(1) = \int_{\lambda_1(\bar{v})}^{\lambda_1(\bar{u})} D(\bar{\mu}(\lambda), \bar{v}) d\lambda \\ + (\lambda_2(\bar{u}) - \lambda_1(\bar{u}))D(\bar{u}, \bar{v}) + \int_{\lambda_2(\bar{u})}^{\lambda_2(u)} D(\bar{v}(\lambda), \bar{v}) d\lambda,$$

$$(43) \quad I_4(0) = (\lambda_2(u) - s)D(u, \bar{v}) \\ = [(\lambda_1(\bar{u}) - s) + (\lambda_2(\bar{u}) - \lambda_1(\bar{u})) + (\lambda_2(u) - \lambda_2(\bar{u}))]D(u, \bar{v}).$$

Thus $I_4(1) - I_4(0) \leq 0$ implies

$$(44) \quad \int_{\lambda_1(\bar{v})}^{\lambda_1(\bar{u})} D(\bar{\mu}(\lambda), \bar{v}) d\lambda - (\lambda_1(\bar{u}) - s) D(u, \bar{v}) \\ \leq (\lambda_2(\bar{u}) - \lambda_1(\bar{u})) [D(u, \bar{v}) - D(\bar{u}, \bar{v})] \\ + \left\{ (\lambda_2(u) - \lambda_2(\bar{u})) D(u, \bar{v}) - \int_{\lambda_2(\bar{u})}^{\lambda_2(u)} D(\bar{v}(\lambda), \bar{v}) d\lambda \right\}.$$

Adding (39) to (44) and dividing by $\lambda_2(\bar{u}) - \lambda_1(\bar{u})$ gives

$$(45) \quad \frac{1}{\lambda_2(\bar{u}) - \lambda_1(\bar{u})} \left\{ \int_{\lambda_1(\bar{v})}^{\lambda_1(\bar{u})} D(\bar{\mu}(\lambda), u) d\lambda + \int_{\lambda_1(\bar{v})}^{\lambda_1(\bar{u})} D(\bar{\mu}(\lambda), \bar{v}) d\lambda \right. \\ \left. - (\lambda_1(\bar{u}) - \lambda_1(\bar{v})) D(u, \bar{v}) \right\} \\ \leq D(u, \bar{v}) - D(\bar{u}, \bar{v}) - D(u, \bar{u}) \\ + \frac{1}{\lambda_2(\bar{u}) - \lambda_1(\bar{u})} \left\{ (\lambda_2(u) - \lambda_2(\bar{u})) D(u, \bar{v}) - \int_{\lambda_2(\bar{u})}^{\lambda_2(u)} D(\bar{v}(\lambda), \bar{v}) d\lambda \right\}.$$

The left-hand side of (45) is positive by (10b) of Lemma 1, so we obtain

$$(46) \quad D(\bar{u}, \bar{v}) - D(u, \bar{v}) \leq -D(u, \bar{u}) + \frac{1}{\lambda_2(\bar{u}) - \lambda_1(\bar{u})} \\ \times \left\{ (\lambda_2(u) - \lambda_2(\bar{u})) D(u, \bar{v}) - \int_{\lambda_2(\bar{u})}^{\lambda_2(u)} D(\bar{v}(\lambda), \bar{v}) d\lambda \right\}.$$

But by uniform continuity, $D(\bar{v}(\lambda), \bar{v}) = D(u, \bar{v}) + \epsilon(\delta)$, so the last term in (46) can be estimated as $O(1)\epsilon(\delta)|u - \bar{u}|$ as in the proof of Lemma 2. This term is dominated by $-D(u, \bar{u})$ in (46) for δ sufficiently small, so the proof of Lemma 3 is complete.

LEMMA 4. *There exists a constant $C > 0$ such that, if u, \bar{u}, v, \bar{v} are states in $\mathcal{N} \equiv \cap_{i=0}^2 \mathcal{N}_i$ and have the configuration of Figure 1, then*

$$(47) \quad D(\bar{u}, \bar{v}) - D(v, u) \leq -C|u - \bar{u}|.$$

PROOF. This follows immediately from Lemmas 2 and 3.

PROOF OF THEOREM 1. We use Lemma 4 to obtain a contradiction. Let u_0, u_1, v_0, v_1 in \mathcal{N} have the configuration of Figure 1 in the order u, \bar{u}, v, \bar{v} . Assume that $|u_0 - v_0|$ is sufficiently small so that there is a point $\tilde{u} \in \mathcal{N} \cap R_2(u_0)$ on the same side of u_0 as u_1 , such that

$$(48) \quad |u_0 - \tilde{u}| > (C_0/C)|u_0 - v_0|$$

(see (6) for definition of the constants C, C_0). Let $\{\bar{v}\} = R_2(v_0) \cap R_1(\tilde{u})$. By choosing $|u_0 - v_0|$ sufficiently small, we can assume that the closure of the region bounded by the curves $R_1(v_0, u_0)$, $R_1(\bar{v}, \tilde{u})$, $R_2(v_0, \bar{v})$, $R_2(u_0, \tilde{u})$ lies in \mathcal{N} . Now construct the sequence of points $\{u_i, v_i\}$ in \mathcal{N} such that

$$(49) \quad \{v_i\} = S_1(u_{i-1}) \cap R_2(v_{i-1}), \\ \{u_i\} = R_1(v_i) \cap R_2(u_{i-1}).$$

Thus for each i , u_{i-1} , u_i , v_{i-1} , v_i have the configuration of Figure 1. Moreover, by our assumption that the shock and rarefaction curves diverge in the first family, we can assume (by taking $|u_0 - v_0|$ sufficiently small if necessary) that there is an integer $N > 0$ such that $u_i \in R_2(u_{i-1}, \tilde{u})$ for $i < N$, and

$$(50) \quad \tilde{u} \in R_2(u_0, u_N).$$

Now we can apply Lemma 4 as follows:

$$(51) \quad D(u_i, v_i) - D(u_{i-1}, v_{i-1}) \leq -C|u_i - u_{i-1}|,$$

so, summing from 1 to N ,

$$(52) \quad \sum_{i=1}^N D(u_i, v_i) - D(u_{i-1}, v_{i-1}) = D(u_N, v_N) - D(u_0, v_0) \\ \leq -C \sum_{i=1}^N |u_i - u_{i-1}| \leq -C \cdot \frac{C_0}{C} |u_0 - v_0|$$

or

$$(53) \quad D(u_N, v_N) \leq D(u_0, v_0) - C_0|u_0 - v_0| \\ \leq C_0|u_0 - v_0| - C_0|u_0 - v_0| \leq 0.$$

Thus $u_N = v_N$ by the compatibility assumption. But this is a contradiction, since $u_N \in R_2(u_0)$ and $v_N \in R_2(v_0)$, and $R_2(u_0) \cap R_2(v_0) \cap \mathcal{N}_0 = \emptyset$ by the assumption of strict hyperbolicity. This completes the proof of Theorem 1.

It is interesting to note that the only use of the triangle inequality is in Lemma 1. Moreover, (10a) follows without the triangle inequality by using the fact that smooth solutions can be run backwards, and thus (2) must be constant in time for smooth solutions, in order for Theorem 1 to fail. Similar methods do not seem to apply to estimate (11b).

Note also that we need not assume that D is symmetric, since if (2) decreases for \bar{D} not symmetric, then (2) decreases for $D(u, v) = \bar{D}(u, v) + \bar{D}(v, u)$, where D is symmetric.

ACKNOWLEDGEMENT. I am indebted to Ronald DiPerna for suggesting this problem to me, and for many helpful discussions and ideas. I also thank Bradly Plohr for reading the first draft.

Appendix. We show that Corollary 1 follows directly from Theorem 1. Let D be any metric compatible with u -space. Assume that u and v are weak solutions of (1) that agree off a compact set in x at $t = 0$, such that

$$(A1) \quad \int_{-\infty}^{\infty} D(u(x, T), v(x, T)) dx - \int_{-\infty}^{\infty} D(u(x, 0), v(x, 0)) dx = d > 0$$

for some $T > 0$. Let $U_c(x, t) = u(cx, ct)$, $V_c(x, t) = v(cx, ct)$ and define

$$(A2) \quad I_c(t) = \int_{-\infty}^{\infty} D(U_c(x, t), V_c(x, t)) dx.$$

Then

$$(A3) \quad I_c(t/c) = I_1(t)/c.$$

Moreover, (A1) reads

$$(A4) \quad I_1(T) - I_1(0) = d > 0,$$

and so by (A3)

$$(A5) \quad I_c(T/c) - I_c(0) = (1/c)[I_1(T) - I_1(0)] = d/c.$$

Thus

$$(A6) \quad I_c(T/c) = I_c(0) + d/c = (1/c)[I_1(0) + d].$$

Now let u be a solution of form (7), let $v(x, t) \equiv u_1$ and assume there exists $\omega > 0$ such that (8) holds for all such u . Then since $U_c(x, t)$ is also of form (7) with a rescaled value of ε , we must have

$$(A7) \quad I_c(T/c) \leq e^{\omega T/c} I_c(0)$$

for all $c > 1$, and by (A6)

$$(A8) \quad \frac{1}{c} [I_1(0) + d] \leq e^{\omega T/c} I_c(0) = e^{\omega T/c} \frac{1}{c} I_1(0)$$

or

$$(A9) \quad d \leq (e^{\omega T/c} - 1) I_1(0).$$

But for c sufficiently large, (A9) fails, thus proving the corollary.

REFERENCES

1. R. Aris and N. Amundson, *Mathematical methods in chemical engineering*, Vol. 2, Prentice-Hall, Englewood Cliffs, N. J., 1966.
2. R. Courant and K. O. Friedrichs, *Supersonic flow and shock waves*, Wiley, New York, 1948.
3. J. Glimm, *Solutions in the large for nonlinear hyperbolic systems of equations*, Comm. Pure Appl. Math. **18** (1965), 697–715.
4. E. Isaacson, *Global solution of a Riemann problem for a non-strictly hyperbolic system of conservation laws arising in enhanced oil recovery*, J. Comp. Phys. (to appear).
5. F. Helfferich and G. Klein, *Multicomponent chromatography*, Dekker, New York, 1970.
6. B. Keyfitz, *Solutions with shocks: An example of an L_1 -contractive semigroup*, Comm. Pure Appl. Math. **24** (1971), 125–132.
7. B. Keyfitz and H. Kranzer, *A system of non-strictly hyperbolic conservation laws arising in elasticity theory*, Arch. Rational Mech. Anal. **72** (1980), 219–241.
8. P. D. Lax, *Hyperbolic systems of conservation laws. II*, Comm. Pure Appl. Math. **19** (1957), 537–566.
9. ———, *Shock waves and entropy*, Contributions to Nonlinear Functional Analysis (E. H. Zarantonello, ed.), Academic Press, New York, 1971, pp. 603–634.
10. T. P. Liu and C. H. Wang, *On a hyperbolic system of conservation laws which is not strictly hyperbolic*, MRC Technical Summary Report #2184, December 29, 1980.
11. D. W. Peaceman, *Fundamentals of numerical reservoir simulation*, Elsevier North-Holland, New York, 1977.
12. H. Rhee, R. Aris and N. R. Amundson, *On the theory of multicomponent chromatography*, Philos. Trans. Roy. Soc. London Ser. A **267** (1970), 419.
13. B. Temple, *Global solution of the Cauchy problem for a class of 2×2 non-strictly hyperbolic conservation laws*, Adv. in Appl. Math. **3** (1982), 335–375.
14. ———, *Systems of conservation laws with invariant submanifolds*, Proc. Amer. Math. Soc. (to appear).
15. M. Walsh, S. Bryant, R. Schechter and L. Lake, *Precipitation and dissolution of solids attending flow through porous media*, University of Texas Preprint, 1982.

COURANT INSTITUTE OF MATHEMATICAL SCIENCES, NEW YORK UNIVERSITY, NEW YORK, NEW YORK 10012

Current address: 2510 Kendall Ave., Madison, Wisconsin 53705