

Name:

Student ID:

Section:

1	2	3	4	5	6	Total

MIDTERM EXAM
Math 167
Temple-W10

Problem 1. (20pts) True or False:

(Here A is an arbitrary $m \times n$ matrix $Col(A)$ denotes the Column Space of A , $Row(A)$ the Row Space, and $Ker(A)$ the Kernel.)

- (a) $dim \{Row(A)\} = dim \{Col(A)\}$ except when $m < n$.
- (b) If $m = n$ and $Det(A) \neq 0$, then $Ax = b$ has a unique solution for every $b \in \mathcal{R}^n$.
- (c) If $m < n$, and A has maximal rank, then we must have $dim \{Ker(A)\} = n - m$.
- (d) If $A = LDL^T$ where L is lower triangular with 1's on the diagonal and D is diagonal, then A is symmetric.
- (e) Let $E = E_{ij}(a)$ denote the matrix obtained from the $m \times m$ identity matrix by putting a in the (i, j) -entry. Then the matrix multiplication $E \cdot A$ makes sense, and the effect is to add a times the j 'th row of A to the i 'th row of A .
- (f) Let $P = P_{ij}$ denote the matrix obtained from the $m \times m$ identity matrix by interchanging the i 'th and j 'th rows. Then the matrix multiplication $P \cdot A$ makes sense, and the effect is to "Put" the i 'th row of A into the j 'th column.
- (g) In general it takes more operations to do back substitution than it does to do Gaussian elimination.
- (h) Let u and v be column vectors in \mathcal{R}^n . Then the rows of the rank-1 matrix $u \cdot v^t$ are all multiples of u .
- (i) If $m < n$, then A cannot have a right inverse.
- (j) The solution space of all x such that $Ax = b$ is always a vector space.

Problem 2. (15pts) Use elementary matrices to find matrices L and U^* , (L lower triangular with 1's on the diagonal, U^* upper triangular), such that $A = LU$, assuming

$$A = \begin{bmatrix} 1 & 2 & 0 \\ 1 & 4 & 1 \\ -2 & -6 & 2 \end{bmatrix}$$

Problem 3. (15pts) Let $A = LU$ where

$$L = \begin{bmatrix} -1 & 0 & 0 \\ 1 & 2 & 0 \\ 1 & -1 & 1 \end{bmatrix}, \quad U = \begin{bmatrix} 2 & 1 & -1 \\ 0 & 1 & -1 \\ 0 & 0 & 1 \end{bmatrix}, \quad \mathbf{b} = \begin{bmatrix} 1 \\ 2 \\ -1 \end{bmatrix}.$$

Solve $Ax = b$ for $x = (x_1, x_2, x_3)$ by the **most efficient method**.

Problem 4. (20pts) Consider the problem $Ax = 0$ where

$$A = \begin{bmatrix} 2 & 1 & 0 & 2 \\ 0 & 0 & -3 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix}, \quad x = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix}.$$

- (a) Find the *pivots* of A , and the rank of A .
- (b) Determine the *pivot variables* and the *free variables* in x .
- (c) Find matrices D and R such that $A = DR$ where D is diagonal and R is the reduced row echelon form of A .
- (d) Find a basis for $Row(A)$.
- (e) Find a basis for $Ker(A)$.

Problem 5. (15pts) Find the matrix A that represents a linear transformation $T : \mathcal{R}^3 \rightarrow \mathcal{R}^5$ in terms of the standard basis \mathbf{e}_i , (the vector with 1 in the i 'th position and zeros elsewhere), if

$$\mathbf{e}_1 \rightarrow \mathbf{e}_2 - \mathbf{e}_1$$

$$\mathbf{e}_2 \rightarrow \mathbf{e}_4 - 3\mathbf{e}_2$$

$$\mathbf{e}_3 \rightarrow -\mathbf{e}_5 + 2\mathbf{e}_3.$$

Find the dimension of the kernel of T .

Problem 6. (15pts) Let $\{v_1, \dots, v_k\}$ be a finite set of vectors in a vector space V .

(a) Complete the definition:

$$\text{Span}\{v_1, \dots, v_k\} = \{v \in V :$$

(b) Define what it means for $\{v_1, \dots, v_k\}$ to be *linearly dependent*.

(c) Prove that if $\{v_1, \dots, v_k\}$ are linearly dependent, then at least one vector can be removed from the list without changing $\text{Span}\{v_1, \dots, v_k\}$.