Name:		
Student ID#:		

Section:

Final Exam Thursday March 21, 1:00-3:00pm MAT 185A, Temple, Winter 2019

Print name and ID's clearly. Have student ID ready. Write solutions clearly and legibly. Do not write near the edge of the paper or the stapled corner. Correct answers with no supporting work will not receive full credit. No calculators, notes, books, cellphones...allowed.

Problem	Your Score	Maximum Score
1		20
2		20
3		20
4		20
5		20
6		20
7		20
8		20
9		20
10		20
Total		200

Problem #1 (20pts): Recall that $\cos z = \frac{e^{iz} + e^{-iz}}{2}$, for $z = x + iy \in C$. (a) Show that $\cos z$ reduces to $\cos x$ when $z = x \in \mathcal{R}$.

(b) Find u(x, y) and v(x, y) real so that $\cos(z) = u(x, y) + iv(x, y)$.

(c) Prove $f(z) = \cos z$ satisfies the Cauchy-Riemann equations $u_x = v_y$, $u_y = -v_x$.

Problem #2 (20pts): Let z = 3i. Find $z^{1/13}$. (That is, find all complex numbers w such that $w^{13} = z$.)

Problem #3 (20pts):

(a) Assume that f^{-1} and f are inverses of each other, and $w = f^{-1}(z)$. Prove that

$$\frac{d}{dz}f^{-1}(z) = \frac{1}{\frac{d}{dw}f(w)}.$$

(b) The *logarithm* is defined as the inverse of the exponential, so w = log(z) if and only if $z = e^w$. Use part (a) together with properties of the exponential to derive $\frac{d}{dz} \log z$.

Problem #4 (20pts): Assume f(z) = u + iv is analytic in an open set containing the closure of the ball $B_R(z_0)$, and let C_R denote the positively oriented closed curve which is its boundary. Prove that u at the center is given by its average value, i.e., prove

$$u(z_0) = \frac{1}{2\pi} \int_0^{2\pi} u(z_0 + Re^{it}) dt.$$

Problem #5 (20pts): Assume only that f is continuous, but that for any points A, B in the complex plane, $\int_C f(z)dz$ is independent of path C taking A to B. Let point A be fixed. Prove: $F(z) = \int_A^z f(z)dz$ is an anti-derivative of f. (Here $\int_A^z denotes the integral along any path from <math>A$ to z.)

Problem #6 (20pts): Recall that Cauchy's Inequality states that if f is analytic in a neighborhood of $\overline{B_R(z_0)}$, then $|f^{(k)}(z_0)| \leq \frac{k!}{R^k}M$, where M is the maximum value of f in $\overline{B_R(z_0)}$. (Here $B_R(z_0)$ denotes the open ball with center z_0 and radius R, and the bar on top denotes its closure.)

(a) Use Cauchy's Inequality to prove Liouville's Theorem, that every bounded entire function is constant.

(b) Use Liouville's Theorem to prove that every polynomial P(z) of order $n \ge 1$ has a complex root. (You may assume that every polynomial P(z) is non-constant and $\lim_{z\to\infty} P(z) = \infty$ when $n \ge 1$.)

Problem #7 (20pts): Assume f is analytic everywhere except for a singularities at $z = \pm 2i$.

(a) Recall the Taylor series $f(z) = \sum_{k=0}^{\infty} c_k (z - z_0)^k$. Assuming you can differentiate the series term by term, derive a formula for the c_k in terms of the value of f and its derivatives at $z = z_0 \neq \pm 2i$.

(b) Give the radius of convergence of the Taylor series at $z_0 \neq 2i$, and state for what radii the Taylor series converges, converges uniformly, diverges, and for which radii it may or may not converge. (No proofs required.)

(c) Recall the Laruent series $f(z) = \sum_{k=1}^{\infty} \frac{c_{-k}}{(z-z_0)^k} + \sum_{0}^{\infty} c_k (z-z_0)^k$. Determine the annulus of convergence of the Laurent series at $z_0 = 2i$. State for which annuli the Laurent series converges, and state for which annuli it converges uniformly. (No proofs required.)

Problem #8 (20pts):(a) Find the residues of the function $f(z) = \frac{1}{(z-2i)(z+i)^2}$ at z = 2i and z = -i.

(b) Find the $\int_{\mathcal{C}} f(z)dz$ where \mathcal{C} is the positively oriented circle of radius r = 3/2 centered at the origin.

Problem #9 (20pts): Evaluate $\int_{-\infty}^{+\infty} \frac{1}{2+z^2} dz$ by the method in class.

Problem #10 (20pts): Evaluate $\int_0^{2\pi} \frac{1}{2+\sin t} dt$ by the method in class.