

Name: _____

Student ID#: _____

Section: _____

Final Exam

Monday March 19, 3:30-5:30pm

MAT 21D, Temple, Winter 2018

Show your work on every problem. Correct answers with no supporting work will not receive full credit. Be organized and use notation appropriately. No calculators, notes, books, cellphones, etc. Please write legibly. Please have your student ID ready to be checked when you turn in your exam.

Problem	Your Score	Maximum Score
1		25
2		25
3		25
4		25
5		25
6		25
7		25
8		25
Total		200

Problem #1 (25pts): (a) Sketch the region of integration \mathbf{R}_{xy} and evaluate the iterated integral

$$\int_{-1}^0 \int_{x^2}^{-x} x^2 y \, dy dx. \quad (1)$$

(b) Rewrite (1) with order of integration reversed. (Do not re-evaluate).

Problem #2 (25pts): Use spherical coordinates (ρ, ϕ, θ) to find the volume of the region obtained by removing the cone $\phi \leq \pi/4$ from the sphere $x^2 + y^2 + z^2 = 9$.

Problem #3 (25pts): A cannonball is shot out of a canon at angle α to the ground. Assuming the acceleration of gravity is exactly $g = 10m/s^2$, and the muzzle velocity of the cannonball is $100m/s$, find the angle α such that the cannonball will hit the ground at a distance of exactly 500 meters.

Problem #4 (25pts): (a) Let $F = M\mathbf{i} + N\mathbf{j} + P\mathbf{k}$, let C be a smooth curve that takes A to B , and let $\vec{\mathbf{r}}(t)$ be a parameterization of C . Use Leibniz's substitution principle to show the following are equal:

$$\int_C \vec{\mathbf{F}} \cdot \vec{\mathbf{T}} \, ds = \int_C \vec{\mathbf{F}} \cdot d\vec{\mathbf{r}} = \int_C \vec{\mathbf{F}} \cdot \vec{\mathbf{v}} \, dt = \int_C Mdx + Ndy + Pdz.$$

s

(b) Assume further that $\mathbf{F} = m\mathbf{a}$, and $\vec{\mathbf{F}}$ is conservative, so $\vec{\mathbf{F}} = -\nabla P$. Derive the principle of conservation of energy

$$\left\{ \frac{1}{2}mv_B^2 - \frac{1}{2}mv_A^2 \right\} + \{P(B) - P(A)\} = 0. \quad (2)$$

(Hint: Integrate $\int_C \vec{\mathbf{F}} \cdot \vec{\mathbf{T}} ds$ two different ways.)

Problem #5 (25pts): Recall Stokes Theorem:

$$\int_{\mathcal{C}} \mathbf{F} \cdot \mathbf{T} ds = \int \int_{\mathcal{S}} \text{Curl}(\mathbf{F}) \cdot \mathbf{n} d\sigma$$

and Green's Theorem

$$\int_{\mathcal{C}} Mdx + Ndy = \int \int_{\mathcal{R}_{xy}} N_x - M_y dA.$$

(a) Use the definition of the *Curl* to derive Green's Theorem from Stokes Theorem.

(b) Let a be any given real number. Use Green's Theorem to construct a vector field $\mathbf{F}(x, y) = M(x, y)\mathbf{i} + N(x, y)\mathbf{j}$ such that

$$\int_C \vec{\mathbf{F}} \cdot \vec{\mathbf{T}} ds = aA,$$

where $A > 0$ is the area enclosed by the simple closed curve C .

(c) What are the possible values for a if $\vec{\mathbf{F}}$ is a *conservative* vector field? Explain.

Problem #6 (25pts): (a) Use Stokes Theorem to derive the meaning of $\text{Curl } \vec{\mathbf{F}}(\mathbf{x}_0) \cdot \mathbf{n}$ at a point $\mathbf{x}_0 = (x_0, y_0, z_0)$ as the circulation per area around axis \mathbf{n} at \mathbf{x}_0 .

(b) Use the Divergence Theorem to derive the meaning of the $Div\vec{\mathbf{F}}(\mathbf{x}_0)$ at a point $\mathbf{x}_0 = (x_0, y_0, z_0)$ as the Flux per volume at \mathbf{x}_0 .

Problem #7 (25pts): Let $\delta \equiv \delta(x, y, z, t)$ denote the density of a gas in motion, (you can think of air flowing around an airplane), and assume it is being transported by velocity

$$\mathbf{v} = \mathbf{v}(x, y, z, t) = M(x, y, z, t)\mathbf{i} + N(x, y, z, t)\mathbf{j} + P(x, y, z, t)\mathbf{k}.$$

Let $\vec{\mathbf{F}} = \delta\mathbf{v}$ denote the mass flux vector. Now not every density and velocity field can be a real flow, and the condition it must satisfy in order that mass be conserved, is the continuity equation

$$\delta_t + \text{Div}(\delta\mathbf{v}) = 0. \tag{3}$$

Use the Divergence Theorem, and the physical interpretation of Flux, to show that if (3) holds, then mass is conserved in every volume $V \subset \mathbf{R}^3$. (Hint, show the rate at which mass changes in V equals the rate at which mass is flowing out through the boundary, at each fixed time. Use enough English words to give an argument that makes sense.)

Problem #8 (25pts): Let S denote the two dimensional surface in the (x, y) -plane bounded by the ellipse $a^2 x^2 + b^2 y^2 = 1$. Verify Stokes Theorem $\int_C \vec{\mathbf{F}} \cdot \vec{\mathbf{T}} ds = \int \int_S \text{Curl} \vec{\mathbf{F}} \cdot \vec{\mathbf{n}} d\sigma$ for $\vec{\mathbf{F}} = y\mathbf{i} - x\mathbf{j}$ by directly evaluating both sides by explicit parametrization, and showing they are equal.

(a) $\int_C \vec{\mathbf{F}} \cdot \vec{\mathbf{T}} ds = ?$

$$(b) \int \int_S \text{Curl} \vec{F} \cdot \vec{n} d\sigma = ?$$