

Math 25  
Vogler

## Some Basics for Proofs

### Sets

Defn 1) A set  $S$  is a collection of objects.

2)  $x \in S$  means  $x$  is an element of  $S$ .

3)  $x \notin S$  means  $x$  is not an element of  $S$ .

4)  $\emptyset$  represents the empty set (i.e. one with no elements)

Note: For this class, we'll be working with sets of numbers.

Ex Here are a collection of sets we'll be dealing with.

a)  $\mathbb{N} := \{1, 2, 3, \dots\}$  are the natural numbers.

b)  $\mathbb{Z} := \{\dots, -2, -1, 0, 1, 2, \dots\}$  are the integers.

c)  $\mathbb{Q} := \{\frac{m}{n} : m, n \in \mathbb{Z} \text{ and } n \neq 0\}$  are the rational numbers.

d)  $\mathbb{R}$  is the set of all real numbers (including decimals)

e)  $\mathbb{I}$  is the set of all irrational numbers.

Notes: 1)  $\mathbb{R} = \mathbb{Q} \cup \mathbb{I}$  2)  $\mathbb{I} = \mathbb{R} - \mathbb{Q}$  3) More info. in appendix.

Quantifiers Let  $P(x)$  be a proposition involving  $x$ .

Ex  $P(x) = (|x| < 7)$

Defn 1)  $(\forall x) P(x)$  means 'for all  $x$ ,  $P(x)$ '

2)  $(\exists x) P(x)$  means 'there exists an  $x$  such that  $P(x)$ '

3)  $(\exists! x) P(x)$  means 'there exists a unique  $x$  such that  $P(x)$ '

Negation of Quantifiers  $\sim$  is the negation of a proposition.

Ex If  $P(x) = (3x = 9)$ , then  $\sim P(x) = (3x \neq 9)$

Lemma 1)  $\sim (\forall x) P(x)$  is equivalent to  $(\exists x) (\sim P(x))$

2)  $\sim (\exists x) P(x)$  is equivalent to  $(\forall x) (\sim P(x))$

Notes a) By (1), we can disprove  $(\forall x) P(x)$  by finding an  $x$  (at least one) for which  $P(x)$  is false, such an  $x$  is called a counter example.

b) We cannot prove  $(\forall x) P(x)$  by giving an example(s).

---

### Subsets

Defn Let  $A$  and  $B$  be sets. We say  $A$  is a subset of  $B$ , written  $A \subseteq B$ , if and only if (iff) every element of  $A$  is an element of  $B$ . Using symbols, we have  $(\forall x) (x \in A \Rightarrow x \in B)$  or  $\forall x \in A \Rightarrow x \in B$

Ex 1)  $\mathbb{N} \subseteq \mathbb{Z} \subseteq \mathbb{Q} \subseteq \mathbb{R}$

Defn Sets  $A$  and  $B$  are equal, written  $A = B$ , if  $A \subseteq B$  and  $B \subseteq A$

Ex 1)  $\{n: n \text{ is even}\} = \{m+1: m \text{ is odd}\}$

---

### Conditionals

1) The conditional  $P \Rightarrow Q$  means 'if  $P$  then  $Q$ '

Note: A conditional is true only when  $P$  forces  $Q$  to be true.

2) The biconditional  $P \Leftrightarrow Q$  means  $P \Rightarrow Q$  and  $Q \Rightarrow P$ .