



Section 3.1


1.



2.



3.



1.) P_1 : slope $\approx \frac{9}{9} = 1$

P_2 : slope $\approx \frac{16}{4} = 4$

2.) P_1 : slope $\approx -\frac{9}{3} = -3$

P_2 : slope ≈ 0

3.) P_1 : slope $\approx \frac{14}{6} = \frac{7}{3}$

P_2 : slope $\approx -\frac{5}{9}$

5.) $f(x) = 4 - x^2$ so

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

$$= \lim_{h \rightarrow 0} \frac{(4 - (x+h)^2) - (4 - x^2)}{h}$$

$$= \lim_{h \rightarrow 0} \frac{4 - x^2 - 2hx - h^2 - 4 + x^2}{h}$$

$$= \lim_{h \rightarrow 0} \frac{h(-2x-h)}{h} = -2x, \text{ i.e.,}$$

$f'(x) = -2x$; so slope of tangent line at $(-1, 3)$ is $m = f'(-1) = 2$ and line is given by

$$Y - 3 = 2(x - (-1)) \rightarrow Y - 3 = 2x + 2 \rightarrow$$

$$Y = 2x + 5$$

8.) $f(x) = \frac{1}{x^2}$ so

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

$$= \lim_{h \rightarrow 0} \frac{\frac{1}{(x+h)^2} - \frac{1}{x^2}}{h}$$

$$= \lim_{h \rightarrow 0} \frac{x^2 - (x+h)^2}{(x+h)^2 \cdot x^2} \cdot \frac{1}{h}$$

$$= \lim_{h \rightarrow 0} \frac{x^2 - x^2 - 2hx - h^2}{(x+h)^2 \cdot x^2 \cdot h}$$

$$= \lim_{h \rightarrow 0} \frac{x(-2x-h)}{(x+h)^2 \cdot x^2 \cdot h} = \frac{-2x}{x^2 \cdot x^2}, \text{ i.e.,}$$

$$f'(x) = -\frac{2}{x^3}; \text{ so slope of tangent line}$$

at $(-1, 1)$ is $m = f'(-1) = -\frac{2}{(-1)^3} = 2$ and line is given by

$$Y - 1 = 2(x - (-1)) \rightarrow Y - 1 = 2x + 2 \rightarrow$$

$$Y = 2x + 3$$

12.) $f(x) = x - x^2$ so

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

$$\begin{aligned}
 &= \lim_{h \rightarrow 0} \frac{(x+h) - (x+h)^2}{h} - (x-x^2) \\
 &= \lim_{h \rightarrow 0} \frac{x+h-x^2-2hx-h^2-x+x^2}{h} \\
 &= \lim_{h \rightarrow 0} \frac{x(1-2x-h)}{h} = 1-2x, \text{ i.e.,} \\
 f'(x) &= 1-2x; \text{ so slope of tangent line} \\
 \text{at } (1, -1) \text{ is } m &= f'(1) = 1-2 = -1 \text{ and} \\
 \text{line is given by } Y-(-1) &= -1(x-1) \rightarrow \\
 Y+1 &= -x+1 \rightarrow \boxed{Y=-x}.
 \end{aligned}$$

$$\begin{aligned}
 18.) \quad f(x) &= \sqrt{x+1} \text{ so} \\
 f'(x) &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} \\
 &= \lim_{h \rightarrow 0} \frac{\sqrt{x+h+1} - \sqrt{x+1}}{h} \cdot \frac{\sqrt{x+h+1} + \sqrt{x+1}}{\sqrt{x+h+1} + \sqrt{x+1}} \\
 &= \lim_{h \rightarrow 0} \frac{(x+h+1) - (x+1)}{h(\sqrt{x+h+1} + \sqrt{x+1})} \\
 &= \lim_{h \rightarrow 0} \frac{x+h+1-x-1}{h(\sqrt{x+h+1} + \sqrt{x+1})} \\
 &= \lim_{h \rightarrow 0} \frac{1}{\sqrt{(\sqrt{x+h+1} + \sqrt{x+1})}} = \frac{1}{\sqrt{x+1} + \sqrt{x+1}}, \\
 \text{i.e., } f'(x) &= \frac{1}{2\sqrt{x+1}}; \text{ so slope of} \\
 \text{tangent line at } (8, 3) \text{ is } f'(8) &= \frac{1}{2\sqrt{9}} = \frac{1}{6} \text{ and line} \\
 \text{is given by } Y-3 &= \frac{1}{6}(x-8) \rightarrow Y-3 = \frac{1}{6}x - \frac{4}{3} \rightarrow \boxed{Y = \frac{1}{6}x + \frac{5}{3}}
 \end{aligned}$$

22.) $f(x) = \frac{x-1}{x+1}$ so

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

$$= \lim_{h \rightarrow 0} \frac{\frac{(x+h)-1}{(x+h)+1} - \frac{x-1}{x+1}}{h}$$

$$= \lim_{h \rightarrow 0} \frac{(x+h-1)(x+1) - (x+h+1)(x-1)}{(x+h+1)(x+1) h} \cdot \frac{1}{h}$$

$$= \lim_{h \rightarrow 0} \frac{x^2 + hx - x + x + h - 1 - (x^2 + hx + x - x - h - 1)}{(x+h+1)(x+1) h}$$

$$= \lim_{h \rightarrow 0} \frac{x^2 + bx + h - x^2 - hx + h + x}{(x+h+1)(x+1) h}$$

$$= \lim_{h \rightarrow 0} \frac{2h}{(x+h+1)(x+1) h} = \frac{2}{(x+1)^2}, \text{ i.e.,}$$

$$f'(x) = \frac{2}{(x+1)^2}; \text{ so slope of tangent line}$$

at $x=0$ (and $y=-1$) is $m = f'(0) = 2$
and line is given by

$$y - (-1) = 2(x - 0) \rightarrow \boxed{y = 2x - 1}$$

24.) $g(x) = x^3 - 3x$ so

$$g'(x) = \lim_{h \rightarrow 0} \frac{g(x+h) - g(x)}{h}$$

$$= \lim_{h \rightarrow 0} \frac{((x+h)^3 - 3(x+h)) - (x^3 - 3x)}{h}$$

$$= \lim_{h \rightarrow 0} \frac{x^3 + 3hx^2 + 3h^2x + h^3 - 3x - 3h - x^3 + 3x}{h}$$

$$= \lim_{h \rightarrow 0} \frac{h(3x^2 + 3hx + h^2 - 3)}{h} = 3x^2 - 3, \text{ i.e.,}$$

$g'(x) = 3x^2 - 3$; horizontal tangent

(slope = 0) means $g'(x) = 0$, i.e.,

$$3x^2 - 3 = 0 \rightarrow 3(x-1)(x+1) = 0 \rightarrow$$

$$\underline{x=1, Y=2} \quad \text{or} \quad \underline{x=-1, Y=2}.$$

$$33.) f(x) = \begin{cases} x^2 \sin(\frac{1}{x}), & \text{if } x \neq 0 \\ 0, & \text{if } x = 0; \end{cases}$$

$$f'(0) = \lim_{h \rightarrow 0} \frac{f(0+h) - f(0)}{h}$$

$$= \lim_{h \rightarrow 0} \frac{f(h) - 0}{h} = \lim_{h \rightarrow 0} \frac{h^2 \sin(\frac{1}{h})}{h}$$

$$= \lim_{h \rightarrow 0} h \sin(\frac{1}{h}) \quad (-1 \leq \sin(\frac{1}{h}) \leq +1)$$

$$\rightarrow \begin{cases} -h \leq h \sin(\frac{1}{h}) \leq h, & \text{if } h > 0 \\ -h \geq h \sin(\frac{1}{h}) \geq h, & \text{if } h < 0 \end{cases}; \text{ but}$$

$$\lim_{h \rightarrow 0} -h = 0 = \lim_{h \rightarrow 0} h, \text{ so by Squeeze}$$


$$\text{Principle } \lim_{h \rightarrow 0} h \sin(\frac{1}{h}) = 0 \}$$

$$= 0, \text{ i.e., } f'(0) = 0.$$

$$34.) g(x) = \begin{cases} x \sin(\frac{1}{x}), & \text{if } x \neq 0 \\ 0, & \text{if } x = 0 \end{cases}$$

$$\begin{aligned}
 g'(0) &= \lim_{h \rightarrow 0} \frac{g(0+h) - g(0)}{h} \\
 &= \lim_{h \rightarrow 0} \frac{g(h) - 0}{h} = \lim_{h \rightarrow 0} \frac{h \sin(\frac{1}{h})}{h} \\
 &= \lim_{h \rightarrow 0} \sin\left(\frac{1}{h}\right) \quad \text{DNE (oscillation between } -1 \text{ and } 1\text{),}
 \end{aligned}$$


i.e., $g'(0)$ DNE.



$$f'(0) = \lim_{h \rightarrow 0} \frac{f(0+h) - f(0)}{h}$$

$$= \lim_{h \rightarrow 0} \frac{f(h) - 0}{h} = \lim_{h \rightarrow 0} \frac{h^{4/5}}{h} = \lim_{h \rightarrow 0} \frac{1}{h^{1/5}}$$

$$\begin{cases} \frac{1}{0^+} = +\infty, & \text{if } h > 0 \\ \frac{1}{0^-} = -\infty, & \text{if } h < 0 \end{cases}, \text{ so } f'(0) \text{ DNE (corner)}$$



$$f'(0) = \lim_{h \rightarrow 0} \frac{f(0+h) - f(0)}{h} = \lim_{h \rightarrow 0} \frac{f(h) - 0}{h}$$

$$= \lim_{h \rightarrow 0} \frac{h^{1/5}}{h} = \lim_{h \rightarrow 0} \frac{1}{h^{4/5}} = \frac{1}{0^+} = +\infty,$$

so $f'(0)$ DNE (vertical tangent line)

Section 3.2

6.) $r(s) = \sqrt{25+s}$ so

$$r'(s) = \lim_{h \rightarrow 0} \frac{r(s+h) - r(s)}{h}$$

$$= \lim_{h \rightarrow 0} \frac{\sqrt{25+h+1} - \sqrt{25+1}}{h} \cdot \frac{\sqrt{25+2h+1} + \sqrt{25+1}}{\sqrt{25+2h+1} + \sqrt{25+1}}$$

$$= \lim_{h \rightarrow 0} \frac{(25+2h+1) - (25+1)}{h \cdot (\sqrt{25+2h+1} + \sqrt{25+1})} \cdot \frac{\sqrt{25+2h+1} + \sqrt{25+1}}{\sqrt{25+2h+1} + \sqrt{25+1}}$$

$$= \lim_{h \rightarrow 0} \frac{2h}{h(\sqrt{25+2h+1} + \sqrt{25+1})} = \frac{2}{2\sqrt{25+1}}, \text{ i.e.,}$$

$$r'(s) = \frac{1}{\sqrt{25+1}}; \text{ then}$$

$$r'(0) = 1, \quad r'(1) = \frac{1}{\sqrt{3}}, \quad r'\left(\frac{1}{2}\right) = \frac{1}{\sqrt{2}}.$$

11.) $p = \frac{1}{\sqrt{q+1}}$ so

$$\frac{dp}{dq} = \lim_{h \rightarrow 0} \frac{p(q+h) - p(q)}{h}$$

$$= \lim_{h \rightarrow 0} \frac{\frac{1}{\sqrt{q+h+1}} - \frac{1}{\sqrt{q+1}}}{h}.$$

$$= \lim_{h \rightarrow 0} \frac{\frac{1}{\sqrt{q+1}} - \frac{1}{\sqrt{q+h+1}}}{\sqrt{q+h+1} \cdot \sqrt{q+1}} \cdot \frac{1}{h} \cdot \frac{\sqrt{q+1} + \sqrt{q+h+1}}{\sqrt{q+1} + \sqrt{q+h+1}}$$

$$= \lim_{h \rightarrow 0} \frac{(q+1) - (q+h+1)}{\sqrt{q+h+1} \cdot \sqrt{q+1} \cdot h (\sqrt{q+1} + \sqrt{q+h+1})}$$

$$= \lim_{h \rightarrow 0} \frac{-h}{\sqrt{q+h+1} \cdot \sqrt{q+1} \cdot h \cdot (\sqrt{q+1} + \sqrt{q+h+1})}$$

$$\begin{aligned}
 &= \frac{-1}{\sqrt{q+1} \cdot \sqrt{q+1} (\sqrt{q+1} + \sqrt{q+1})} \\
 &= \frac{-1}{(q+1) \cdot 2\sqrt{q+1}} = \frac{-1}{2(q+1)^{3/2}}, \text{ i.e., } \frac{dp}{dq} = \frac{-1}{2(q+1)^{3/2}}
 \end{aligned}$$

13.) $f(x) = x + \frac{q}{x}$ so

$$\begin{aligned}
 f'(x) &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} \\
 &= \lim_{h \rightarrow 0} \frac{\left((x+h) + \frac{q}{x+h}\right) - \left(x + \frac{q}{x}\right)}{h} \\
 &= \lim_{h \rightarrow 0} \frac{x+h + \frac{q}{x+h} - x - \frac{q}{x}}{h} \\
 &= \lim_{h \rightarrow 0} \left(\frac{h}{x} + \frac{\frac{q}{x+h} - \frac{q}{x}}{h}\right) \\
 &= \lim_{h \rightarrow 0} \left(1 + \frac{qx - q(x+h)}{(x+h)x} \cdot \frac{1}{h}\right) \\
 &= \lim_{h \rightarrow 0} \left(1 + \frac{qx - qx - qh}{(x+h)x \cdot h}\right) \\
 &= \lim_{h \rightarrow 0} \left(1 - \frac{qh}{(x+h)x \cdot h}\right) = 1 - \frac{q}{x^2}, \text{ i.e.,}
 \end{aligned}$$

$$f'(x) = 1 - \frac{q}{x^2} \text{ so } f'(-3) = 1 - \frac{q}{9} = 0$$


27.) b.)

28.) a.)

29.) d.)

30.) c.)

32.) a.)



- 44.) a.) diff. for $-2 \leq x \leq 3$
b.) cont., but not diff., for no
 x -values
c.) neither cont. nor diff. for no
 x -values

46.) a.) diff for $-2 \leq x < -1$, $-1 < x < 0$,
 $0 < x < 2$, $2 < x < 3$

- b.) cont., but not diff., for $x = -1$
c.) neither cont. nor diff. for
 $x = 0$, $x = 2$

47.) a.) diff. for $-1 \leq x < 0$, $0 < x \leq 2$
b.) cont., but not diff., for $x = 0$
c.) neither cont. nor diff. for
no x -values