## Homework 12 Solutions

11.3) (i) For  $s_n = \cos\left(\frac{n\pi}{3}\right)$ , we have

$$s_n: \frac{1}{2}, -\frac{1}{2}, -1, -\frac{1}{2}, \frac{1}{2}, 1, \frac{1}{2}, \dots$$

(a) Consider the subsequence  $\{s_{6k}\}$ , so  $s_{6k} = 1 \quad \forall k \in \mathbb{N}$ . This subsequence is clearly monotonically non-decreasing (or increasing). Moreover, it is also monotonically non-increasing (or decreasing).

(b) The set of subsequential limits is  $S = \left\{\frac{1}{2}, -\frac{1}{2}, -1, 1\right\}.$ 

(c) By Theorem 11.7, we have

$$\limsup_{n \to \infty} s_n = \sup S = 1 \text{ and } \liminf_{n \to \infty} s_n = \inf S = -1.$$

(d) This sequence does not exist (i.e. diverges) by Theorem 11.7 since S has more than one element.

(e) This sequence is bounded since its lim sup and lim inf are both finite.

(ii) For 
$$t_n = \frac{3}{4n+1}$$
, we have  
 $t_n : \frac{3}{5}, \frac{3}{9}, \frac{3}{13}, \frac{3}{17}, \frac{3}{21}, \dots$ 

(a) Consider the (trivial) subsequence  $\{t_k\}$ , so  $t_k = \frac{3}{4k+1}$   $\forall k \in \mathbb{N}$ . This subsequence is clearly monotonically non-increasing (or decreasing).

- (b) The set of subsequential limits is  $S = \{0\}$ .
- (c) By Theorem 11.7, we have

$$\limsup_{n \to \infty} s_n = \sup S = 0 \text{ and } \liminf_{n \to \infty} s_n = \inf S = 0.$$

- (d) This sequence converges to 0 by Theorem 11.7 since S has exactly one element.
- (e) This sequence is bounded since its lim sup and lim inf are both finite.

(iii) For 
$$u_n = \left(-\frac{1}{2}\right)^n$$
, we have  
 $u_n : -\frac{1}{2}, \frac{1}{4}, -\frac{1}{8}, \frac{1}{16}, -\frac{1}{32}, \dots$ 

(a) Consider the (even) subsequence  $\{t_{2k}\}$ , so  $t_{2k} = \frac{1}{2^{2k}} \quad \forall k \in \mathbb{N}$ . This subsequence is clearly monotonically non-increasing (or decreasing).

- (b) The set of subsequential limits is  $S = \{0\}$ .
- (c) By Theorem 11.7, we have

$$\limsup_{n \to \infty} s_n = \sup S = 0 \text{ and } \liminf_{n \to \infty} s_n = \inf S = 0.$$

- (d) This sequence converges to 0 by Theorem 11.7 since S has exactly one element.
- (e) This sequence is bounded since its lim sup and lim inf are both finite.
- (iv) For  $v_n = (-1)^n + \frac{1}{n}$ , we have

$$v_n: 0, \frac{3}{2}, -\frac{2}{3}, \frac{5}{4}, -\frac{4}{5}, \dots$$

(a) Consider the (odd) subsequence  $\{t_{2k+1}\}$ , so  $t_{2k+1} = -1 + \frac{1}{2k+1}$   $\forall k \in \mathbb{N}$ . This subsequence is clearly monotonically non-increasing (or decreasing).

- (b) The set of subsequential limits is  $S = \{1, -1\}$ .
- (c) By Theorem 11.7, we have

$$\limsup_{n \to \infty} s_n = \sup S = 1 \text{ and } \liminf_{n \to \infty} s_n = \inf S = -1.$$

(d) This sequence does not exist (i.e. diverges) by Theorem 11.7 since S has more than one element.

(e) This sequence is bounded since its lim sup and lim inf are both finite.

11.5) Suppose  $\{q_n\}$  is the enumeration of all the rational numbers in the interval (0, 1]. Let S denote the set of its subsequential limits.

(a) Since every number can be the limit of a subsequence of the enumeration of the rational numbers (see example 11.3), every number in the interval (0, 1] can be the limit of the subsequence of  $\{q_n\}$ , so we must have  $(0, 1] \subset S$ . Moreover, since we could easily construct a sequence in (0, 1] that converges to 0 (i.e.  $t_n = \frac{1}{n} \forall n \in \mathbb{N}$ ), 0 must be in S too by Theorem 11.8. Therefore, we conclude S = [0, 1].

(b) By Theorem 11.7, we have

$$\limsup_{n \to \infty} q_n = \sup S = 1 \text{ and } \liminf_{n \to \infty} q_n = \inf S = 0.$$

11.10) Suppose  $\{s_n\}$  is the sequence of numbers illustrated in Figure 11.2, listed in the indicated order. Let S denote the set of its subsequential limits.

(a) Notice that the set of terms in the sequence  $\{s_n\}$  is

$$E := \{s_n : n \in \mathbb{N}\} = \left\{1, \frac{1}{2}, \frac{1}{3}, \frac{1}{4}, \frac{1}{5}, \frac{1}{6}, \dots\right\}$$

Moreover, each term will appear an infinite amount of times since there are an infinite amount of rows in Figure 11.2. So for each each element in E, we could construct a subsequence  $\{s_{n_k}\}$  that is identically one of these elements (i.e.  $s_{n_k} = \frac{1}{2} \quad \forall k \in \mathbb{N}$ ). Also, we can construct a subsequence  $\{s_{n_k}\}$  with every value in E (i.e.  $s_{n_k} = \frac{1}{k} \quad \forall k \in \mathbb{N}$ ), and this subsequence clearly converges to 0. Therefore, we conclude

$$S = E \cup \{0\} = \left\{0, 1, \frac{1}{2}, \frac{1}{3}, \frac{1}{4}, \frac{1}{5}, \frac{1}{6}, \dots\right\}.$$

(b) By Theorem 11.7, we have

$$\limsup_{n \to \infty} s_n = \sup S = 1 \text{ and } \liminf_{n \to \infty} s_n = \inf S = 0.$$